

NEWTON POLYTOPES AND SYMMETRIC GROTHENDIECK POLYNOMIALS

LAURA ESCOBAR AND ALEXANDER YONG

ABSTRACT. *Symmetric Grothendieck polynomials* are inhomogeneous versions of Schur polynomials that arise in combinatorial K -theory. A polynomial has *saturated Newton polytope* (SNP) if every lattice point in the polytope is an exponent vector. We show Newton polytopes of these Grothendieck polynomials and their homogeneous components have SNP. Moreover, the Newton polytope of each homogeneous component is a permutahedron. This addresses recent conjectures of C. Monical–N. Tokcan–A. Yong and of A. Fink–K. Mészáros–A. St. Dizier in this special case.

Let $s_\lambda(x_1, \dots, x_n)$ be the *Schur polynomial*, which is the generating series for semistandard Young tableaux of shape λ with entries in $[n] := \{1, 2, \dots, n\}$. By work of C. Lenart [Le00, Theorem 2.2], the **symmetric Grothendieck polynomial** is given by

$$(1) \quad G_\lambda(x_1, \dots, x_n) = \sum_{\mu} a_{\lambda\mu} s_\mu(x_1, \dots, x_n).$$

The sum is over partitions μ (identified with their Young diagrams in English notation) with $\leq n$ rows. The $(-1)^{|\mu| - |\lambda|} a_{\lambda\mu}$ counts the number of row and column strictly increasing skew tableaux of shape μ/λ with entries in $[n]$ such that the entries in row r are weakly less than $r - 1$.

By (1), $G_\lambda(x_1, \dots, x_n)$ is an inhomogeneous deformation of $s_\lambda(x_1, \dots, x_n)$ and itself symmetric. For example, if $n = 3$ and $\lambda = (3, 1, 0)$,

$$G_\lambda(x_1, x_2, x_3) = s_{(3,1)} - (2s_{(3,1,1)} + s_{(3,2,0)}) + 2s_{(3,2,1)} - s_{(3,2,2)}.$$

These polynomials appear in the study of K -theoretic Schubert calculus; we refer the reader to [Le00, Bu02] and the references therein for additional discussion.

More generally, A. Lascoux–M.-P. Schützenberger [LaSc82] recursively defined (possibly nonsymmetric) Grothendieck polynomials associated to any permutation $w \in \mathfrak{S}_n$. We mention that A. Buch [Bu02] discovered the *set-valued tableaux* formula for $G_\lambda(x_1, \dots, x_n)$; this formula is often taken as a definition in the literature. (Recently, C. Monical [Mo16] found a bijection between the aforementioned rules of C. Lenart and of A. Buch.)

The **Newton polytope** of a polynomial $f = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} c_\alpha x^\alpha \in \mathbb{C}[x_1, \dots, x_n]$ is the convex hull of its exponent vectors, i.e., $\text{Newton}(f) = \text{conv}(\{\alpha : c_\alpha \neq 0\}) \subseteq \mathbb{R}^n$. In [MoToYo17], f is said to have **saturated Newton polytope** (SNP) if $c_\alpha \neq 0$ whenever $\alpha \in \text{Newton}(f)$. A study of SNP and algebraic combinatorics was given in *loc. cit.*

If $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0)$, the **permutahedron** $\mathcal{P}_\lambda \subseteq \mathbb{R}^n$ is the convex hull of the \mathfrak{S}_n -orbit of λ . This theorem extends the old fact that $\text{Newton}(s_\lambda(x_1, \dots, x_n)) = \mathcal{P}_\lambda$:

Theorem. $G_\lambda(x_1, \dots, x_n)$ has SNP. In addition, each homogeneous component has SNP with Newton polytope being a permutahedron (as specified below in (3)).

Date: May 22, 2017.

The first assertion addresses [MoToYo17, Conjecture 5.5] for the case that the permutation w is **Grassmannian** at position n ; that is $w(i) < w(i+1)$ unless $i = n$. The second assertion responds, in this case, to a conjecture of A. Fink-K. Mészáros-A. St. Dizier [MeSt17, Conjecture 5.1]. In *loc. cit.*, these conjectures were proved for the case that $w = 1w'$ where w' is a **dominant permutation**, i.e., w is 132-avoiding.

Proof of the Theorem: Let $\mu^{(0)} := \lambda$. For $1 \leq k \leq n$ define $\mu^{(k)}$ to be $\mu^{(k-1)}$ with a box added in the northmost row r such that $\mu_r^{(k-1)} - \mu_r^{(0)} < r - 1$ and the addition of the box gives a Young diagram. Stop when no such r exists or $k = n$. Suppose we obtain N such partitions.

Recall that **dominance order** on partitions of a fixed size is defined by $\theta \leq_D \delta$ if $\sum_{j=1}^t \theta_j \leq \sum_{j=1}^t \delta_j$ for $t \geq 1$.

Claim A. $\mu^{(k)}$ is the \leq_D -maximum among all shapes μ of size $|\lambda| + k$ such that $a_{\lambda, \mu} \neq 0$.

Proof of Claim A: The skew shape $\mu^{(k)}/\lambda$ consists of the k boxes added to λ . We can inductively define a skew tableau T_k of this shape by adding the minimum possible label to T_{k-1} (in the box $\mu^{(k)}/\mu^{(k-1)}$) that maintains row and column strictness. It is straightforward that this tableau exists and witnesses $a_{\lambda, \mu^{(k)}} \neq 0$.

Let μ be a shape such that $a_{\lambda, \mu} \neq 0$. Then $\mu_i \leq \lambda_i + (i-1)$ for all i . Suppose that $\mu \not\leq_D \mu^{(k)}$ and let $r (> 1)$ be the first row such that $\mu_1 + \dots + \mu_r > \mu_1^{(k)} + \dots + \mu_r^{(k)}$. Then

$$\mu_1 + \dots + \mu_{r-1} \leq \mu_1^{(k)} + \dots + \mu_{r-1}^{(k)} \quad \text{and} \quad \mu_r > \mu_r^{(k)}.$$

This contradicts the construction of $\mu^{(k)}$ because by $\mu_r^{(k)} - \mu_r^{(0)} < \mu_r - \mu_r^{(0)} \leq r - 1$ $\mu^{(k)}$ must have another box in row r . \square

R. Rado's theorem [Ra52] states that for two partitions θ, δ of the same size,

$$(2) \quad \mathcal{P}_\theta \subseteq \mathcal{P}_\delta \iff \theta \leq_D \delta.$$

The Theorem's second assertion is immediate from (2) and Claim A. In fact if $G_\lambda[k]$ denotes the degree k homogeneous component of $G_\lambda(x_1, \dots, x_n)$ then

$$(3) \quad \text{Newton}(G_\lambda[k]) = \mathcal{P}_{\mu^{(k)}}.$$

Let $v^{(k)} \in \mathcal{P}_{\mu^{(k)}}$. Thus $v^{(k)}$ is a nonnegative vector (being a convex combination of nonnegative vectors). Rado's theorem implies that $v^{(k)}$ is *majorized* by $\mu^{(k)}$. That is, the rearrangement $(v^{(k)})^\downarrow$ of the components of $v^{(k)}$ into decreasing order satisfies

$$(4) \quad (v^{(k)})^\downarrow \leq_D \mu^{(k)}.$$

Suppose

$$v = \sum_{k=0}^N c_k v^{(k)} \quad \text{where} \quad \sum_{k=0}^N c_k = 1 \quad \text{and} \quad c_k \geq 0$$




FIGURE 1. The Newton polytope of $G_\lambda(x_1, x_2, x_3)$ for $\lambda = (3, 1, 0)$. Each color indicates degree.

is a convex combination of the vectors $v^{(k)}$.

Claim B. v is majorized by $\bar{\mu} := \sum_{k=0}^N c_k \mu^{(k)}$.

Proof of Claim B: Let $v^* := \sum_{k=0}^N c_k (v^{(k)})^\downarrow$. By (4), for any $t \geq 1$ we have $c_k \sum_{j=1}^t (v^{(k)})^\downarrow_j \leq c_k \sum_{j=1}^t \mu_j^{(k)}$. By summing both sides over all k and interchanging the order of summation we conclude v^* is majorized by $\bar{\mu}$. It is a standard property of majorization that $a + b$ is majorized by $a^\downarrow + b^\downarrow$ [MaOlAr11, Proposition A.1.b]. Thus v is majorized by v^* . Now use that majorization (being a preorder) is transitive. \square

Claim C. Suppose $|\bar{\mu}| - |\mu^{(0)}| = K$, then $\bar{\mu}$ is majorized by $\mu^{(K)}$.

Proof of Claim C: Let $r_k :=$ row on which k -th box gets added to $\mu^{(k)}$ (so $\mu^{(k)} = \mu^{(0)} + e_{r_1} + \dots + e_{r_k}$, where e_i is a standard basis vector).

Lemma 0.1. For any (row) r

$$\bar{\mu}_1 + \dots + \bar{\mu}_r = \mu_1^{(0)} + \dots + \mu_r^{(0)} + c_1 + 2c_2 + \dots + \ell c_\ell + \dots + \ell c_N,$$

where ℓ is the largest i such that $r_i = r$.

Proof. Suppose we added boxes $a, a+1, \dots, a+b$ to row r of $\mu^{(0)}$ in order to obtain $\mu^{(N)}$.

We write

$$\bar{\mu}_r = \underbrace{c_0 \mu_r^{(0)} + \dots + c_{a-1} \mu_r^{(a-1)}}_{\textcircled{1}} + \underbrace{c_a \mu_r^{(a)} + \dots + c_{a+b} \mu_r^{(a+b)}}_{\textcircled{2}} + \underbrace{c_{a+b+1} \mu_r^{(a+b+1)} + \dots + c_N \mu_r^{(N)}}_{\textcircled{3}}.$$

Next, $\mu_r^{(0)} = \dots = \mu_r^{(a-1)}$, so

$$\textcircled{1} = (c_0 + \dots + c_{a-1}) \mu_r^{(0)}.$$

Since $\mu_r^{(a+i)} = \mu_r^{(0)} + (i+1)$ for $i = 0, \dots, b$, then

$$\textcircled{2} = (c_a + \dots + c_{a+b}) \mu_r^{(0)} + (c_a + 2c_{a+1} + \dots + (b+1)c_{a+b}).$$

Finally, $\mu_r^{(a+b)} = \mu_r^{(a+b+1)} = \dots = \mu_r^{(n)}$, so

$$\textcircled{3} = (c_{a+b+1} + \dots + c_N) \mu_r^{(0)} + (c_{a+b+1} + \dots + c_N)(b+1).$$

Therefore we conclude $\bar{\mu}_r = \mu_r^{(0)} + c_a + 2c_{a+1} + \dots + (b+1)c_{a+b} + \dots + (b+1)c_N$. The lemma then follows by a simple induction. \square

The following is immediate from the definitions.

Lemma 0.2. Let $b_r = \mu_r^{(N)} - \mu_r^{(0)}$, i.e. b_r is the number of extra boxes $\mu^{(N)}$ has in row r . For any $r < r_K$

$$\mu_1^{(k)} + \dots + \mu_r^{(k)} = \mu_1^{(0)} + \dots + \mu_r^{(0)} + (b_1 + \dots + b_r).$$

Let ℓ be the largest i such that $r_i = r$. We consider two cases.

Case 1 ($r < r_K$): Observe that

$$c_1 + 2c_2 + \dots + \ell c_\ell + \dots + \ell c_N = (c_1 + \dots + c_N) + (c_2 + \dots + c_N) + \dots + (c_\ell + \dots + c_N) \leq \ell.$$

Since $b_1 + \cdots + b_r$ equals the number of boxes placed from rows 1 through r and the ℓ -th box is the last box placed in row r , then $\ell = b_1 + \cdots + b_r$. Combining this equality with the inequality just derived, we see

$$(5) \quad c_1 + 2c_2 + \cdots + \ell c_\ell + \cdots + \ell c_N \leq b_1 + \cdots + b_r.$$

By (5) together with Lemmas 0.1 and 0.2,

$$\bar{\mu}_1 + \cdots + \bar{\mu}_r \leq \mu_1^{(K)} + \cdots + \mu_r^{(K)}.$$

Case 2 ($r \geq r_K$): Here, we notice that

$$(6) \quad \mu_1^{(K)} + \cdots + \mu_r^{(K)} = \mu_1^{(0)} + \cdots + \mu_r^{(0)} + K.$$

Observe that

$$(7) \quad c_1 + 2c_2 + \cdots + \ell c_\ell + \cdots + \ell c_N \leq c_1 + 2c_2 + \cdots + Nc_N = K,$$

where the equality follows from Lemma 0.1. Apply Lemma 0.1 to the left hand side of (7) and use (6) to replace K on the right hand side, to conclude $\bar{\mu}_1 + \cdots + \bar{\mu}_r \leq \mu_1^{(K)} + \cdots + \mu_r^{(K)}$. Hence in either case, $\bar{\mu} \leq_D \mu^{(K)}$, as required. \square

Let w_1, \dots, w_M be any exponent vectors of $G_\lambda(x_1, \dots, x_n)$. SNPness means that if $w \in \text{conv}\{w_1, \dots, w_M\}$ is a lattice point then $[x^w]G_\lambda(x_1, \dots, x_n) \neq 0$. Suppose that $|w| = K$. Without loss of generality, $M = N$ and there is a unique vector w_k with $|w_k| = k$. Then by Claim B, w is majorized by $\bar{\mu}$. Claim C says $\bar{\mu}$ is majorized by $\mu^{(K)}$ and hence $w^\downarrow \leq_D \mu^{(K)}$. By (2) we conclude $w \in \mathcal{P}_{\mu^{(K)}}$, which by (3) completes the proof of the Theorem. Indeed, we have shown that $\text{Newton}(G_\lambda(x_1, \dots, x_n)) = \bigcup_{k=0}^N \mathcal{P}_{\mu^{(k)}}$. \square

ACKNOWLEDGEMENTS

We thank Karola Mészáros and Avery St. Dizier for a helpful communication. We used SAGE code of C. Monical during our investigation. AY was supported by an NSF grant.

REFERENCES

- [Bu02] A. Buch. A Littlewood-Richardson rule for the K -theory of Grassmannians. *Acta Math.*, 189(1):37–78, 2002.
- [LaSc82] A. Lascoux and M.-P. Schützenberger. Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux. *C. R. Acad. Sci. Paris Sér. I Math.*, 295(11):629–633, 1982.
- [Le00] C. Lenart. Combinatorial aspects of the K -theory of Grassmannians. *Ann. Comb.*, 4(1):67–82, 2000.
- [MaOlAr11] A. W. Marshall, I. Olkin and B. C. Arnold. *Inequalities: theory of majorization and its applications*. Springer Series in Statistics. Springer, New York, second edition, 2011.
- [MeSt17] K. Mészáros and A. St. Dizier. Generalized permutahedra to Grothendieck polynomials via flow polytopes. *Preprint*, arXiv:1705.02418
- [Mo16] C. Monical. Set-Valued Skyline Fillings. *Preprint*, arXiv:1611.08777
- [MoToYo17] C. Monical, N. Tokcan and A. Yong. Newton polytopes in algebraic combinatorics. *Preprint*, arXiv:1703.02583
- [Ra52] R. Rado. An inequality. *J. London Math. Soc.*, 27:1–6, 1952.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA–CHAMPAIGN, URBANA, IL 61801, USA

E-mail address: lescobar@illinois.edu, ayong@uiuc.edu