

EQUIVARIANT K -THEORY OF GRASSMANNIANS

OLIVER PECHENIK AND ALEXANDER YONG

ABSTRACT. We address a unification of the Schubert calculus problems solved by [A. Buch '02] and [A. Knutson-T. Tao '03]. That is, we prove a combinatorial rule for the structure coefficients in the torus-equivariant K -theory of Grassmannians with respect to the basis of Schubert structure sheaves. We thereby deduce the conjectural rule of [H. Thomas-A. Yong '13] for the same coefficients. Both rules are positive in the sense of [D. Anderson-S. Griffeth-E. Miller '11] (and moreover in a stronger form). Our work is based on the combinatorics of *genomic tableaux* and a generalization of [M.-P. Schützenberger '77]'s *jeu de taquin*.

CONTENTS

1. Introduction	2
1.1. Overview	2
1.2. Genomic tableaux	3
1.3. The ballot property	4
1.4. Tableau weights and the main theorem	4
1.5. Organization	6
2. Bundled tableaux and a reformulation of Theorem 1.3	7
3. Structure of the proof of Theorem 1.3	8
4. The base case of the recurrence	9
5. Good tableaux	12
6. Snakes of good tableaux	16
6.1. Snakes	16
6.2. Snake sections	18
7. Genomic jeu de taquin	19
7.1. Miniswaps	19
7.2. Swaps and slides	22
7.3. Examples	23
8. Ladders	25
9. Reverse genomic jeu de taquin	27
10. The reversal tree	32
10.1. Walkways	32
10.2. Walkway reversal	33

Date: June 5, 2015.

11. The recurrence coefficients	37
12. Weight preservation	43
12.1. Fine tableaux and their weights	43
12.2. Main claim about weight preservation	44
12.3. Pairing maps	46
13. Proof of the conjectural K_T rule from [ThYo13]	53
Appendix A. Proof of Proposition 7.7	58
Appendix B. Proof of Proposition 9.5	71
Appendix C. Block decomposition; completion of proof of Proposition 12.3(II)	82
Acknowledgments	91
References	92

1. INTRODUCTION

1.1. **Overview.** Let $X = \text{Gr}_k(\mathbb{C}^n)$ denote the Grassmannian of k -dimensional subspaces of \mathbb{C}^n . The action of $\text{GL}_n(\mathbb{C})$ on X restricts to an action of the Borel subgroup B of invertible upper triangular matrices and its subgroup T of invertible diagonal matrices. The T -fixed points $e_\lambda \in X$ are naturally indexed by Young diagrams λ contained in the rectangle $k \times (n - k)$. The **Schubert varieties** are the orbit closures $X_\lambda = \overline{B} \cdot e_\lambda$. The Poincaré duals $[X_\lambda]$ of their classes form a \mathbb{Z} -basis of the cohomology ring $H^*(X, \mathbb{Z})$.

The (classical) **Schubert structure constants** $c_{\lambda, \mu}^\nu$ are defined by $[X_\lambda] \cdot [X_\mu] = \sum_\nu c_{\lambda, \mu}^\nu [X_\nu]$. In Schubert calculus, one interprets $c_{\lambda, \mu}^\nu \in \mathbb{Z}_{\geq 0}$ as the number of points (when finite) in a generic triple intersection of Schubert varieties. Combinatorially, $c_{\lambda, \mu}^\nu$ is computed, in a manifestly nonnegative manner, by Littlewood-Richardson rules. The first such rule was stated by D. E. Littlewood-A. R. Richardson in the 1930s [LiRi34] in their study of the representation theory of the symmetric group. The first rigorous proof of a rule was given by M.-P. Schützenberger [Sc77] in the 1970s. These rules describe $c_{\lambda, \mu}^\nu$ as a count of certain Young tableaux.

In the modern Schubert calculus, there is significant attention on the problem of generalizing the above work to richer cohomology theories. Early last decade, two problems of this type were solved. A. Buch [Bu02] found the first rule for the multiplication of the Schubert structure sheaves in K -theory. His rule is positive after accounting for a predictable alternation of sign. Separately, A. Knutson-T. Tao [KnTa03] introduced *puzzles* to give the first rule for equivariant Schubert calculus that is positive in the sense of [Gr01].

We turn to a unification of these problems. Let $K_T(X)$ denote the Grothendieck ring of T -equivariant vector bundles over X . This ring has a natural $K_T(\text{pt})$ -module structure and an additive basis given by the classes of Schubert structure sheaves; for background, we refer the reader to, e.g., [KoKu90, AnGrMi11] and the references therein. The analogues of Littlewood-Richardson coefficients are the Laurent polynomials $K_{\lambda, \mu}^\nu \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}] \cong K_T(\text{pt})$ defined by

$$[\mathcal{O}_{X_\lambda}] \cdot [\mathcal{O}_{X_\mu}] = \sum_{\nu \subseteq k \times (n-k)} K_{\lambda, \mu}^\nu [\mathcal{O}_{X_\nu}],$$

where $[\mathcal{O}_{X_\lambda}]$ is the class of the structure sheaf of X_λ . These coefficients may be algebraically computed using *double Grothendieck polynomials*; see [LaSc82, FuLa94]. The problem addressed by this paper is to prove a combinatorial rule for $K_{\lambda,\mu}^\nu$.

We give a summary of earlier contributions to the problem: A. Knutson-R. Vakil conjectured a (still-open) formula for $K_{\lambda,\mu}^\nu$ in terms of puzzles (reported in [CoVa05, §5]). V. Kreiman [Kr05] proved a rule for the case $\lambda = \nu$, corresponding to a certain localization (cf. Section 4). C. Lenart-A. Postnikov [LePo07] determined a rule for the case $\lambda = (1)$ (in a broader context applicable to any generalized flag variety); we use this result. Later, W. Graham-S. Kumar [GrKu08] determined the coefficients in the case $X = \mathbb{P}^{n-1}$. “Positivity” of $K_{\lambda,\mu}^\nu$ (in a more general context) was geometrically established by D. Anderson-S. Griffeth-E. Miller [AnGrMi11]. More recently, A. Knutson [Kn10] obtained a puzzle rule in $K_\tau(X)$ for the different problem of multiplying the class of a Schubert structure sheaf by that of an *opposite* Schubert structure sheaf. Finally, H. Thomas and the second author conjectured the first Young tableau rule for $K_{\lambda,\mu}^\nu$ [ThYo13, Conjecture 4.7]; they showed their conjectural rule is positive in the sense of [AnGrMi11]; see [ThYo13, §4.1].

This paper introduces and proves an [AnGrMi11]-positive rule for $K_{\lambda,\mu}^\nu$ (Theorem 1.3); in fact our rule exhibits a further property of the coefficients which seems at present not to have a geometric explanation. Our rule allows us to deduce the aforementioned conjecture of [ThYo13]. Indeed, our approach completes the strategy set out in *loc. cit.* and our Theorem 1.3 is a generalization of the rule of [ThYo13] for T -equivariant cohomology. The general approach of our proof is to relate our combinatorial rule to a K -theoretic generalization of a recurrence proven by A. Molev-B. Sagan [MoSa99] and A. Knutson-T. Tao [KnTa03] (who also credit A. Okounkov). A similar approach was employed by A. Buch [Bu15] who gave a rule for the equivariant quantum cohomology of Grassmannians, cf. [BuMi11]. (The case of non-equivariant quantum cohomology had been previously handled geometrically by [Co09] and combinatorially by [BKPT13], cf. [BuKrTa03].)

We introduce *genomic tableaux* and a generalization of M.-P. Schützenberger’s *jeu de taquin* [Sc77]. C. Monical has reported applications of genomic tableaux in the study of *Lascoux polynomials* (see, e.g., [RoYo13] and references therein) and K -theoretic analogues of *Demazure atoms*, extending results of [HLMvW11]. These tableaux also give a new rule for (non-equivariant) K -theory of Grassmannians; the announcement [PeYo15] outlines applications to analogous problems when X is replaced by Lagrangian or maximal orthogonal Grassmannians. In addition, one hopes to use our results to shed light on the A. Knutson-R. Vakil’s puzzle conjecture for $K_{\lambda,\mu}^\nu$. Note the puzzle conjecture does not directly recover an earlier ordinary K -theory puzzle rule; this is a qualitative difference with the rules of this text. Moreover, closely related to the equivariant Schubert calculus of X , the combinatorial rule of A. Molev-B. Sagan [MoSa99] solves a *triple Schubert calculus* problem in $H^*(\mathrm{GL}_n(\mathbb{C})/\mathrm{B} \times X \times \mathrm{GL}_n(\mathbb{C})/\mathrm{B})$ (see [KnTa03, §6]). Our methods should extend to give a K -theoretic analogue, cf. [KnTa03, §6.2].

1.2. Genomic tableaux. A **genomic tableau** is a Young diagram filled with (subscripted) labels i_j where $i \in \mathbb{Z}_{>0}$ and the j ’s that appear for each i form an initial segment of $\mathbb{Z}_{>0}$. It is **edge-labeled** of shape ν/λ if each horizontal edge of a box weakly below the southern border of λ (viewed as a lattice path from $(0, 0)$ to $(k, n - k)$) is filled with a subset of $\{i_j\}$.

Let $x \rightarrow$ be the box immediately east of x , $x \uparrow$ the box immediately north of x , etc. For a box x , let \bar{x} denote the upper horizontal edge of x and \underline{x} denote the lower horizontal edge.

We write $\text{family}(i_j) = i$. We distinguish two orders on subscripted labels. Say $i_j < k_\ell$ if $i < k$. Write $i_j \prec k_\ell$ if $i < k$ or $i = k$ with $j < \ell$. Note that \prec is a total order, while $<$ is not.

A genomic tableau T is **semistandard** if the following four conditions hold:

- (S.1) $\text{label}(x) \prec \text{label}(x^{\rightarrow})$;
- (S.2) every label is $<$ -strictly smaller than any label South¹ in its column;
- (S.3) if i_j, k_ℓ appear on the same edge then $i \neq k$;
- (S.4) if i_j is West of i_k , then $j \leq k$.

Refer to the multiset $\{i_j\}$ (for fixed i and j) collectively as a **gene**. The **content** of T is (c_1, c_2, c_3, \dots) where c_i is the number of genes of family i . Suppose x is in row r . A label i_j is **too high** if $i \geq r$ and $i_j \in \bar{x}$, or alternatively if $i > r$ and $i_j \in x$ or $i_j \in \underline{x}$.

Example 1.1. For $\lambda = (4, 2, 2, 1)$ and $\nu = (6, 5, 4, 3, 2)$ consider the genomic tableau T :

					1 ₂	1 ₃
			1 ₂	2 ₁	2 ₂	
			2 ₁	3 ₂		
		1 ₁	3 ₂			
	2 ₁	3 ₂	4 ₂			
2 ₁	3 ₂					
3 ₁	4 ₁					

The content of T is $(3, 2, 2, 2)$. The tableau T is not semistandard, since the second column from the left fails (S.2). If we deleted the 3_2 from the edge, the result would be semistandard. No label is too high.

□

1.3. The ballot property. A **genotype** G of T is a choice of one label from each gene of T . Let $\text{word}(G)$ be obtained by reading G down columns from right to left. (If there are multiple labels on an edge, read them from smallest to largest in \prec -order.) Then G is **ballot** if in every initial segment of $\text{word}(G)$, there are at least as many labels of family i as of family $i + 1$, for each $i \geq 1$. We say T is **ballot** if all of its genotypes are ballot. Let $\text{BallotGen}(\nu/\lambda)$ be the set of ballot, semistandard, edge-labeled genomic tableaux of shape ν/λ where no label is too high.

Example 1.2. Let $T = \begin{array}{|c|c|} \hline & 1_2 \\ \hline 1_1 & 2_1 \\ \hline \end{array}$ and $U = \begin{array}{|c|c|} \hline & 1_1 \\ \hline 1_1 & 2_1 \\ \hline \end{array}$. Then T is ballot: the one genotype (itself) has reading word is $1_2 2_1 1_1$, which is a ballot sequence. U is not ballot: it has two genotypes $\begin{array}{|c|c|} \hline & \\ \hline 1_1 & 2_1 \\ \hline \end{array}$ and $\begin{array}{|c|c|} \hline 1_1 & \\ \hline & 2_1 \\ \hline \end{array}$ and the word for the former is $2_1 1_1$, which is not ballot. □

1.4. Tableau weights and the main theorem. Let $T \in \text{BallotGen}(\nu/\lambda)$. For a box x , $\text{Man}(x)$ is the “Manhattan distance” from the southwest corner (point) of $k \times (n - k)$ to the northwest corner (point) of x (the length of any north-east lattice path between the corners).

For a gene \mathcal{G} , let $N_{\mathcal{G}}$ be the number of genes \mathcal{G}' with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$ and $\mathcal{G}' \succ \mathcal{G}$. For instance, in Example 1.1, $N_{1_1} = 2$ since the genes 1_2 and 1_3 are of the same family as 1_1 (namely family 1) but $1_1 \prec 1_2, 1_3$.

¹Throughout, we write “West”, “west” and “NorthWest” to mean “strictly west”, “weakly west” and “strictly north and strictly west” respectively, etc.

If $\ell = i_j \in \underline{x}$ and x is in row r , then

$$(1.1) \quad \text{edgefactor}(\ell) := \text{edgefactor}_{\underline{x}}(i_j) := 1 - \frac{t_{\text{Man}(x)}}{t_{r-i+N_{i_j}+1+\text{Man}(x)}}.$$

The **edge weight** $\text{edgewt}(T)$ is $\prod_{\ell} \text{edgefactor}(\ell)$; the product is over edge labels of T .

A nonempty box x in row r is **productive** if $\text{label}(x) < \text{label}(x^{\rightarrow})$. If $i_j \in x$, set

$$(1.2) \quad \text{boxfactor}(x) := \frac{t_{\text{Man}(x)+1}}{t_{r-i+N_{i_j}+1+\text{Man}(x)}}.$$

The **box weight** of a tableau T is $\text{boxwt}(T) := \prod_x \text{boxfactor}(x)$, where the product is over all productive boxes of T . The **weight** of T is $\text{wt } T := (-1)^{d(T)} \times \text{boxwt}(T) \times \text{edgewt}(T)$. Here $d(T) = \sum_{\mathcal{G}} (|\mathcal{G}| - 1)$, where the sum is over all genes \mathcal{G} and $|\mathcal{G}|$ is the (multiset) cardinality of \mathcal{G} . Set

$$L_{\lambda, \mu}^{\nu} := \sum_T \text{wt } T,$$

where the sum is over all $T \in \text{BallotGen}(\nu/\lambda)$ that have content μ .

Theorem 1.3 (Main Theorem). $K_{\lambda, \mu}^{\nu} = L_{\lambda, \mu}^{\nu}$.

This provides the first proved rule for $K_{\lambda, \mu}^{\nu}$ that is manifestly [AnGrMi11]-positive. That is, let $z_i := \frac{t_i}{t_{i+1}} - 1$. For $j > i$, we have

$$(1.3) \quad \frac{t_i}{t_j} = \prod_{k=i}^{j-1} (z_k + 1) \quad \text{and} \quad 1 - \frac{t_i}{t_j} = - \left(\prod_{k=i}^{j-1} (z_k + 1) - 1 \right).$$

Therefore, $(-1)^{\#\text{edge labels}} \times \text{boxwt}(T) \times \text{edgewt}(T)$ is z -positive. Since clearly $d(T) = |\nu| - |\lambda| - |\mu| + \#\text{edge labels}$, we have that $(-1)^{|\nu|-|\lambda|-|\mu|} L_{\lambda, \mu}^{\nu} = \sum_T (-1)^{|\nu|-|\lambda|-|\mu|} \text{wt } T$ is z -positive. This positivity is the same as that of [AnGrMi11, Corollary 5.3] after the substitution $z_i \mapsto e^{\alpha_i} - 1$ where α_i is the i -th simple root for the root system A_{n-1} .

Example 1.4. To compute $K_{(2),(2,1)}^{(2,2)}$ for $\text{Gr}_2(\mathbb{C}^4)$, the required tableaux are

$$T_1 = \begin{array}{|c|c|} \hline & \\ \hline 1_1 & 1_2 \\ \hline 2_1 & \\ \hline \end{array}, \quad T_2 = \begin{array}{|c|c|} \hline & \\ \hline 1_1 & 1_2 \\ \hline & 2_1 \\ \hline \end{array}, \quad T_3 = \begin{array}{|c|c|} \hline & \\ \hline 1_1 & 1_2 \\ \hline 2_1 & 2_1 \\ \hline \end{array}, \quad T_4 = \begin{array}{|c|c|} \hline & 1_2 \\ \hline 1_1 & 2_1 \\ \hline \end{array}, \quad T_5 = \begin{array}{|c|c|} \hline & 1_2 \\ \hline 1_1 & 2_1 \\ \hline 2_1 & \\ \hline \end{array}$$

Then

- $\text{edgewt}(T_1) = 1 - \frac{t_1}{t_2}$, $\text{boxwt}(T_1) = \frac{t_3}{t_4}$ and $d(T_1) = 0$;
- $\text{edgewt}(T_2) = 1 - \frac{t_2}{t_3}$, $\text{boxwt}(T_2) = \frac{t_3}{t_4}$ and $d(T_2) = 0$;
- $\text{edgewt}(T_3) = (1 - \frac{t_1}{t_2})(1 - \frac{t_2}{t_3})$, $\text{boxwt}(T_3) = \frac{t_3}{t_4}$ and $d(T_3) = 1$;
- $\text{edgewt}(T_4) = (1 - \frac{t_3}{t_4})$, $\text{boxwt}(T_4) = \frac{t_2}{t_4}$ and $d(T_4) = 0$; and
- $\text{edgewt}(T_5) = (1 - \frac{t_1}{t_2})(1 - \frac{t_3}{t_4})$, $\text{boxwt}(T_5) = \frac{t_2}{t_4}$ and $d(T_5) = 1$.

Hence

$$K_{(2),(2,1)}^{(2,2)} = \left(1 - \frac{t_1}{t_2}\right) \frac{t_3}{t_4} + \left(1 - \frac{t_2}{t_3}\right) \frac{t_3}{t_4} - \left(1 - \frac{t_1}{t_2}\right) \left(1 - \frac{t_2}{t_3}\right) \frac{t_3}{t_4} + \left(1 - \frac{t_3}{t_4}\right) \frac{t_2}{t_4} - \left(1 - \frac{t_1}{t_2}\right) \left(1 - \frac{t_3}{t_4}\right) \frac{t_2}{t_4}.$$

Observe that, after rewriting using (1.3), each term is z -negative, in agreement with the discussion above; that is,

$$\begin{aligned}
(-1)^{|(2,2)|-|(2)|-|(2,1)|} K_{(2),(2,1)}^{(2,2)} &= -(-z_1)(z_3 + 1) - (-z_2)(z_3 + 1) + (-z_1)(-z_2)(z_3 + 1) \\
&\quad -(-z_3)(z_2 + 1)(z_3 + 1) + (-z_1)(-z_3)(z_2 + 1)(z_3 + 1) \\
&= z_1(z_3 + 1) + z_2(z_3 + 1) + z_1 z_2(z_3 + 1) \\
&\quad + z_3(z_2 + 1)(z_3 + 1) + z_1 z_3(z_2 + 1)(z_3 + 1)
\end{aligned}$$

is z -positive (without any cancellation needed). \square

There is a stronger positivity property exhibited by the rule of Theorem 1.3. The work of [AnGrMi11] generalizes the positivity of W. Graham [Gr01]: the equivariant Schubert structure coefficients are polynomials with nonnegative integer coefficients in the simple roots α_i . In [Kn10], A. Knutson observes W. Graham’s geometric argument further implies the coefficients can be expressed as polynomials with nonnegative integer coefficients in the positive roots *such that each monomial is square-free*. Moreover, A. Knutson raises the issue of finding a “proper analogue” in equivariant K -theory for this square-free property. For X , we offer:

Corollary 1.5 (Strengthened [AnGrMi11]-positivity). *Let $z_{ij} := \frac{t_i}{t_j} - 1$. Then $(-1)^{|\nu|-|\lambda|-|\mu|} K_{\lambda,\mu}^\nu$ is expressible as a polynomial with nonnegative integer coefficients in the z_{ij} ’s such that each monomial is square-free.*

Proof. The nonnegativity of the coefficients is immediate from each z_{ij} being positive in the z_i ’s. It remains to show each monomial in our expression $L_{\lambda,\mu}^\nu$ is square-free.

Consider a $T \in \text{BallotGen}(\nu/\lambda)$. Every edgefactor(ℓ) is of the form $-z_{ij}$, while every boxfactor(x) is of the form $z_{ij} + 1$. Define an (i, j) -label to be either an edge label with edgefactor(ℓ) = $-z_{ij}$ or a label ℓ in a productive box x with boxfactor(x) = $z_{ij} + 1$.

Suppose ℓ, ℓ' are (i, j) -labels of T . Say $\ell \in x$ or \bar{x} and $\ell' \in y$ or \bar{y} . Since both are $(i, -)$ -labels, $\text{Man}(x) = \text{Man}(y)$. Hence x and y are boxes of the same diagonal. We may assume x northwest of y . Let ℓ be an instance of m_n and ℓ' an instance of p_q . Since both are $(-, j)$ -labels, $\text{row}(x) - m + N_{m_n} = \text{row}(y) - p + N_{p_q}$. By (S.1) and (S.2), $m + r(y) - r(x) \leq p$, so $N_{m_n} = r(y) - r(x) + m - p + N_{p_q} \leq p - p + N_{p_q} = N_{p_q}$. Hence by ballotness of T , $x = y$ and moreover $m = p$. Therefore by (S.2) and (S.3), $\ell = \ell'$, and thus T contains at most one (i, j) -label and each monomial in our expression is square-free. \square

We do not know a geometric explanation for Corollary 1.5. However, based on this result, one speculates that for any G/P , if for each positive root α we set $z_\alpha := e^\alpha - 1$, then the corresponding Schubert structure coefficients for $K_T(G/P)$ may be expressed in a square-free manner with nonnegative coefficients in the z_α ’s.

1.5. Organization. The first key to the proof is to reformulate Theorem 1.3 in terms of the more technical *bundled* tableaux that are appropriate for the inductive argument; this is presented in Section 2. In Section 3, we outline this inductive argument that the rule of Theorem 1.3 satisfies the key recurrence alluded to above. The base case is in Section 4. Both the plan of induction and the base case may be considered routine.

The core of the argument lies in Sections 5–12. There we construct local swapping rules, defining a genomic generalization of M.-P. Schützenberger’s *jeu de taquin*. This permits us

to establish a combinatorial map of formal sums of tableaux. This part of the argument is developed as a sequence of four main ideas:

- (1) To show well-definedness of the map, we identify and characterize the class of *good tableaux* that arise via genomic *jeu de taquin* (Sections 5, 6 and 7).
- (2) To establish surjectivity, we define reverse swaps and slides and prove these operations keep one in the class of good tableaux (Sections 8 and 9).
- (3) To prove that the map respects the coefficients of the key recurrence, we define and prove properties of a *reversal tree* (Sections 10 and 11).
- (4) The map is weight-preserving. However, a significant subtlety is that it is not generally weight-preserving on *individual* tableaux. To establish this property of the map, we need involutions that pair tableaux (Section 12).


In Section 13, we recall the conjecture of [ThYo13] and prove it from Theorem 1.3; this argument is essentially independent of the rest of the paper. The three appendices isolate essentially straightforward but long technical proofs of important propositions.

2. BUNDLED TABLEAUX AND A REFORMULATION OF THEOREM 1.3

A tableau $T \in \text{BallotGen}(\nu/\lambda)$ is **bundled** if every edge label is the westmost label of its gene. For example, in Example 1.4, only T_3 is *not* bundled (the eastmost 2_1 is to blame). We denote the set of bundled tableaux of shape ν/λ by $\text{Bundled}(\nu/\lambda)$.

Define a surjection $\text{Bun} : \text{BallotGen}(\nu/\lambda) \twoheadrightarrow \text{Bundled}(\nu/\lambda)$. This sends T to $\text{Bun}(T)$ by deleting each edge label of T that is not maximally west in its gene. If $B \in \text{Bundled}(\nu/\lambda)$, then any $T \in \text{Bun}^{-1}(B)$ differs from B by having (possibly 0) additional edge labels. Let E_{i_j} be the edges where i_j appears in some $T \in \text{Bun}^{-1}(B)$ but not in B , i.e., the set of edges of B where adding an i_j would yield an element of $\text{BallotGen}(\nu/\lambda)$. We say B has a **virtual label** i_j on each edge of E_{i_j} . We denote a virtual label i_j by $\textcircled{i_j}$.

Example 2.1. All virtual labels are depicted below:



□

For $B \in \text{Bundled}(\nu/\lambda)$, let

$$(2.1) \quad \text{wt}(B) = \sum_{T \in \text{Bun}^{-1}(B)} \text{wt}(T).$$

Let $B_{\lambda, \mu}^\nu$ denote the set of tableaux in $\text{Bundled}(\nu/\lambda)$ with content μ .

Proposition 2.2. $L_{\lambda, \mu}^\nu = \sum_{B \in B_{\lambda, \mu}^\nu} \text{wt}(B)$.

Proof. Immediate from (2.1) and the definition of $L_{\lambda,\mu}^\nu$. \square

Compute $\widetilde{\text{wt}}(B)$ as a product: an edge label ℓ contributes a factor of $\text{edgefactor}(\ell)$ and each productive box x contributes a factor of $\text{boxfactor}(x)$. Each virtual label $\mathbb{L} \in \underline{x}$ contributes $1 - \text{edgefactor}_{\underline{x}}(\ell)$ (where the latter is calculated as if \mathbb{L} were instead ℓ). Multiply by $(-1)^{d(T)}$ where $d(T) = \sum_{\mathcal{G}} (|\mathcal{G}| - 1)$ and here $|\mathcal{G}|$ is interpreted to be the multiset cardinality of non-virtual \mathcal{G} in T .

Example 2.3. For B from Example 2.1, $\widetilde{\text{wt}}(B) = (-1)^1 \cdot \left(1 - \frac{t_2}{t_8}\right) \cdot \frac{t_2}{t_4} \frac{t_4}{t_6} \frac{t_6}{t_9} \frac{t_8}{t_8} \frac{t_{11}}{t_{11}} \cdot \frac{t_3}{t_5} \frac{t_4}{t_9} \frac{t_5}{t_7} \frac{t_8}{t_{10}} \frac{t_9}{t_{11}}$. \square

Lemma 2.4. $\text{wt}(B) = \widetilde{\text{wt}}(B)$.

Proof. Let m be the number of virtual labels in B and a_i be the non-virtual weight of the i -th virtual label (listed in some given order). By the weights' definitions, the lemma follows from the ‘‘inclusion-exclusion’’ identity $\prod_{i \in [m]} a_i = \sum_{S \subseteq [m]} (-1)^{|S|} \prod_{i \in S} (1 - a_i)$. \square

3. STRUCTURE OF THE PROOF OF THEOREM 1.3

Let $\lambda^+ := \{\rho \supseteq \lambda : \rho/\lambda \text{ has no two boxes in the same row or column}\}$ and $\nu^- := \{\delta \subsetneq \nu : \nu/\delta \text{ has no two boxes in the same row or column}\}$. For a set D of boxes, let $\text{wt } D := \prod_{x \in D} \frac{t_{\text{Man}(x)}}{t_{\text{Man}(x)+1}}$.

Proposition 3.1 (Key recurrence).

$$(3.1) \quad \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} K_{\rho,\mu}^\nu = K_{\lambda,\mu}^\nu (1 - \text{wt } \nu/\lambda) + \sum_{\delta \in \nu^-} (-1)^{|\nu/\delta|+1} K_{\lambda,\mu}^\delta \text{wt } \delta/\lambda.$$

Proof. The Chevalley formula in equivariant K -theory [LePo07, Corollary 8.2] implies:

$$[\mathcal{O}_{X_\lambda}][\mathcal{O}_{X_\square}] = [\mathcal{O}_{X_\lambda}](1 - \text{wt } \lambda) + \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} [\mathcal{O}_{X_\rho}] \text{wt } \lambda.$$

Thus, the coefficient of $[\mathcal{O}_{X_\nu}]$ in $([\mathcal{O}_{X_\lambda}][\mathcal{O}_{X_\square}])[\mathcal{O}_{X_\mu}]$ is

$$K_{\lambda,\mu}^\nu (1 - \text{wt } \lambda) + \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} K_{\rho,\mu}^\nu \text{wt } \lambda.$$

On the other hand, the coefficient of $[\mathcal{O}_{X_\nu}]$ in $([\mathcal{O}_{X_\lambda}][\mathcal{O}_{X_\mu}])[\mathcal{O}_{X_\square}]$ is

$$K_{\lambda,\mu}^\nu (1 - \text{wt } \nu) + \sum_{\delta \in \nu^-} (-1)^{|\nu/\delta|+1} K_{\lambda,\mu}^\delta \text{wt } \delta.$$

The proposition then follows from associativity and commutativity:

$$([\mathcal{O}_{X_\lambda}][\mathcal{O}_{X_\square}])[\mathcal{O}_{X_\mu}] = ([\mathcal{O}_{X_\lambda}][\mathcal{O}_{X_\mu}])[\mathcal{O}_{X_\square}]. \quad \square$$

To prove $K_{\lambda,\mu}^\nu = L_{\lambda,\mu'}^\nu$ we induct on $|\nu/\lambda|$. Proposition 4.1 is the base case: $K_{\lambda,\mu}^\lambda = L_{\lambda,\mu'}^\lambda$; this is proved using the description of $L_{\lambda,\mu}^\lambda$ from Section 1.

The remaining cases use the description of $L_{\lambda,\mu}^\nu$ from Proposition 2.2. Assume $K_{\theta,\mu}^\tau = L_{\theta,\mu}^\tau$ when $|\tau/\theta| \leq h$. Suppose we are given λ, ν with $|\nu/\lambda| = h + 1$. We show that $L_{\lambda,\mu}^\nu$ satisfies (3.1). Since Proposition 3.1 asserts $K_{\lambda,\mu}^\nu$ also satisfies (3.1) we will be done by induction.

Fix λ, μ, ν with $\lambda \subsetneq \nu$. Define the formal sum

$$\Lambda^+ := \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} \sum_{T \in B_{\rho, \mu}^\nu} T.$$

Similarly define

$$\Lambda := (1 - \text{wt } \nu/\lambda) \sum_{T \in B_{\lambda, \mu}^\nu} T \quad \text{and} \quad \Lambda^- := \sum_{\delta \in \nu^-} (-1)^{|\nu/\delta|+1} (\text{wt } \delta/\lambda) \sum_{T \in B_{\lambda, \mu}^\delta} T.$$

In Section 7.2, we define an operation $\text{slide}_{\rho/\lambda}$ that takes as input $T \in \Lambda^+$ and returns a formal sum of tableaux with coefficients from $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. The construction of $\text{slide}_{\rho/\lambda}$ and proof of its correctness are found in Sections 5–7. Specifically, Corollary 7.11 shows the tableaux in the formal sum are from $B_{\lambda, \mu}^\nu \cup (\bigcup_{\delta \in \nu^-} B_{\lambda, \mu}^\delta)$.

In Section 11 we prove that

$$\text{slide}(\Lambda^+) := \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} \sum_{T \in B_{\rho, \mu}^\nu} \text{slide}_{\rho/\lambda}(T) = \Lambda + \Lambda^-;$$

see Proposition 11.8 for the precise statement. Finally Proposition 12.3 shows that $\text{wt } \Lambda^+ = \text{wt } \text{slide}(\Lambda^+)$, so $\sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} L_{\rho, \mu}^\nu = L_{\lambda, \mu}^\nu (1 - \text{wt } \nu/\lambda) + \sum_{\delta \in \nu^-} (-1)^{|\nu/\delta|+1} L_{\lambda, \mu}^\delta \text{wt } \delta/\lambda$. This completes the proof that the Laurent polynomials $L_{\lambda, \mu}^\nu$ defined by the rule of Proposition 2.2 equal $K_{\lambda, \mu}^\nu$. Hence we have completed our proof of Theorem 1.3. \square

4. THE BASE CASE OF THE RECURRENCE

A different rule for the case $K_{\lambda, \mu}^\lambda$ was given by V. Kreiman [Kr05]. We give an independent proof of the following:

Proposition 4.1 (Base case of the recurrence). $K_{\lambda, \mu}^\lambda = L_{\lambda, \mu}^\lambda$.

Proof. We use the original (unbundled) definition of $L_{\lambda, \mu}^\lambda$ from Section 1.

One says that $\pi \in S_n$ is a **Grassmannian permutation** if there is at most one k such that $\pi(k) > \pi(k+1)$. The Grassmannian permutation for $\lambda \subseteq k \times (n-k)$ is the (unique) Grassmannian permutation $\pi_\lambda \in S_n$ defined by $\pi_\lambda(i) = i + \lambda_{k-i+1}$ for $1 \leq i \leq k$ and $\pi(i) < \pi(i+1)$ for $i \neq k$.

Let $w', v' \in S_n$ be the Grassmannian permutations for the conjugate diagrams $\lambda', \mu' \subseteq (n-k) \times k$. The following identity relates $K_{\lambda, \mu}^\lambda$ to the localization of the class $[\mathcal{O}_{X_\lambda}]$ at the T -fixed point e_μ , expressed as a specialization of a double Grothendieck polynomial:

Lemma 4.2. $K_{\lambda, \mu}^\lambda = \overline{\mathfrak{G}_{v'}(t_{w'(1)}, \dots, t_{w'(n)}; t_1, \dots, t_n)}$, where $\overline{f(t_1, \dots, t_n)}$ is obtained by applying the substitution $t_j \mapsto t_{n-j+1}$ to $f(t_1, \dots, t_n)$.

Proof. This lemma is known to experts, but for completeness we give details and references. Suppose X_w is a Schubert variety in $\text{GL}_n(\mathbb{C})/\text{B}$. We have in $K_{\text{T}}(\text{GL}_n(\mathbb{C})/\text{B})$,

$$(4.1) \quad [\mathcal{O}_{X_v}][\mathcal{O}_{X_w}] = K_{v,w}^w[\mathcal{O}_{X_w}] + \sum_{\theta \neq w} K_{v,w}^\theta[\mathcal{O}_{X_\theta}].$$

It is known that $K_{v,w}^\theta = 0$ unless $v \leq \theta$ in Bruhat order; this follows for instance from the equivariant K -theory localization formula of M. Willems [Wi06] together with the *mutatis mutandis* modification of the proof of [KnTa03, Proposition 1].

where the factors correspond to the entries of $\xi(T)$ as read up columns from left to right (i.e., consistent with the order of factors of $\text{wt}(T)$ above).

Since $\lambda' = (3, 2, 1, 1)$ we have $w' = 2357146$ (one-line notation). So substituting, we get

$$\text{SVSSYTEqwt}(\xi(T)) = (-1)^1 \left(1 - \frac{t_7}{t_2}\right) \left(1 - \frac{t_5}{t_2}\right) \left(1 - \frac{t_3}{t_1}\right) \left(1 - \frac{t_2}{t_1}\right) \left(1 - \frac{t_7}{t_4}\right) \left(1 - \frac{t_5}{t_4}\right).$$

The reader can check $\overline{\text{SVSSYTEqwt}(\xi(T))} = \text{wt}(T)$, in agreement with the lemma. \square

(ξ^{-1} is well-defined and is weight-preserving): Let $U \in \mathcal{B}$. That $\xi^{-1}(U)$ is an edge-labeled genomic tableau is immediate from the column strictness of U . Ballotness follows from the row increasingness of U .

We now check that no label of $\xi^{-1}(U)$ is too high. Suppose c is a **bad** label in U in (matrix) position (r, i) , i.e., one such that the label $i_{\mu_{i+1-r}}$ placed in column c of λ/λ is too high. Observe that every label c' North of c and in the same column of U is also bad: this is since c' corresponds to placing another label of family i in the weakly shorter column c' East of column c (since $c' < c$). Thus we may assume c is in the northmost row of U , i.e., $r = 1$. Now if $i = 1$, then since c is bad, it must be that $\lambda'_{n-k-c+1} = 0$, so $w'(c) = c + 0$. Now c contributes a factor of $1 - \frac{x_c}{y_c}$ to $\text{SVSSYTw}(U)$ and hence a factor of $1 - \frac{t_{c+0}}{t_c} = 0$ to $\text{SVSSYTEqwt}(U)$. That is, $\text{SVSSYTEqwt}(U) = 0$, so $U \notin \mathcal{B}$, a contradiction. Otherwise, we may also assume $i > 1$ is smallest such that a label in $(r = 1, i)$ is bad. Since no label c' in $(r = 1, i - 1)$ of U is bad, it must be that c is “barely” bad, i.e.,

$$(4.2) \quad \lambda'_{n-k-c+1} = i - 1$$

(column c is one box too short). However, c contributes a factor of $1 - \frac{x_c}{y_{c+i-1}}$ to $\text{SVSSYTw}(U)$ and hence a factor of $1 - \frac{t_{c+\lambda'_{n-k-c+1}}}{t_{c+i-1}}$ to $\text{SVSSYTEqwt}(U)$. This latter factor is 0 precisely by (4.2). Hence again $U \notin \mathcal{B}$, a contradiction. Thus U has no bad labels and thus no label of $\xi^{-1}(U)$ is too high, as desired.

The sign appearing in $\text{wt} \xi^{-1}(U)$ records the difference between $|\mu|$ and the number of labels in $\xi^{-1}(U)$, while the sign in $\overline{\text{SVSSYTEqwt}(U)}$ records the difference between $|\mu|$ and number of labels in U . Since the number of labels in U is clearly the same as the number of labels in $\xi^{-1}(U)$, these signs are equal.

We check that the weight assigned to a label c of U in position (r, i) is the same as the edgefactor assigned to the corresponding label $i_{\mu_{i+1-r}}$ at the bottom of column c in $\xi^{-1}(U)$. The label c is assigned the weight

$$\text{SSYTEqfactor}_{(r,i)}(c) := 1 - \frac{x_c}{y_{c+i-r}} = 1 - \frac{t_{c+\lambda'_{n-k+1-c}}}{t_{c+i-r}}.$$

Hence we must show the equality of these two quantities:

$$\overline{\text{SSYTEqfactor}_{(r,i)}(c)} = 1 - \frac{t_{n+1-c-\lambda'_{n-k+1-c}}}{t_{n+1-c+r-i}} \text{ and}$$

$$\text{edgefactor}_{\underline{x}}(i_{\mu_{i+1-r}}) = 1 - \frac{t_{\text{Man}(\underline{x})}}{t_{\lambda'_{n-k+1-c}^{-i+r+\text{Man}(\underline{x})}}},$$

where \underline{x} is the southern edge of λ in column c .

Now, counting the rows and columns separating x from the southwest corner of $k \times (n - k)$, we have

$$\text{Man}(x) = (n - k - c) + (k - \lambda'_{n-k+1-c} + 1) = n + 1 - c - \lambda'_{n-k+1-c}.$$

Thus, the numerators of the quotients of $\overline{\text{SSYTeqfactor}(c)}$ and $\text{edgefactor}(c)$ are equal. To see that the denominators are also equal, observe

$$\begin{aligned} \text{Man}(x) + \lambda'_{n-k+1-c} - i + r &= (n + 1 - c - \lambda'_{n-k+1-c}) + \lambda'_{n-k+1-c} - i + r \\ &= n + 1 - c - i + r. \end{aligned}$$

(ξ is well-defined and weight-preserving): Let $T \in \mathcal{A}$. We must show $\xi(T)$ is strictly increasing along columns. This is clear since T satisfies (S.3) and (S.4).

Now we show that $\xi(T)$ is weakly increasing along rows. Suppose we have a in position (r, i) and b in position $(r, i + 1)$. This a comes from an i_{μ_i+1-r} in column a in T , while this b comes from an $(i + 1)_{\mu_{i+1}+1-r}$ in column b . By ballotness of T , each i_{μ_i+1-r} must be weakly right of each $(i + 1)_{\mu_{i+1}+1-r}$. Thus $a \leq b$.

Hence $\xi(T)$ is a set-valued semistandard tableau of shape μ' . The same computations showing ξ^{-1} is weight preserving shows $0 \neq \text{wt}(T) = \overline{\text{SSYTeqwt}(\xi(T))}$ and so the desired conclusions hold. \square

The proposition now follows immediately from Lemmas 4.2 and 4.3. \square


5. GOOD TABLEAUX

In this section, we give an intrinsic description of the tableaux that will appear during our generalized *jeu de taquin* process (defined in Section 7). Since we will use *box* labels $\bullet_{\mathcal{G}}$, we distinguish labels i_j as **genetic labels**. As a visual aid, we mark genetic labels \mathcal{F} southeast of a $\bullet_{\mathcal{G}}$ with $\mathcal{F} \prec \mathcal{G}$ as $\mathcal{F}^!$. For a gene \mathcal{G} , let \mathcal{G}^+ (respectively, \mathcal{G}^-) denote the successor (respectively, predecessor) of \mathcal{G} in the total order \prec on genes. For example, $1_1^+ = 2_1$ if $\mu_1 = 1$, and $1_1^+ = 1_2$ if $\mu_1 > 1$. Let \mathcal{G}_{\max} be the maximum gene that can appear, namely $\ell(\mu)_{\mu_{\ell(\mu)}}$ where $\ell(\mu)$ is the number of nonzero rows of μ . Declare $\mathcal{G}_{\max}^+ := (\ell(\mu) + 1)_1$.

A **\mathcal{G} -good tableau** is an edge-labeled filling T of ν/λ by genetic labels i_j (such that $i \in \mathbb{Z}_{>0}$ and the j 's that appear for each i form an initial segment of $\mathbb{Z}_{>0}$) and *box* labels $\bullet_{\mathcal{G}}$, satisfying the conditions (G.1)–(G.13) below:

- (G.1) no genetic label is too high;
- (G.2) no $\bullet_{\mathcal{G}}$ is southeast of another (in particular, $\bullet_{\mathcal{G}}$'s are in distinct rows and columns);
- (G.3) the labels \prec -increase along rows (ignoring any $\bullet_{\mathcal{G}}$'s), except for possibly three consecutive labels $\boxed{\mathcal{H} \mid \bullet_{\mathcal{G}} \mid \mathcal{F}^!}$ with $\mathcal{H} > \mathcal{F}$;
- (G.4) the labels $<$ -increase down columns (ignoring any $\bullet_{\mathcal{G}}$'s), except that unmarked \mathcal{F} may appear adjacent and above $\mathcal{F}^!$ when both are box labels;
- (G.5) if i_j, k_ℓ appear on the same edge, then $i \neq k$;
- (G.6) if i_j is West of i_k , then $j \leq k$;
- (G.7) each edge label is maximally west in its gene;
- (G.8) each genotype G obtained by choosing one label of each gene of T is ballot in the sense defined in Section 1.3.
- (G.9) if \mathcal{F} appears northwest of $\bullet_{\mathcal{G}}$, then $\mathcal{F} \prec \mathcal{G}$;
- (G.10) if $\mathcal{F}^! \in x$ or $\mathcal{F}^! \in \underline{x}$, then $\bullet_{\mathcal{G}}$ appears in x 's row;

- (G.11) $\bullet_{\mathcal{G}}$ does not appear in a column containing a marked label;
(G.12) if ℓ and ℓ' are genetic labels of the same family with ℓ NorthWest of ℓ' , then there are boxes x, z in row r with x West of z , $\ell \in x$ or \bar{x} , and $\ell' \in z$ or \underline{z} ; further, $\bullet_{\mathcal{G}}$ appears in some box y of r that is East of x and west of z . Pictorially, the scenarios are:



Furthermore, if $y = z = x^{\rightarrow}$ in the last scenario, then y^{\rightarrow} does not contain a marked label nor another instance of the gene of ℓ' .

We place a **virtual label** \mathcal{H} on each edge \underline{x} where $\mathcal{H} \in \underline{x}$ would

- (V.1) *not* be marked (hence if \mathcal{H} appears southeast of a $\bullet_{\mathcal{G}}$, then $\mathcal{H} \succeq \mathcal{G}$);
(V.2) *not* be maximally west in its gene (hence violating condition (G.7)); and
(V.3) satisfy the conditions (G.1), (G.4), (G.5), (G.6), (G.8), (G.9) and (G.12).
(G.13) If $\mathcal{E}^! \in x$ or $\mathcal{E}^! \in \underline{x}$, then there is \mathcal{F} or \mathcal{F} on \underline{x} with $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$.

A tableau is **good** if it is \mathcal{G} -good for some \mathcal{G} .

Example 5.1. The tableau

$\frac{1}{2_1}$	\bullet_{2_2}	$\frac{1}{2_2}$


 is 2_2 -good. Although the labels in the second row do not increase left to right, they satisfy (G.3). Furthermore, notice the 1_1 and $1_2^!$ satisfy (G.12), as do the 2_1 and 2_2 .

The tableau

	1_1	\bullet_{2_1}
\bullet_{2_1}	$\frac{1}{2_1}$	

 is also good. Although the label 1_1 appears twice in the same column, the lower instance is marked in accordance with (G.4). □

Example 5.2. The following tableaux are *not* good:



The first fails conditions (G.1) and (G.7) because of the edge label 2_1 . The second fails (G.8), as the unique genotype is not ballot. Although the marked $1_1^!$ in the third tableau has a label of family 2 on the lower edge of its box, the tableau fails (G.13) as $1 = N_{1_1} \neq N_{2_1} = 0$. It also fails (G.11) by having both a \bullet_{2_1} and a marked label in the second column. The fourth tableau fails (G.12). □

Lemma 5.3. *If $T \in \text{Bundled}(\nu/\lambda)$, then T is \mathcal{G} -good for every \mathcal{G} . Moreover the virtual labels of the \mathcal{G} -good tableau T (as defined by (V.1)–(V.3)) are the same as the virtual labels of the bundled tableau T (as defined in Section 2).*

Proof. Since T is bundled, (S.1), (S.2), (S.3) and (S.4) hold. These conditions respectively imply (G.3), (G.4), (G.5) and (G.6). (G.1), (G.7) and (G.8) are part of the definition of a bundled tableau. For (G.12), if ℓ is NorthWest of ℓ' and both are from the same family,

(S.1) or (S.2) is violated. The remaining conditions are vacuous since T has no $\bullet_{\mathcal{G}}$'s. Hence T is \mathcal{G} -good.

The claim about virtual labels is then clear from the definitions. \square

Lemma 5.4 (Strong form of (G.10)). *Assume T is \mathcal{G} -good. Let x be a box of T in row r .*

(I) *If $\mathcal{F}^! \in x$, then $\text{label}(x)$ is marked.*

(II) *If $\mathcal{F}^! \in x$, then there is a y West of x in r such that $\bullet_{\mathcal{G}} \in y$. Every box label of r between x and y is marked.*

Proof. (I): Since $\mathcal{F}^! \in x$, x (and hence also x) is southeast of a $\bullet_{\mathcal{G}}$. By (G.11), $\bullet_{\mathcal{G}} \notin x$. Hence some $\mathcal{E} \in x$. By (G.4), $\mathcal{E} < \mathcal{F}$. Therefore the $\mathcal{E} \in x$ is marked.

(II): Since $\mathcal{F}^! \in x$, there is a $\bullet_{\mathcal{G}}$ northwest of x . By (G.10), there is a $\bullet_{\mathcal{G}}$ in x 's row. If this latter $\bullet_{\mathcal{G}}$ is East of x , these two $\bullet_{\mathcal{G}}$'s are distinct and violate (G.2). Hence the $\bullet_{\mathcal{G}}$ in x 's row is in some box y West of x . If \mathcal{E} is a box label between x and y (and in the same row), it is southeast of the $\text{label}(y) = \bullet_{\mathcal{G}}$. By (G.3) $\mathcal{E} \prec \mathcal{F}$. Hence this \mathcal{E} is also marked. \square

Lemma 5.5 (Strong form of (G.13)). *Let T be \mathcal{G} -good. Suppose $\mathcal{E}^! \in x$ or $\mathcal{E}^! \in \underline{x}$ with $\text{family}(\mathcal{G}) - \text{family}(\mathcal{E}) = k > 0$. For each $0 < h < k$, there is $\mathcal{H}^! \in \underline{x}$ with $N_{\mathcal{H}} = N_{\mathcal{E}}$ and $\text{family}(\mathcal{H}) = \text{family}(\mathcal{E}) + h$. Also, there is a \mathcal{G}' or $(\mathcal{G}') \in \underline{x}$ with $N_{\mathcal{G}'} = N_{\mathcal{E}}$ and $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$.*

Proof. This follows by repeated application of (G.13). Note that none of the \mathcal{H} 's of the statement can be virtual since they must be marked. \square

Lemma 5.6. *If $\mathcal{E} < \mathcal{F}$ are genes of a good tableau T with $N_{\mathcal{E}} = N_{\mathcal{F}}$, then no \mathcal{F} or (\mathcal{F}) is East of any \mathcal{E} .*

Proof. First suppose that some \mathcal{F} is East of some \mathcal{E} . Let G be a genotype of T with $\mathcal{F} \in G$ that is East of some $\mathcal{E} \in G$. Then \mathcal{F} appears before \mathcal{E} in $\text{word}(G)$. By (G.6), the initial segment W of $\text{word}(G)$ ending at \mathcal{F} contains $N_{\mathcal{F}} + 1$ labels of $\text{family}(\mathcal{F})$ and at most $N_{\mathcal{E}}$ labels of $\text{family}(\mathcal{E})$. Thus T 's (G.8) is violated for some $\text{family}(\mathcal{E}) \leq i < \text{family}(\mathcal{F})$, a contradiction. Finally, if some (\mathcal{F}) is East of some \mathcal{E} , then by (V.3) the tableau T' obtained by replacing that (\mathcal{F}) by \mathcal{F} satisfies (G.6) and (G.8). Now we derive the same contradiction as before, using T' in place of T . \square

Lemma 5.7. *If $\mathcal{E}^!$ appears in a good tableau T , then it is maximally west in its gene.*

Proof. Suppose $\mathcal{E}^! \in x$ or $\mathcal{E}^! \in \underline{x}$. By (G.13), there is an \mathcal{F} or $(\mathcal{F}) \in \underline{x}$ with $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\mathcal{E} < \mathcal{F}$. Thus we are done by Lemma 5.6. \square

Lemma 5.8. *Suppose column c of good tableau T contains labels \mathcal{H} and \mathcal{J} with $\mathcal{H} < \mathcal{J}$ and $N_{\mathcal{H}} = N_{\mathcal{J}}$. Then for every i such that $\text{family}(\mathcal{H}) < i < \text{family}(\mathcal{J})$, there is a label \mathcal{I} of family i in column c such that $N_{\mathcal{H}} = N_{\mathcal{I}}$.*

Proof. Suppose not. By (G.8), there is some $\mathcal{I} \in T$ of family i such that $N_{\mathcal{H}} = N_{\mathcal{J}} = N_{\mathcal{I}}$. If this \mathcal{I} is not in column c , we contradict Lemma 5.6. \square

Lemma 5.9. *Suppose \mathcal{E} and \mathcal{F} satisfy $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$. Let T be a \mathcal{G} -good tableau with $(\mathcal{F}) \in \underline{x}$ and either $\mathcal{E}^! \in x$ or $\mathcal{E}^! \in \underline{x}$. Then $\bullet_{\mathcal{G}} \in x^{\leftarrow}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$.*

Proof. If $\bullet_{\mathcal{G}} \notin x^{\leftarrow}$, then by Lemma 5.4, $\mathcal{D}^! \in x^{\leftarrow}$. By (G.3) and (G.4), $\mathcal{D} \prec \mathcal{E}$. Also $\mathcal{E} \prec \mathcal{G}$ since $\mathcal{E}^! \in T$. Thus by (G.6) and Lemma 5.5, there is a $\tilde{\mathcal{E}}^! \in \underline{x^{\leftarrow}}$ or $\tilde{\mathcal{E}}^! \in x^{\leftarrow}$ with $\text{family}(\mathcal{E}) = \text{family}(\tilde{\mathcal{E}})$ and $N_{\tilde{\mathcal{E}}} = N_{\mathcal{D}}$. By (G.13), there is $\tilde{\mathcal{F}}$ or $(\tilde{\mathcal{F}}) \in \underline{x^{\leftarrow}}$ with $\text{family}(\mathcal{F}) = \text{family}(\tilde{\mathcal{F}})$ and $N_{\tilde{\mathcal{E}}} = N_{\tilde{\mathcal{F}}}$. Thus, by Lemma 5.6, $\mathcal{F} \neq \tilde{\mathcal{F}}$, contradicting $(\mathcal{F}) \in \underline{x}$. Finally, $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$ by Lemma 5.5. \square

Lemma 5.10. *If T is \mathcal{G} -good, then no \mathcal{H} is southEast of another.*

Proof. If some \mathcal{H} is SouthEast of another \mathcal{H} , by (G.12) there is a $\bullet_{\mathcal{G}}$ in between the two \mathcal{H} 's. If two \mathcal{H} 's are box labels of the same row, then by (G.3) we reach the same conclusion that there is a $\bullet_{\mathcal{G}}$ in between the two \mathcal{H} 's. In either case, since this $\bullet_{\mathcal{G}}$ is southeast of the western \mathcal{H} we have $\mathcal{H} \prec \mathcal{G}$ by (G.9). Since this $\bullet_{\mathcal{G}}$ is northwest of the eastern \mathcal{H} , this eastern \mathcal{H} is marked. This contradicts Lemma 5.7. Finally, suppose two \mathcal{H} 's are edge labels on the bottom of the same row. This contradicts (G.7). \square

Lemma 5.11. *Let T be a \mathcal{G} -good tableau. Suppose $\text{family}(\mathcal{F}) \leq \text{family}(\mathcal{G})$, $\bullet_{\mathcal{G}} \in y$ and $\mathcal{F} \in z$ or \underline{z} . Then z is not SouthEast of y .*

Proof. Suppose z is SouthEast of y . First assume $\mathcal{F} < \mathcal{G}$. Consider the box a that is in y 's column and z 's row. By Lemma 5.4, either a contains a marked label (contradicting (G.11)) or $\bullet_{\mathcal{G}} \in a$ Southeast of y (contradicting (G.2)).

Now assume $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$. (We do not assume $\mathcal{F} \preceq \mathcal{G}$.) Consider the box b of T that is in y 's row and z 's column. By (G.2), b contains a genetic label. By (G.4), $\text{label}(b) < \mathcal{F}$. Hence $\text{label}(b)$ is marked in T . By Lemma 5.5, b then contains a (possibly virtual) label of the same family as \mathcal{F} and \mathcal{G} . This contradicts (G.4). \square

Lemma 5.12. *Let U be a \mathcal{G}^+ -good tableau. Suppose that $\bullet_{\mathcal{G}^+} \in x$ and that either $\mathcal{G} \in y$ or $\mathcal{G} \in \underline{y}$. Then y is not NorthWest of x .*

Proof. Suppose otherwise. Consider the box b that is in y 's column and x 's row. By (G.2) it contains a genetic label. By (G.4) either $\text{label}(b) > \mathcal{G}$ or else $\mathcal{G}^! \in b$. If $\mathcal{G}^! \in b$, then b is southeast of a $\bullet_{\mathcal{G}^+}$ by definition. This contradicts (G.2). If $\mathcal{G} < \text{label}(b)$, we contradict (G.9). \square

Lemma 5.13. *Let c be a column of a \mathcal{G} -good tableau T . Suppose $\bullet_{\mathcal{G}} \in c$ and either $\mathcal{G} \in c$ or $(\mathcal{G}) \in c$. Further suppose that $\mathcal{E}^! \in y$, where y is a box of column c^{\rightarrow} . Then $(\mathcal{G}) \in \underline{y}$.*

Proof. Since $\mathcal{E}^!$ appears in T , $\mathcal{E} \prec \mathcal{G}$. Since \mathcal{E} appears East of some \mathcal{G} , by (G.6) this implies $\mathcal{E} < \mathcal{G}$.

Hence by Lemma 5.5, there is either $\mathcal{G}' \in \underline{y}$ or $(\mathcal{G}') \in \underline{y}$ with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$. It remains to show $\mathcal{G}' = \mathcal{G}$, for then by (G.7), $(\mathcal{G}) \in \underline{y}$.

Suppose $\mathcal{G}' \neq \mathcal{G}$. Then by (G.4), (G.5) and (G.6), $\mathcal{G}' = \mathcal{G}^+$. By Lemma 5.5, $N_{\mathcal{E}} = N_{\mathcal{G}^+}$; thus $\text{family}(\mathcal{E}^-) = \text{family}(\mathcal{E})$ by (G.8). Also by (G.8), every instance of \mathcal{E}^- must be read before any \mathcal{G} or (\mathcal{G}) . By (G.4), $\mathcal{E}^- \notin c^{\rightarrow}$. By (G.6), \mathcal{E}^- does not appear East of c^{\rightarrow} . But by assumption either $\mathcal{G} \in c$ or $(\mathcal{G}) \in c$, so \mathcal{E}^- must appear in c .

Consider the box y^{\leftarrow} . By Lemma 5.4, either $\bullet_{\mathcal{G}} \in y^{\leftarrow}$ or some $\mathcal{D}^! \in y^{\leftarrow}$. The latter is impossible by (G.11), since $\bullet_{\mathcal{G}} \in c$. Hence $\bullet_{\mathcal{G}} \in y^{\leftarrow}$.

Now \mathcal{E}^- cannot appear South of y^{\leftarrow} in c , for then it would be marked, in violation of (G.11). We have $\mathcal{E}^- \notin y^{\leftarrow}$, since $\bullet_{\mathcal{G}} \in y^{\leftarrow}$. By (G.12), \mathcal{E}^- cannot appear North of y^{\leftarrow} in c . This contradicts that \mathcal{E}^- must appear in c , and therefore the assumption $\mathcal{G}' \neq \mathcal{G}$. \square

6. SNAKES OF GOOD TABLEAUX

In this section, we give structural results about certain subsets of a good tableau; these will play a critical role in the definition of our generalized *jeu de taquin* (given in Section 7).

6.1. Snakes. Let T be a \mathcal{G} -good tableau. Let $\mathcal{G} = g_k$ and consider the set of *boxes* in T that contain either $\bullet_{\mathcal{G}}$ or \mathcal{G} . This set decomposes into edge-connected components R that we call **presnakes**. A **short ribbon** is a connected skew shape without a 2×2 subshape and where each row and column contains at most two boxes.

Lemma 6.1. *Each presnake R is a short ribbon. Any row of R with two boxes is $\boxed{\bullet_{\mathcal{G}} \mathcal{G}}$. Any column of R with two boxes is $\begin{array}{c} \boxed{\bullet_{\mathcal{G}}} \\ \boxed{\mathcal{G}} \end{array}$.*


Proof. Since T is \mathcal{G} -good, there is no $\mathcal{G}^!$. So any column of R has at most one \mathcal{G} by (G.4) and at most one $\bullet_{\mathcal{G}}$ by (G.2). Hence any column of R has at most two boxes. By (G.9) if $\bullet_{\mathcal{G}}$ and \mathcal{G} are in the same column, the $\bullet_{\mathcal{G}}$ is to the north. The description of rows of R holds by (G.2), (G.3) and (G.9). That R is a skew shape with no 2×2 subshape then follows immediately. \square

A **snake** S is a presnake R extended by (R.1)–(R.3):

- (R.1) If the box immediately right of the northmost $\bullet_{\mathcal{G}}$ in R contains \mathcal{G}^+ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, then adjoin this box to R .
- (R.2) If the box immediately left of the southmost \mathcal{G} in R contains a marked label, adjoin this box to R .
- (R.3) If x in the northmost row of R contains $\bullet_{\mathcal{G}}$, $\text{label}(x^{\rightarrow})$ is marked and either \mathcal{G} or $\mathcal{G} \in x^{\rightarrow}$, then adjoin x^{\rightarrow} to R .

The entries of S are its box labels and labels appearing on the bottom edges of its boxes.

Example 6.2. Below are snakes for $\mathcal{G} = 2_2$:



On the other hand, $\begin{array}{c} \bullet_{2_2} \\ \bullet_{2_2} \end{array} \begin{array}{c} 2_2 \\ 2_2 \end{array}$ is *not* a snake, even if $3_1 = 2_2^+$ ((R.1) does not apply). \square

Example 6.3 (Snakes can share a row). $\begin{array}{ccc|c} & & & 1_2 \\ \bullet_{2_2} & 1_1! & 2_2 & \\ \hline & 2_1, 3_1 & 3_2 & \end{array}$ contains two snakes as colored. \square

Example 6.4 (Snakes can share a column). $\begin{array}{c|cc} & \bullet_{1_1} & 1_2 \\ \bullet_{1_1} & 1_2 & \\ \hline 1_1 & & \end{array}$ has two snakes as colored. \square

Lemma 6.5. *Every snake S is a short ribbon.*

Proof. S is built by adjoining boxes to a presnake R . By Lemma 6.1, R is a short ribbon. In view of Lemma 6.1, (R.1) and (R.3) only apply if the northmost row of R is a single box with $\bullet_{\mathcal{G}}$. So adjoining a box to the right maintains shortness. Similarly, (R.2) maintains shortness. \square

Lemma 6.6 (Disjointness and relative positioning of snakes). *Suppose S, S' are snakes obtained from distinct presnakes R, R' respectively. Up to relabeling of the snakes, one of the following holds:*

- (I) S is entirely SouthWest of the S' (that is, if b, b' are respectively boxes of these snakes, then b is SouthWest of b').
- (II) S consists of a single box containing $\bullet_{\mathcal{G}}$ with neither \mathcal{G} nor $\textcircled{\mathcal{G}}$ on its lower edge; further, this box appears West of and in the same row as the southmost row of S' , and all intervening box labels are marked; cf. Example 6.3.
- (III) S involves an (R.1) extension, adjoining a \mathcal{G}^+ in some box w , while $S' = \{\bullet_{\mathcal{G}} \in w^\uparrow\}$ or $S' = \{\bullet_{\mathcal{G}} \in w^\uparrow, \mathcal{G}^+ \in w^{\uparrow\rightarrow}\}$; cf. Example 6.4.

In particular, S and S' are box disjoint.

Proof. By Lemma 6.1, (G.2) and/or (G.4), R and R' share at most one row and do not share a column. Moreover, one sees that R is southWest of R' (say). By (R.1)–(R.3), S and S' share a row if and only if R and R' do.

Case 1: (R and R' share a row r): The northmost row of R and the southmost row of R' are in row r . We must show that (II) holds and that S, S' are box disjoint.

By (G.2), (G.9) and Lemma 5.10, R has in row r only a $\bullet_{\mathcal{G}} \in x$ while R' has in r only $\mathcal{G} \in y$. Since $S \neq S'$, $y \neq x^\rightarrow$. By (G.3), $\text{label}(y^\leftarrow) \prec \mathcal{G}$, so we have some marked label $\mathcal{F}^\dagger \in y^\leftarrow$. Therefore R' extends to S' by (R.2).

Claim 6.7. *No \mathcal{G} or $\textcircled{\mathcal{G}}$ appears in columns west of y^\leftarrow .*

Proof. Since $\mathcal{F} \prec \mathcal{G}$, we are done by (G.4) and (G.6) if $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$. Thus assume $\mathcal{F} < \mathcal{G}$. By Lemma 5.5, there is either $\mathcal{G}' \in y^\leftarrow$ or $\textcircled{\mathcal{G}'} \in y^\leftarrow$ such that $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$ and $N_{\mathcal{F}} = N_{\mathcal{G}'}$. By (G.6), $\mathcal{G}' \preceq \mathcal{G}$ because $\mathcal{G} \in y$. If $\mathcal{G}' = \mathcal{G}$, then since $N_{\mathcal{F}} = N_{\mathcal{G}'(=\mathcal{G})}$, the $\mathcal{G} \in y$ and $\mathcal{F}^\dagger \in y^\leftarrow$ combine to contradict Lemma 5.6. Thus $\mathcal{G}' \prec \mathcal{G}$ and we are done by (G.6) and (G.4). \square

By Claim 6.7, $R = \{\bullet_{\mathcal{G}} \in x\}$ without \mathcal{G} or $\textcircled{\mathcal{G}} \in x$. Observe that R cannot extend to S by (R.1), since (R.1) requires $\mathcal{G}^+ \in x^\rightarrow$, which contradicts (G.3) in view of $\mathcal{G} \in y$. It cannot be extended by (R.2) since $\mathcal{G} \notin x$. If R were extended by (R.3), there would be a \mathcal{G} or $\textcircled{\mathcal{G}}$ in x^\rightarrow in violation of Claim 6.7. Thus $R = S = \{x\}$.

By Lemma 5.4(II), all labels strictly between x and y are marked. Hence (II) holds. Since $y^\leftarrow \notin S$, we see by (R.1)–(R.3) that S and S' are box disjoint.

Case 2: (R and R' do not share a row): We may assume S and S' share a column, for if they do not, then clearly (I) and box-disjointness both hold. Since R and R' do not share a column, S and S' can only share a column if R is extended East by (R.1) or (R.3) or if R' is extended West by (R.2). Let x be the northeastmost box of R and y be the southwestmost box of R' .

Subcase 2.1: (R is extended by (R.1)): Since $\text{label}(x^\rightarrow) = \mathcal{G}^+$ and $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, by (G.6) R' cannot contain any \mathcal{G} 's and therefore $R' = \{\bullet_{\mathcal{G}} \in y\}$. Hence (R.2) does not extend R' . By assumption, x^\rightarrow and y are in the same column. Hence by (G.4) and (G.11), $y = x^{\rightarrow\uparrow}$. By (G.6), R' is not extended by (R.3), since $\mathcal{G}^+ \in x^\rightarrow$ and (R.3) requires $\mathcal{G} \in \underline{y}$ or $\textcircled{\mathcal{G}} \in \underline{y}$. If R' is extended by (R.1), we obtain the second scenario described by (III) (and S, S' are box disjoint). If R' is not extended by any of (R.1)–(R.3), then we have the first scenario described by (III) (and S, S' are box disjoint).

Subcase 2.2: (R is extended by (R.3)): Let c be x^\rightarrow 's column. We have $\mathcal{F}^! \in x^\rightarrow$ and either $\mathcal{G} \in \underline{x^\rightarrow}$ or $\textcircled{\mathcal{G}} \in \underline{x^\rightarrow}$. Moreover $N_{\mathcal{F}} = N_{\mathcal{G}}$. Hence by Lemma 5.6, no \mathcal{G} appears East of c . Thus $R' = \{\bullet_{\mathcal{G}} \in y\}$. By (G.11), $y \notin c$. Thus S and R' do not share a column. Since $\bullet_{\mathcal{G}} \in y$, R' is not extended by (R.2). Thus S and S' do not share a column.

Subcase 2.3: (R' is extended by (R.2); R is not extended by either (R.1) or (R.3)): Here $\mathcal{G} \in y$ and $\mathcal{F}^! \in y^\leftarrow$. By Lemma 5.5, either $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$ or else we have $\mathcal{G}' \in \underline{y^\leftarrow}$ or $\textcircled{\mathcal{G}'} \in \underline{y^\leftarrow}$ such that $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$. Hence by (G.4) and (G.11), R cannot contain a box in the column of y^\leftarrow . Hence R, S' do not share a column. Hence by the assumption of the subcase, S and S' do not share a column. \square

6.2. Snake sections. We decompose each snake S into three **snake sections** denoted $\text{head}(S)$, $\text{body}(S)$ and $\text{tail}(S)$ as follows:

Definition-Lemma 6.8.

- (I) If a snake S has at least two rows and its southmost row has two boxes, then $\text{head}(S)$ is the southmost row of S , $\text{tail}(S)$ is the northmost row and $\text{body}(S)$ is the remaining rows.
- (II) If a snake S has at least two rows and its southmost row has exactly one box, then $\text{head}(S)$ is empty, $\text{tail}(S)$ is the northmost row and $\text{body}(S)$ is the other rows.
- (III) If S has exactly one row, then S is one of the following (edge labels not depicted):

- (i) $S = \boxed{\mathcal{G}} = \text{body}(S)$; (ii) $S = \boxed{\bullet_{\mathcal{G}}} = \text{head}(S)$; (iii) $S = \boxed{\bullet_{\mathcal{G}} \mathcal{G}} = \text{head}(S)$;
- (iv) $R = \boxed{\bullet_{\mathcal{G}} \mathcal{G}^+} = \text{head}(S)$; (v) $S = \boxed{\mathcal{F}^! \mathcal{G}} = \text{head}(S)$;
- (vi) $S = \boxed{\bullet_{\mathcal{G}} \mathcal{F}^!} = \text{tail}(S)$ (with \mathcal{G} or $\textcircled{\mathcal{G}}$ on the lower right edge).

Proof. It is only required to verify that in (III) all possible one-row snakes are shown. This is done by combining Lemma 6.1 and (R.1)–(R.3). \square

Lemma 6.9 (Properties of head, body, tail).

- (I) If $\text{head}(S) = \{x\}$, then $\bullet_{\mathcal{G}} \in x$.
- (II) If $\text{head}(S) = \{x, x^\rightarrow\}$, then $\text{head}(S) = \boxed{\mathcal{F}^! \mathcal{G}}, \boxed{\bullet_{\mathcal{G}} \mathcal{G}}$ or $\boxed{\bullet_{\mathcal{G}} \mathcal{G}^+}$.
- (III) $\text{body}(S)$ is a short ribbon consisting only of $\bullet_{\mathcal{G}}$'s and \mathcal{G} 's (with no edge label \mathcal{G} 's or $\textcircled{\mathcal{G}}$'s).
- (IV) If $\text{tail}(S) = \{x\}$, then $\text{tail}(S) = \boxed{\bullet_{\mathcal{G}}}$ and S has at least two rows.
- (V) If $\text{tail}(S) = \{x, x^\rightarrow\} = \boxed{\bullet_{\mathcal{G}} \mathcal{G}}$ or $\boxed{\bullet_{\mathcal{G}} \mathcal{G}^+}$, then S has at least two rows, $\mathcal{G} \notin \underline{x}$ and $\textcircled{\mathcal{G}} \notin \underline{x}$.
- (VI) If $\text{tail}(S) = \{x, x^\rightarrow\}$ and \mathcal{G} or $\textcircled{\mathcal{G}} \in \underline{x^\rightarrow}$, then $\text{tail}(S) = \boxed{\bullet_{\mathcal{G}} \mathcal{F}^!}$ or $\boxed{\bullet_{\mathcal{G}} \textcircled{\mathcal{F}^!}}$.
- (VII) If S has at least two rows, then $\mathcal{G} \in \underline{x^\downarrow}$ where x is the westmost box of $\text{tail}(S)$.

Proof. If S has one row, then by Definition-Lemma 6.8(III) these claims are clear (or irrelevant). Thus assume S has at least two rows.

(I): Under the assumption that S has at least two rows, the claim is vacuous since by Definition-Lemma 6.8(I,II) we know $|\text{head}(S)| \neq 1$.

(II): Either the southmost row of S is $\boxed{\mathcal{F}! \mathcal{G}}$ if (R.2) was used, or it is $\boxed{\bullet \mathcal{G}}$ if (R.2) was not used; cf. Lemma 6.1.

(III): That $\text{body}(S)$ is a short ribbon is clear, since S is a short ribbon by Lemma 6.5. Boxes of $\text{body}(S)$ only contain \mathcal{G} or $\bullet_{\mathcal{G}}$ because (R.1)–(R.3) adjoin boxes only to the northmost or southmost row (and if the southmost row of S has two boxes, then by definition that row is not part of $\text{body}(S)$). By (G.12), an edge label \mathcal{G} or $\textcircled{\mathcal{G}}$ can only appear in the northmost or southmost row of S . In those cases, the row is not part of $\text{body}(S)$ by Definition-Lemma 6.8(I,II).

(IV): $\text{tail}(S)$ is the northmost row of S and, since $|\text{tail}(S)| = 1$, it is the northmost row of the presnake of S . Thus we are done by Lemma 6.1.

(V): $\text{tail}(S)$ is the northmost row and by Lemma 6.1, $\mathcal{G} \in x^\downarrow$ (x^\downarrow is in the presnake of S) so $\mathcal{G}, \textcircled{\mathcal{G}} \notin x$ by (G.4).

(VI): x is in the presnake of S and so by Lemma 6.1, $\bullet_{\mathcal{G}} \in x$. By (G.2), $\bullet_{\mathcal{G}} \notin x^\rightarrow$. By (G.4), $\text{label}(x^\rightarrow) < \mathcal{G}$ and so $\text{label}(x^\rightarrow)$ is marked, since it is southeast of the $\bullet_{\mathcal{G}} \in x$.

(VII): x and x^\downarrow are part of the presnake of S . Now apply Lemma 6.1. □

7. GENOMIC JEU DE TAQUIN

7.1. Miniswaps. We first define **miniswaps** on snake sections of a \mathcal{G} -good tableau. The output is a formal sum of tableaux. Below, interpret $\bullet = \bullet_{\mathcal{G}}$ before the miniswap and $\bullet = \bullet_{\mathcal{G}^+}$ after the miniswap. We depict $\textcircled{\mathcal{G}}$ whenever it exists. Labels and virtual labels from other genes are not depicted unless relevant to the miniswap's definition. For a box x , define

$$\beta(x) := 1 - \frac{t_{\text{Man}(x)}}{t_{\text{Man}(x)+1}} \quad \text{and} \quad \hat{\beta}(x) := 1 - \beta(x) = \frac{t_{\text{Man}(x)}}{t_{\text{Man}(x)+1}}.$$

Note that if $x = \alpha/\beta$, then $\hat{\beta}(x) = \text{wt } \alpha/\beta$, as defined in Section 3. If a snake section is empty, then mswap acts trivially, so below we assume otherwise.

7.1.1. Miniswaps on $\text{head}(S)$.

(Case H1: $\text{head}(S) = \{x\}$ and $\mathcal{G} \in x$):

$$\text{head}(S) = \boxed{\bullet_{\mathcal{G}}} \mapsto \text{mswap}(\text{head}(S)) = \beta(x) \cdot \boxed{\mathcal{G}} + \gamma \cdot \boxed{\bullet_{\mathcal{G}}}$$

Set $\gamma := 0$ if $\text{row}(x) = \text{family}(\mathcal{G})$ (that is, if $\mathcal{G} \in \bar{x}$ would be too high); otherwise set $\gamma := 1$.

(Case H2: $\text{head}(S) = \{x\}$ and $\textcircled{\mathcal{G}} \in x$):

$$\text{head}(S) = \boxed{\bullet_{\textcircled{\mathcal{G}}}} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\bullet} + \beta(x) \cdot \boxed{\mathcal{G}}$$

(Case H3: $\text{head}(S) = \{x\}$ and Cases H1/H2 do not apply):

$$\text{head}(S) = \boxed{\bullet} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\bullet}$$

(Case H4: $\text{head}(S) = \{x, x^\rightarrow\}$, $\mathcal{G} \in x^\rightarrow$, and $\mathcal{G} \in \underline{x}$):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}} \mapsto \text{mswap}(\text{head}(S)) = 0$$

(Case H5: $\text{head}(S) = \{x, x^\rightarrow\}$, $\mathcal{G} \in x^\rightarrow$, and $\mathcal{G} \notin \underline{x}$):

(Subcase H5.1: $\mathcal{H} \in \underline{x^\rightarrow}$, $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}}_{\mathcal{H}} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\bullet \mathcal{G}^!}_{\mathcal{H}}$$

(Subcase H5.2: $\mathcal{H} \in \underline{x^\rightarrow}$, $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}}_{\mathcal{H}} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\bullet \mathcal{G}^!}_{\mathcal{H}} + \hat{\beta}(x) \cdot \boxed{\mathcal{G} \bullet}$$

(Subcase H5.3: Subcases H5.1/H5.2 do not apply):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}}_{\mathcal{G}} \text{ or } \boxed{\bullet \mathcal{G}} \mapsto \text{mswap}(\text{head}(S)) = \hat{\beta}(x) \cdot \boxed{\mathcal{G} \bullet}$$

(Case H6: $\text{head}(S) = \{x, x^\rightarrow\}$, $\mathcal{G}^+ \in x^\rightarrow$, and $\mathcal{G} \in \underline{x}$):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}^+}_{\mathcal{G}} \mapsto \text{mswap}(\text{head}(S)) = \beta(x) \cdot \boxed{\mathcal{G} \mathcal{G}^+} + \alpha \cdot \boxed{\mathcal{G} \bullet}_{\mathcal{G}^+}$$

Set $\alpha := 0$ if the second tableau has two $\bullet_{\mathcal{G}^+}$'s in the same column; otherwise set $\alpha := \hat{\beta}(x)$.

(Case H7: $\text{head}(S) = \{x, x^\rightarrow\}$, $\mathcal{G}^+ \in x^\rightarrow$, and $\mathcal{G} \in \underline{x}$):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}^+}_{\mathcal{G}} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\bullet \mathcal{G}^+} + \beta(x) \cdot \boxed{\mathcal{G} \mathcal{G}^+} + \alpha \cdot \boxed{\mathcal{G} \bullet}_{\mathcal{G}^+}$$

Set $\alpha := 0$ if the third tableau has two $\bullet_{\mathcal{G}^+}$'s in the same column; otherwise set $\alpha := \hat{\beta}(x)$.

(Case H8: $\text{head}(S) = \{x, x^\rightarrow\}$, $\mathcal{G}^+ \in x^\rightarrow$, and Cases H6 and H7 do not apply):

$$\text{head}(S) = \boxed{\bullet \mathcal{G}^+} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\bullet \mathcal{G}^+}$$

(Case H9: $\text{head}(S) = \{x, x^\rightarrow\}$, $\mathcal{F}^! \in x$, and $\mathcal{G} \in x^\rightarrow$):

$$\text{head}(S) = \boxed{\mathcal{F}^! \mathcal{G}} \mapsto \text{mswap}(\text{head}(S)) = \boxed{\mathcal{F}^! \mathcal{G}^!}$$

Lemma 7.1. *Every nonempty $\text{head}(S)$ falls into exactly one of H1–H9.*

Proof. Since $\text{head}(S) \neq \emptyset$, $|\text{head}(S)| \in \{1, 2\}$ by Lemma 6.5. If $\text{head}(S) = \{x\}$, then by Lemma 6.9(I), $\bullet_{\mathcal{G}} \in x$. Then \underline{x} contains exactly one of \mathcal{G} , \mathcal{G} or neither; these are respectively Cases H1, H2 and H3. If $\text{head}(S) = \{x, x^\rightarrow\}$, see Lemma 6.9(II): one possibility is $\mathcal{F}^! \in x$ and $\mathcal{G} \in x^\rightarrow$; this is H9. Otherwise, $\bullet_{\mathcal{G}} \in x$ and x^\rightarrow contains \mathcal{G} or \mathcal{G}^+ . The cases where $\mathcal{G} \in x^\rightarrow$ are covered by H4–H5. The cases where $\mathcal{G}^+ \in x^\rightarrow$ are covered by H6–H8. \square

7.1.2. *Miniswaps on $\text{body}(S)$.* Let $\text{body}_{\bullet_{\mathcal{G}}}(S) = \{x \in \text{body}(S) : \bullet_{\mathcal{G}} \in x\}$.

(Case B1: $\text{body}(S) = S$): By Definition-Lemma 6.8, $S = \boxed{\mathcal{G}}$. Define

$$\text{body}(S) = \boxed{\mathcal{G}} \mapsto \text{mswap}(\text{body}(S)) = \boxed{\mathcal{G}}$$

(Case B2: The southmost row of $\text{body}(S)$ contains two boxes): Replace each \mathcal{G} in $\text{body}(S)$ with $\bullet_{\mathcal{G}^+}$ and each $\bullet_{\mathcal{G}}$ with \mathcal{G} , emitting a weight $\prod_{x \in \text{body}_{\bullet_{\mathcal{G}}}(S)} \hat{\beta}(x)$. That is (cf. Lemma 6.9(III)),

$$\text{body}(S) = \begin{array}{c} \bullet \mathcal{G} \\ \bullet \mathcal{G} \\ \bullet \mathcal{G} \end{array} \mapsto \text{mswap}(\text{body}(S)) = \prod_{x \in \text{body}_{\bullet_{\mathcal{G}}}(S)} \hat{\beta}(x) \cdot \begin{array}{c} \mathcal{G} \bullet \\ \mathcal{G} \bullet \\ \mathcal{G} \bullet \end{array}$$

(Case B3: Cases B1/B2 do not apply): Replace each \mathcal{G} in $\text{body}(S)$ with $\bullet_{\mathcal{G}^+}$ and each $\bullet_{\mathcal{G}}$ with \mathcal{G} , emitting $-\prod_{x \in \text{body}_{\bullet_{\mathcal{G}}}(S)} \hat{\beta}(x)$. That is (cf. Lemma 6.9(III)),

$$\text{body}(S) = \begin{array}{|c|c|} \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \bullet \\ \hline \end{array} \mapsto \text{mswap}(\text{body}(S)) = -\prod_{x \in \text{body}_{\bullet_{\mathcal{G}}}(S)} \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{G} & \bullet \\ \hline \mathcal{G} & \bullet \\ \hline \end{array}$$

Lemma 7.2. *Every nonempty $\text{body}(S)$ falls into exactly one of B1–B3.*

Proof. If B1 applies, then by Definition-Lemma 6.8, $S = \boxed{\mathcal{G}}$. The lemma follows. \square

7.1.3. *Miniswaps on $\text{tail}(S)$.*

(Case T1: $\text{tail}(S) = \{x\}$):

$$\text{tail}(S) = \boxed{\bullet} \mapsto \text{mswap}(\text{tail}(S)) = -\hat{\beta}(x) \cdot \boxed{\mathcal{G}}$$

(Case T2: $\text{tail}(S) = \{x, x^{\rightarrow}\}$ and $\mathcal{G} \in x^{\rightarrow}$):

$$\text{tail}(S) = \boxed{\bullet \mathcal{G}} \mapsto \text{mswap}(\text{tail}(S)) = \hat{\beta}(x) \cdot \boxed{\mathcal{G} \bullet}$$

(Case T3: $\text{tail}(S) = \{x, x^{\rightarrow}\}$ and $\mathcal{G}^+ \in x^{\rightarrow}$):

$$\text{tail}(S) = \boxed{\bullet \mathcal{G}^+} \mapsto \text{mswap}(\text{tail}(S)) = -\hat{\beta}(x) \cdot \boxed{\mathcal{G} \mathcal{G}^+} + \alpha \cdot \boxed{\mathcal{G} \bullet_{\mathcal{G}^+}}$$

Set $\alpha := 0$ if the second tableau has two $\bullet_{\mathcal{G}^+}$'s in the same column; otherwise set $\alpha := \hat{\beta}(x)$.

(Case T4: $\text{tail}(S) = \{x, x^{\rightarrow}\}$, $\mathcal{G} \in x^{\rightarrow}$): Let $Z = \{\ell \in x^{\rightarrow} : \mathcal{F} \prec \ell \prec \mathcal{G}\}$.

(Subcase T4.1: $\mathcal{H} \in x^{\rightarrow}$, $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$):

$$\text{tail}(S) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline z, \mathcal{G}, \mathcal{H} & \\ \hline \end{array} \mapsto \text{mswap}(\text{tail}(S)) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline z, \mathcal{G}^!, \mathcal{H} & \\ \hline \end{array}$$

(Subcase T4.2: $\mathcal{H} \in x^{\rightarrow}$, $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$):

$$\text{tail}(S) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline z, \mathcal{G}, \mathcal{H} & \\ \hline \end{array} \mapsto \text{mswap}(\text{tail}(S)) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline z, \mathcal{G}^!, \mathcal{H} & \\ \hline \end{array} + \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{F}, Z & \\ \hline \mathcal{G} & \bullet \\ \hline \end{array}$$

(Subcase T4.3: Subcases T4.1/T4.2 do not apply):

$$\text{tail}(S) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline z, \mathcal{G} & \\ \hline \end{array} \mapsto \text{mswap}(\text{tail}(S)) = \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{F}, Z & \\ \hline \mathcal{G} & \bullet \\ \hline \end{array}$$

(Case T5: $\text{tail}(S) = \{x, x^{\rightarrow}\}$, $\mathcal{G} \in x^{\rightarrow}$, $\mathcal{G} \notin x$): Let $Z = \{\ell \in x^{\rightarrow} : \mathcal{F} \prec \ell \prec \mathcal{G}\}$.

$$\text{tail}(S) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline z, \mathcal{G} & \\ \hline \end{array} \mapsto \text{mswap}(\text{tail}(S)) = \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{F}, Z & \\ \hline \mathcal{G} & \bullet \\ \hline \end{array}$$

(Case T6: $\text{tail}(S) = \{x, x^{\rightarrow}\}$, $\mathcal{G} \in x^{\rightarrow}$, $\mathcal{G} \in x$):

$$\text{tail}(S) = \begin{array}{|c|c|} \hline \bullet & \mathcal{F}^! \\ \hline \mathcal{G} & \mathcal{G} \\ \hline \end{array} \mapsto \text{mswap}(\text{tail}(S)) = 0$$

Lemma 7.3. *Every nonempty $\text{tail}(S)$ falls into exactly one of T1–T6.*

Proof. Since $\text{tail}(S) \neq \emptyset$, $|\text{tail}(S)| \in \{1, 2\}$ by Lemma 6.5. If $|\text{tail}(S)| = 1$, then by Lemma 6.9(IV), $\text{tail}(S) = \boxed{\bullet_{\mathcal{G}}}$; this is covered by T1. Suppose $\text{tail}(S) = \{x, x^{\rightarrow}\}$. By Lemma 6.1, (R.1)–(R.3) and Definition-Lemma 6.8, $\text{tail}(S) = \boxed{\bullet_{\mathcal{G}} \mathcal{G}}$ (handled by T2),

$\text{tail}(S) = \boxed{\bullet \mathcal{G}^+}$ (handled by T3) or $\text{tail}(S) = \boxed{\bullet \mathcal{F}^!}$ with \mathcal{G} or $\mathcal{G} \in \underline{x}^{\rightarrow}$ (handled by T4, T5 or T6). \square

7.2. Swaps and slides. We define $\text{swap}_{\mathcal{G}}(T)$ and $\text{slide}_{\{x_i\}}(T)$ for a good tableau T . Define $\text{swap}_{\mathcal{G}}$ on a single snake S by applying mswap simultaneously to $\text{head}(S)$, $\text{body}(S)$, and $\text{tail}(S)$ (where the conditions on each mswap refer to the original S).

Lemma 7.4. *On the edges shared by two adjacent snake sections, the modifications to the labels given by the two miniswaps are compatible.*

Proof. Suppose the lower of the two adjacent sections is $\text{head}(S)$. The only miniswap that introduces a label to the northeast edge (i.e., \bar{x} if $\text{head}(S) = \{x\}$ or \bar{x}^{\rightarrow} if $\text{head}(S) = \{x, x^{\rightarrow}\}$) is H1. However in that case $\text{head}(S) = S$ and the compatibility issue is moot. Since body miniswaps do not affect edge labels, the remaining check is when a tail miniswap involves \underline{x} where x is the left box of $\text{tail}(S)$. This only occurs in T6. In this case $\text{tail}(S) = S$, so compatibility is again moot. \square

Lemma 7.5 (Swap commutation). *If S_1, S_2 are distinct snakes in a \mathcal{G} -good tableau T , then applying $\text{swap}_{\mathcal{G}}$ to S_1 commutes with applying $\text{swap}_{\mathcal{G}}$ to S_2 .*

Proof. By definition, the locations of virtual labels in one snake are unaffected by swapping another snake. Hence if the snakes do not share a horizontal edge, there is no concern. If they do, this is the situation of Lemma 6.6(III). The northmost row r of the lower snake (say S_1) is $\{x, x^{\rightarrow}\}$ with $\mathcal{G}^+ \in x^{\rightarrow}$. Hence by (G.4), $\mathcal{G}, \mathcal{G} \notin \bar{x}^{\rightarrow}$. By inspection, no miniswap involving r affects \bar{x}^{\rightarrow} . Now, the upper snake S_2 has a single row, which by the previous sentence is either an H3 or H8 head, irregardless of whether we have acted on r already. Therefore, $\text{swap}_{\mathcal{G}}$ acts trivially on S_2 whether we act on S_1 or S_2 first. \square

Lemma 7.5 permits us to define the **swap** operation $\text{swap}_{\mathcal{G}}$ on a \mathcal{G} -good tableau as the result of applying $\text{swap}_{\mathcal{G}}$ to all snakes (in arbitrary order). Extend $\text{swap}_{\mathcal{G}}$ to a $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ -linear operator.

An **inner corner** of ν/λ is a maximally southeast box of λ . An **outer corner** of ν/λ is a maximally southeast box of ν/λ .

Let $T \in \text{Bundled}(\nu/\lambda)$ and $\{x_i\}$ be a subset of the inner corners of ν/λ . Define $T^{(1_1)}$ to be T with \bullet_{1_1} placed in each x_i .

Lemma 7.6. *Each $T^{(1_1)}$ is 1_1 -good.*

Proof. (G.2) is clear. By Lemma 5.3, T is good; (G.1), (G.3)–(G.8) and (G.12) are unaffected by adding \bullet_{1_1} 's to inner corners. (G.9)–(G.11) and (G.13) hold vacuously. \square

The **slide** of T at $\{x_i\}$ is

$$(7.1) \quad \text{slide}_{\{x_i\}}(T) := \text{swap}_{\mathcal{G}_{\max}} \circ \text{swap}_{\mathcal{G}_{\max}^-} \circ \dots \circ \text{swap}_{1_1}(T^{(1_1)}),$$

with all $\bullet_{\mathcal{G}_{\max}^+}$'s deleted. If Σ is a formal $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ -linear sum of tableaux we write $V \in \Sigma$ to mean V occurs in Σ with nonzero coefficient. The following proposition, proved in Appendix A, shows (7.1) is well-defined.

Proposition 7.7 (Swaps preserve goodness). *If T is a \mathcal{G} -good tableau, then each $U \in \text{swap}_{\mathcal{G}}(T)$ is \mathcal{G}^+ -good.*

Lemma 7.8 (Swaps preserve content). *If T is a \mathcal{G} -good tableau of content μ , then each $U \in \text{swap}_{\mathcal{G}}(T)$ has content μ .*

Proof. No miniswap eliminates genes in a section. We consider each miniswap that introduces a new gene to a section; this gene must be \mathcal{G} . We show that \mathcal{G} appears elsewhere in T . The first case is H2, which produces a \mathcal{G} in its section, where there was only a (\mathcal{G}) previously. (\mathcal{G}) only appears if some \mathcal{G} is west of it in T . The same analysis applies *verbatim* to H7 and T5. The remaining cases are T1 and T3. By Lemma 6.9(IV, V), the snake on which these miniswaps act has at least two rows. Moreover, there is a \mathcal{G} directly below the $\bullet_{\mathcal{G}}$ under consideration. In particular, \mathcal{G} already appeared in T . \square

Lemma 7.9. *No label is strictly southeast of a $\bullet_{\mathcal{G}_{\max}^+}$ in any $U \in \text{swap}_{\mathcal{G}_{\max}} \circ \text{swap}_{\mathcal{G}_{\max}^-} \circ \dots \circ \text{swap}_{1_1}(T^{(1_1)})$. In particular, all $\bullet_{\mathcal{G}_{\max}^+}$'s are at outer corners of ν/λ .*

Proof. By Proposition 7.7, U is \mathcal{G}_{\max}^+ -good. Let x be a box of U and $\bullet_{\mathcal{G}_{\max}^+} \in x$. There is no $\bullet_{\mathcal{G}_{\max}^+}$ strictly southeast of x by (G.2). By definition, there is no label Q in $T^{(1_1)}$ with $\text{family}(Q) \geq \text{family}(\mathcal{G}_{\max}^+)$. Hence by Lemma 7.8, there are no such labels in U . Therefore, any genetic label ℓ southeast of x is marked. Clearly, we may assume ℓ is in x 's row or column. If ℓ is in x 's column, we contradict (G.11). If ℓ is in x 's row, we contradict Lemma 5.5. \square

Clearly,


Lemma 7.10. *If T is a good tableau with no genetic label southeast of a \bullet , then deleting all \bullet 's gives a bundled tableau.* \square

Corollary 7.11. *Given $\rho \in \lambda^+$ and a tableau $T \in B_{\rho, \mu}^{\nu}$, any tableau $U \in \text{slide}_{\rho/\lambda}(T)$ is in either $B_{\lambda, \mu}^{\nu}$ or $B_{\lambda, \mu}^{\delta}$ for some $\delta \in \nu^-$.*


Proof. By Lemma 5.3, T is a good tableau. By Lemma 7.6, adding \bullet_{1_1} to each box of ρ/λ gives a good tableau $T^{(1_1)}$. By Proposition 7.7, each swap gives a formal sum of good tableaux. By Lemma 7.9, after all swaps, $\bullet_{\mathcal{G}_{\max}^+}$'s are at outer corners with no labels strictly southeast. By Lemma 7.10, deleting these $\bullet_{\mathcal{G}_{\max}^+}$'s gives a bundled tableau (namely U). U has shape ν/λ or δ/λ for $\delta \in \nu^-$, since there is at most one $\bullet_{\mathcal{G}_{\max}^+}$ deleted in any row or column by (G.2). Content preservation is Lemma 7.8. \square

7.3. Examples. We give a number of examples of computing $\text{slide}_{\{x_i\}}(T)$. It is convenient to encode the computations in a diagram. Each non-terminal tableau has its snakes differentiated by color. The notation above each arrow indicates the types of the snakes from southwest to northeast, for example H5.3/ \emptyset/\emptyset /T2 means the head of the snake is H5.3, the body is empty and the tail is T2. The notation below arrows indicates the product of the coefficients coming from each miniswap (we will assume for this purpose that the lower left corner of T coincides with the lower left corner of $k \times (n - k)$). Each $U \in \text{slide}_{\{x_i\}}(T)$ is a terminal tableau of the diagram. Moreover, $[U]\text{slide}_{\{x_i\}}(T)$ is the sum of the products of the coefficients over all directed paths from T to U .


Example 7.12.




Example 7.13.



Example 7.14.




Example 7.15.




□

Example 7.16.




□

Example 7.17.



□

Example 7.18.




□

8. LADDERS

Let U be a \mathcal{G}^+ -good tableau. Consider the boxes of U containing $\bullet_{\mathcal{G}^+}$ or unmarked \mathcal{G} . This set decomposes into maximal edge-connected components, which we call **ladders**.


Example 8.1.



This 2_2 -good tableau has three ladders; we have given each ladder a separate color. (All virtual labels are depicted.)

□

Lemma 8.2. *A row r of a ladder L is one of the following (edge labels other than \mathcal{G} and virtual labels are not shown):*



Proof. By (G.2), at most one $\bullet_{\mathcal{G}^+}$ occurs in each row. By Lemma 5.10, at most one \mathcal{G} appears in each row. Thus r has at most two boxes. If it has one box, r is clearly L1, L2 or L3. If r has two boxes, then it has one box label \mathcal{G} and one box label $\bullet_{\mathcal{G}^+}$. Since the \mathcal{G} is not marked, it is West of the $\bullet_{\mathcal{G}^+}$. By (G.4) and (G.7), no edge label \mathcal{G} is possible in this two-box scenario. Thus L4 is the only two box possibility. □

Lemma 8.3. *A ladder L is a short ribbon where each column with 2 boxes is $\begin{matrix} \boxed{\mathcal{G}} \\ \boxed{\bullet} \end{matrix}$.*

Proof. In each column, there is at most one $\bullet_{\mathcal{G}^+}$ by (G.2) and at most one \mathcal{G} by (G.4). If the column consists of $\bullet_{\mathcal{G}^+}$ and \mathcal{G} , then the \mathcal{G} is North of the $\bullet_{\mathcal{G}^+}$, since otherwise the \mathcal{G} is marked. Therefore the columns are as described.

If L has a 2×2 subsquare the North box of each column must contain \mathcal{G} , violating (G.3). Each row has at most two boxes by Lemma 8.2. That L is a skew shape is now immediate from the descriptions of L 's rows and columns. □

Lemma 8.4 (Relative positioning of ladders). *Suppose U is \mathcal{G}^+ -good, and that L, M are distinct ladders of U . Then, up to relabeling of the ladders, L is entirely SouthWest of M (that is, if b, b' are boxes of L, M respectively, then b is SouthWest of b').*

Proof. Suppose not. There are three cases to consider:

Case 1: ($b \in L$ is NorthWest of $b' \in M$): By definition, b and b' contain either $\bullet_{\mathcal{G}^+}$ or \mathcal{G} . By (G.2) and Lemmas 5.10, 5.11 and 5.12, we see that no combination of these choices is possible.

Case 2: (b is North and in the same column as b'): If $\bullet_{\mathcal{G}^+} \in b$ and $\bullet_{\mathcal{G}^+} \in b'$, we violate (G.2). If $\bullet_{\mathcal{G}^+} \in b$ and $\mathcal{G} \in b'$, then the latter would be marked. Hence $\mathcal{G} \in b$. Since $\mathcal{G} \in b'$ or $\bullet_{\mathcal{G}^+} \in b'$, we have by (G.4) and (G.9) that $b^\downarrow = b'$ and so b, b' are in the same ladder, contradicting $L \neq M$.

Case 3: (b is West and in the same row as b'): By (G.2), at least one of b, b' contains \mathcal{G} . By Lemma 5.10, at least one of b, b' contains $\bullet_{\mathcal{G}^+}$. If $\mathcal{G} \in b$ and $\bullet_{\mathcal{G}^+} \in b'$, then by (G.3) and (G.9), $b' = b^\rightarrow$, contradicting $L \neq M$. If $\bullet_{\mathcal{G}^+} \in b$ and $\mathcal{G} \in b'$, then the latter is marked. □

9. REVERSE GENOMIC JEU DE TAQUIN

Let r be a ladder row in a \mathcal{G}^+ -good tableau U and let x be the westmost box in r . We define the **reverse miniswap** operation revmswap on r . The cases below are labeled in accordance with the classification of Lemma 8.2. Below, each \bullet on the left of the “ \mapsto ” is a $\bullet_{\mathcal{G}^+}$, while on the right it is a $\bullet_{\mathcal{G}}$.

(Case L1):

(Subcase L1.1: $\mathcal{G} \in x^\uparrow$):

$$r = \boxed{\bullet} \mapsto \text{revmswap}(r) = \boxed{\mathcal{G}}$$

(Subcase L1.2: $\mathcal{G} \notin x^\uparrow$):

$$r = \boxed{\bullet} \mapsto \text{revmswap}(r) = \boxed{\bullet}$$

(Case L2):

(Subcase L2.1: $\bullet_{\mathcal{G}^+} \in x^\downarrow$ or $\mathcal{G}^! \in x^\downarrow$):

$$r = \boxed{\mathcal{G}} \mapsto \text{revmswap}(r) = \boxed{\bullet}$$

(Subcase L2.2: $\bullet_{\mathcal{G}^+} \notin x^\downarrow$, $\mathcal{G}^! \notin x^\downarrow$, $\mathcal{G}^! \notin x$, x contains the westmost \mathcal{G}):

$$r = \boxed{\mathcal{G}} \mapsto \text{revmswap}(r) = \boxed{\mathcal{G}} + \boxed{\bullet_{\mathcal{G}}}$$

(Subcase L2.3: $\bullet_{\mathcal{G}^+} \notin x^\downarrow$, $\mathcal{G}^! \notin x^\downarrow$, $\mathcal{G}^! \notin x$, x does *not* contain the westmost \mathcal{G}):

$$r = \boxed{\mathcal{G}} \mapsto \text{revmswap}(r) = \boxed{\mathcal{G}} + \boxed{\bullet_{\mathcal{G}}}$$

(Case L3):

$$r = \boxed{\begin{smallmatrix} \mathcal{G} \\ \bullet \end{smallmatrix}} \mapsto \text{revmswap}(r) = \boxed{\begin{smallmatrix} \bullet \\ \mathcal{G} \end{smallmatrix}}$$

(Case L4):

(Subcase L4.1: $\mathcal{G}^+ \in x^\rightarrow$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, and either $\bullet_{\mathcal{G}^+} \in x^\downarrow$ or $\mathcal{G}^! \in x^\downarrow$):

$$r = \boxed{\begin{smallmatrix} \mathcal{G} & \bullet \\ \mathcal{G}^+ & \end{smallmatrix}} \mapsto \text{revmswap}(r) = \boxed{\begin{smallmatrix} \bullet & \mathcal{G}^+ \\ & \end{smallmatrix}}$$

(Subcase L4.2: $\mathcal{G}^+ \in x^\rightarrow$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, $\bullet_{\mathcal{G}^+} \notin x^\downarrow$, $\mathcal{G}^! \notin x^\downarrow$ and x contains the westmost \mathcal{G}):

$$r = \boxed{\begin{smallmatrix} \mathcal{G} & \bullet \\ \mathcal{G}^+ & \end{smallmatrix}} \mapsto \text{revmswap}(r) = \boxed{\begin{smallmatrix} \bullet & \mathcal{G}^+ \\ \mathcal{G} & \end{smallmatrix}}$$

(Subcase L4.3: $\mathcal{G}^+ \in x^\rightarrow$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, $\bullet_{\mathcal{G}^+} \notin x^\downarrow$, $\mathcal{G}^! \notin x^\downarrow$ and x does *not* contain the westmost \mathcal{G}):

$$r = \boxed{\begin{smallmatrix} \mathcal{G} & \bullet \\ \mathcal{G}^+ & \end{smallmatrix}} \mapsto \text{revmswap}(r) = \boxed{\begin{smallmatrix} \bullet & \mathcal{G}^+ \\ \mathcal{G} & \end{smallmatrix}}$$

(Subcase L4.4: there is *no* $\mathcal{G}^+ \in x^\rightarrow$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, and x contains the westmost \mathcal{G}): Let A be the labels in \bar{x} , $Z = \{\mathcal{E} \in A : N_{\mathcal{G}} = N_{\mathcal{E}}\}$, $Z^\# = Z \cup \{\mathcal{G}\}$, $\mathcal{F} = \min Z^\#$, $A'' = Z^\# \setminus \{\mathcal{F}\}$, and $A' = A \setminus Z$.

$$r = \boxed{\begin{smallmatrix} A \\ \mathcal{G} \end{smallmatrix}} \boxed{\bullet} \mapsto \text{revmswap}(r) = \boxed{\begin{smallmatrix} A' \\ \bullet \end{smallmatrix}} \boxed{\begin{smallmatrix} \mathcal{F} \\ A'' \end{smallmatrix}}$$

(Subcase L4.5: there is no $\mathcal{G}^+ \in \underline{x}$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$, and x does not contain the westmost \mathcal{G}): Let $A, Z, Z^\sharp, \mathcal{F}$ and A' be as in L4.4; also let $A''' = Z \setminus \{\mathcal{F}\}$.

$$r = \begin{array}{|c|c|} \hline A \\ \hline \mathcal{G} & \bullet \\ \hline \end{array} \mapsto \text{revmswap}(r) = \begin{array}{|c|c|} \hline A' \\ \hline \bullet & \mathcal{F} \\ \hline \end{array}_{A''', \mathcal{G}}$$

Lemma 9.1. *Every ladder row falls into exactly one of the above cases.*

Proof. This is tautological, given Lemma 8.2. □

Lemma 9.2. *No revmswap affects an edge that is shared by two rows of the same ladder L .*

Proof. No revmswap affects the upper (virtual) edge labels of the right box of a ladder row. Hence it suffices to analyze those cases that affect the lower (virtual) edge labels of the left box of a ladder row. These are L2.2, L2.3, L3, L4.2 and L4.3. In each case there can be no ladder row of L below, by Lemma 8.3. Hence that edge is not shared. □

Thus it makes sense to define $\text{revswap}_{\mathcal{G}^+}$ on a ladder L , by applying revmswap to each row of L simultaneously (where the conditions on each revmswap refer to the original ladder L).

Lemma 9.3. *If L_1, L_2 are distinct ladders in a \mathcal{G}^+ -good tableau U , then applying $\text{revswap}_{\mathcal{G}^+}$ to L_1 commutes with applying $\text{revswap}_{\mathcal{G}^+}$ to L_2 .*

Proof. This follows, since by definition L_1 and L_2 do not share any edges. □

Lemma 9.3 permits us to define the **reverse swap** $\text{revswap}_{\mathcal{G}^+}$ on a \mathcal{G}^+ -good tableau by applying $\text{revswap}_{\mathcal{G}^+}$ to all ladders (in arbitrary order). We extend this to a $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ -linear operator.

Lemma 9.4 (Reverse swaps preserve content). *If U is \mathcal{G}^+ -good and of content μ , then each $T \in \text{revswap}_{\mathcal{G}^+}(U)$ has content μ .*

Proof. Let \mathcal{H} be a gene in U . We must show $\mathcal{H} \in T$. Let ℓ be the westmost instance of \mathcal{H} in U . If ℓ is not part of a ladder, \mathcal{H} appears in the same location in T and we are done. Thus suppose ℓ is in a ladder row r . Consider the reverse miniswap applied to r . If it is anything but L2.1 or L4.1, then there is an \mathcal{H} in that row of T . If it is L2.1 or L4.1, let x be the box containing ℓ . By definition, U has $\bullet_{\mathcal{H}^+} \in x^\downarrow$ or $\mathcal{H}^! \in x^\downarrow$. In the former case, the miniswap applied at x^\downarrow is L1.1, so \mathcal{H} appears in x^\downarrow in T . In the latter case, x^\downarrow is not in a ladder, so \mathcal{H} appears in x^\downarrow in T .

Conversely suppose \mathcal{H} is not a gene in U . We must show it does not appear in T . If it appeared in T , it must be created by some miniswap. Clearly no miniswap but L1.1 could possibly introduce a new gene. But if we apply L1.1 at some box x of U , introducing $\mathcal{H} \in x$ in T , then U has $\mathcal{H} \in x^\uparrow$ by definition, so \mathcal{H} was indeed a gene of U . □

We prove the following proposition in Appendix B.

Proposition 9.5 (Reverse swaps preserve goodness). *If U is \mathcal{G}^+ -good, each $T \in \text{revswap}_{\mathcal{G}^+}(U)$ is \mathcal{G} -good.*

Lemma 9.6. *Let T be a \mathcal{G} -good tableau and $U \in \text{swap}_{\mathcal{G}}(T)$.*

(I) *If $\text{label}_U(x) = \mathcal{G}$, then $\text{label}_T(x) \in \{\bullet_{\mathcal{G}}, \mathcal{G}\}$.*

- (II) If $\text{label}_U(x) = \bullet_{\mathcal{G}^+}$, then $\text{label}_T(x) \in \{\bullet_{\mathcal{G}}, \mathcal{G}, \mathcal{F}^!, \mathcal{G}^+\}$.
- (III) If $\text{label}_U(x) = \mathcal{G}^!$, then $\text{label}_T(x) = \mathcal{G}$.

Proof. By inspection of the miniswaps. □

Lemma 9.7. *Let U be a \mathcal{G}^+ -good tableau and $T \in \text{revswap}_{\mathcal{G}^+}(U)$.*

- (I) If $\text{label}_U(x) = \mathcal{G}^!$, then $\text{label}_T(x) = \mathcal{G}$.
- (II) If $\text{label}_U(x) = \mathcal{G}$, then $\text{label}_T(x) \in \{\mathcal{G}, \bullet_{\mathcal{G}}\}$.
- (III) If $\text{label}_U(x) = \bullet_{\mathcal{G}^+}$, then $\text{label}_T(x) \in \{\bullet_{\mathcal{G}}, \mathcal{G}, \mathcal{G}^+, \mathcal{F}^!\}$. If moreover $\text{label}_T(x) = \mathcal{G}^+$, then $\text{label}_T(x^{\leftarrow}) = \bullet_{\mathcal{G}}$, while if moreover $\text{label}_T(x) = \mathcal{F}^!$, then $N_{\mathcal{F}} = N_{\mathcal{G}}$, $\text{label}_T(x^{\leftarrow}) = \bullet_{\mathcal{G}}$ and either $\mathcal{G} \in \underline{x}$ or $(\mathcal{G}) \in \underline{x}$.

Proof. By inspection of the reverse miniswaps. □

For a good tableau T of shape ν/λ , define a T -**patch** of ν/λ as one of the following:

- (Pat.1) A row of a snake of T (including both upper and lower edges of the row).
- (Pat.2) A box not in a snake (the box excludes the edges).
- (Pat.3) A horizontal edge not bounding a box of a snake.

Clearly, the set $\{P\}$ of T -patches covers ν/λ . Given a tableau W of shape ν/λ , let $W|_P$ be the tableau obtained by restricting W to P .

Proposition 9.8. *Let T, U be good. Then $U \in \text{swap}_{\mathcal{G}}(T)$ if and only if $T \in \text{revswap}_{\mathcal{G}^+}(U)$.*

Proof. (\Rightarrow) Suppose $U \in \text{swap}_{\mathcal{G}}(T)$. We show $T \in \text{revswap}_{\mathcal{G}^+}(U)$.

Claim 9.9. *Every ladder row r of U is contained in a distinct T -patch.*

Proof. Distinctness is clear. We now argue containment. If r has one box, containment is trivial. Otherwise, r has two boxes, and we are in case L4 of the ladder row classification of Lemma 8.2. So, in U , each box of r contains $\bullet_{\mathcal{G}^+}$ or \mathcal{G} . One considers all possibilities, under Lemma 9.6, for the entries in T of the boxes of r . Since T is good, these boxes of T either form a row of a snake section or are $\boxed{\mathcal{G}} \boxed{\mathcal{G}^+}$. We are done by (Pat.1) in the former case. The latter case cannot occur, since by inspection of the miniswaps, this cannot swap to L4. □

By the definitions, notice that $\text{revswap}_{\mathcal{G}^+}(U) \neq 0$. Moreover:

Claim 9.10. *For each T -patch P , there exists $W \in \text{revswap}_{\mathcal{G}^+}(U)$ such that $W|_P = T|_P$ (ignoring virtual labels).*

Proof. If P is type (Pat.2), then by definition $T|_P = U|_P$, since P is not part of a snake. In particular $U|_P$ does not contain \mathcal{G} or $\bullet_{\mathcal{G}^+}$. So $U|_P$ is not part of a ladder of U . Hence for any $W \in \text{revswap}_{\mathcal{G}^+}(U)$, $W|_P = U|_P = T|_P$ as desired.

If P is type (Pat.3), then $T|_P = U|_P$, since P is not part of a snake. Moreover, by definition, no box y bounded by the edge P is part of a snake in T . Therefore, $\bullet_{\mathcal{G}}, \mathcal{G} \notin y$ in T . Hence $\bullet_{\mathcal{G}^+}, \mathcal{G} \notin y$ in U . So P does not bound a box of a ladder of U . Thus for any $W \in \text{revswap}_{\mathcal{G}^+}(U)$, $W|_P = U|_P = T|_P$.

Finally if P is type (Pat.1), by inspection of the miniswaps, combined with Claim 9.9, $U|_P$ contains at most one ladder row r , and possibly a non-ladder box y . Since $\text{revswap}_{\mathcal{G}^+}$

does not affect y , it suffices to indicate the reverse miniswap on r to give our desired $W|_P = T|_P$. We refer to the list of outputs described in Section 7.

H1: Use L2.2 or L3 respectively on the two mswap outputs.

H2: Use L1.2 or L2.3 respectively on the two mswap outputs.

H3: Use L1.2: By T 's (G.2) and (G.9) and Lemma 9.6(I) applied to T , we have $\mathcal{G} \notin x^\uparrow$ in U .

H4: This case does not arise, since here U does not exist.

H5.1: Use L1.2.

H5.2: For the first output, use L1.2. For the second output, use L4.4 or L4.5. We must show in the latter cases that $Z = \emptyset$. Otherwise if $\mathcal{E} \in Z$, then $\mathcal{E} \in \bar{x}$ in T . Since $N_{\mathcal{E}} = N_{\mathcal{G}}$ in both T and U , this contradicts Lemma 5.6 for T .

H5.3: Use L4.4 or L4.5. The argument that these apply is the same as for H5.2.

H6: Use L2.2 for the first output and L4.2 for the second. By Lemma 9.6(II) and T 's (G.2) and (G.4), $\bullet_{\mathcal{G}^+} \notin x^\downarrow$; by Lemma 9.6(I) and T 's (G.2) or (G.4), $\mathcal{G}^\downarrow \notin x^\downarrow$; that the $\mathcal{G} \in x$ is westmost follows from T 's (G.7) and Claim A.3 applied to T .

H7: Use L1.2 for the first output: By Lemma 9.6(I) and T 's (G.2) and (G.9), U has $\mathcal{G} \notin x^\uparrow$. Use L2.3 for the second output and L4.3 for the third: By Lemma 9.6(II) and T 's (G.2) or (G.4), $\bullet_{\mathcal{G}^+} \notin x^\downarrow$; by Lemma 9.6(I) and T 's (G.2) or (G.4), $\mathcal{G}^\downarrow \notin x^\downarrow$; that the $\mathcal{G} \in x$ is not westmost follows from T 's $\mathcal{G} \in \underline{x}$.

H8: Use L1.2: By T 's (G.2) and (G.9) and Lemma 9.6(I), U has $\mathcal{G} \notin x^\uparrow$.

H9: Here r does not exist.

B1: Use L2.2 or L2.3: By Lemma 9.6(II) and T 's (G.2) or (G.4), $\bullet_{\mathcal{G}^+} \notin x^\downarrow$; by Lemma 9.6(III) and T 's (G.4), $\mathcal{G}^\downarrow \notin x^\downarrow$.

B2: Use L4.4 or L4.5; applicability is as for H5.2.

B3: If we are not in the bottom row, we may use L4.4 or L4.5 as for B2. Otherwise, use L1.1.

T1: Use L2.1: By Lemma 6.9(IV, VII), T has $\mathcal{G} \in x^\downarrow$, so the hypothesis holds by inspection of the miniswaps.

T2: Use L4.4 or L4.5; applicability is as for H5.2.

T3: Use L2.1 or L4.1; applicability is as for T1.

T4.1: Use L1.2: By T 's (G.2) and (G.12) and Lemma 9.6(I), U has $\mathcal{G} \notin x^\uparrow$ and $\mathcal{G} \notin \bar{x}$.

T4.2: Use L1.2 on the first output; applicability is as for T4.1. Use L4.4 on the second output; applicability is as for H5.2.

T4.3: Use L4.4; applicability is as for H5.2.

T5: Use L4.5; applicability is as for H5.2.

T6: This case does not arise, since here U does not exist. □

By definition, $\text{revswap}_{\mathcal{G}^+}(U)$ is obtained by acting on ladder rows of U independently. By Claim 9.10, it follows that $\text{revswap}_{\mathcal{G}^+}(U)$ is also obtained by acting on the T -patches of U independently. Thus (\Leftarrow) holds by Claim 9.9.

(\Leftarrow) Suppose $T \in \text{revswap}_{\mathcal{G}^+}(U)$. We show U is in $\text{swap}_{\mathcal{G}}(T)$.

Recall $\text{swap}_{\mathcal{G}}(T)$ is a formal sum, given by independently replacing each snake section in each prescribed way. Trivially, by (Pat.1), each snake section is a union of T -patches. Moreover, if a snake section σ consists of more than one T -patch, then σ is a body with at least two rows, and hence either B2 or B3. Therefore $\text{mswap}(\sigma)$ has a unique output in this case. Since $\text{swap}_{\mathcal{G}}$ acts trivially on the T -patches of types (Pat.2) and (Pat.3), by Lemma 7.5, it follows that $\text{swap}_{\mathcal{G}}(T)$ is also given by acting independently on the T -patches of T . It remains to show that locally at P , we may swap $T|_P$ to obtain $U|_P$.

To make these local verifications, we use:

Claim 9.11.

- (I) *Every ladder row of U sits in a distinct T -patch of type (Pat.1).*
- (II) *Every T -patch P of type (Pat.1) not coming from an H9 snake section, contains a ladder row of U .*

Proof. (I): By Lemma 9.7, every ladder row of U is contained in a T -patch of type (Pat.1). Consider a T -patch P of type (Pat.1); P consists of at most two boxes. If P does not consist of two boxes, clearly at most one ladder row of U can be contained in it. If P consists of two boxes, they are joined by a vertical edge. Since distinct ladder rows do not share a vertical edge, it follows that distinct ladder rows of U are contained in distinct T -patches.

(II): By inspection of the reverse miniswaps. □

If P is type (Pat.2) or (Pat.3), then by Claim 9.11, P does not intersect any ladder row of U . Thus $T|_P = U|_P$. By definition, P is not part of any snake in T . Hence for any $V \in \text{swap}_{\mathcal{G}}(T)$, $V|_P = T|_P = U|_P$ as desired.

Finally suppose P is a patch of type (Pat.1). If it comes from an H9 snake section, then $V \in \text{swap}_{\mathcal{G}}(T)$, $V|_P = T|_P = U|_P$. Otherwise, by Claim 9.11, P contains a unique ladder row in U . We consider each ladder row type in turn and indicate the miniswaps on $T|_P$ that give our desired $V|_P = U|_P$. We refer to the list of outputs described at the beginning of Section 9. The following case analysis completes the proof of (\Rightarrow) .

L1.1: Use B3: Since $\text{label}_U(x^\uparrow) = \mathcal{G}$, we apply at x^\uparrow either L2.1, L4.1, L4.4 or L4.5. In each case $\text{label}_T(x^\uparrow) = \bullet_{\mathcal{G}^+}$. Hence x and x^\uparrow are part of the southmost two rows of a snake of T . We claim x^\leftarrow is not part of this snake. Note that by assumption x^\leftarrow is not part of any ladder of U . Thus $\text{label}_U(x^\leftarrow) = \text{label}_T(x^\leftarrow)$ and $\text{label}_T(x^\leftarrow) \notin \{\bullet_{\mathcal{G}}, \mathcal{G}\}$. If x^\leftarrow is part of x 's snake in T , then $\text{label}_T(x^\leftarrow) = \mathcal{F}^\dagger \prec \mathcal{G}$ and southeast of some $\bullet_{\mathcal{G}}$. Hence in U , x^\leftarrow is southeast of some $\bullet_{\mathcal{G}^+}$; this contradicts U 's (G.2) in view of U 's $\bullet_{\mathcal{G}} \in x$. Thus x is the unique box of the southmost row of its snake and by Definition-Lemma 6.8, it is the southmost row of a B3 snake section.

L1.2: Use H2, H3, H7, H8, T4.1 or T4.2: Since $\text{label}_T(x^\downarrow) = \mathcal{G}$, $\text{label}_U(x^\downarrow) \in \{\bullet_{\mathcal{G}^+}, \mathcal{G}\}$. Hence by Lemma 8.3, x^\downarrow is not in x 's snake in T . Since $\text{label}_T(x) = \bullet_{\mathcal{G}}$, x^\uparrow is not in x 's snake in T . Hence x is in a one-row snake. Since L1.2 applies, $\text{label}_U(x^\rightarrow) \neq \mathcal{G}$, so $\text{label}_T(x^\rightarrow) \neq \mathcal{G}$. Thus x 's snake in T is type (ii), (iv) or (vi) in Definition-Lemma 6.8(III). Type (ii) uses H2 or H3; type (iv) uses H7 or H8; type (vi) uses T4.1 or T4.2.

L2.1: Use T1 or T3: By assumption, $\text{label}_U(x^\downarrow) \in \{\bullet_{\mathcal{G}^+}, \mathcal{G}^\dagger\}$. Hence by inspection of the reverse miniswaps, $\text{label}_T(x^\downarrow) = \mathcal{G}$. Since $\text{label}_T(x) = \bullet_{\mathcal{G}}$, x^\uparrow is not in x 's snake. Hence by Definition-Lemma 6.8(I,II), x is in its snake's tail. By T 's (G.3), $\text{label}_T(x^\rightarrow) \succ \mathcal{G}$, so $\text{label}_U(x^\rightarrow) \neq \mathcal{F}^\dagger$. Thus either T1 or T3 applies.

L2.2: Use B1 for the first output. By assumption and U 's (G.9), U has no $\bullet_{\mathcal{G}^+}$ adjacent to x . Moreover by U 's (G.4), no box adjacent to x is in any ladder. Hence T has no $\bullet_{\mathcal{G}}$ adjacent to x . If $\mathcal{F}^! \in x^{\leftarrow}$ in T , then (possibly marked) $\mathcal{F} \in x^{\leftarrow}$ in U . If $\text{label}_U(x^{\leftarrow}) = \mathcal{F}^!$, then we contradict unmarked $\mathcal{G} \in x$ in U . If $\text{label}_U(x^{\leftarrow})$ is unmarked, then U has no $\bullet_{\mathcal{G}^+}$ northwest of x^{\leftarrow} . By U 's (G.3) and (G.4), U has no \mathcal{G} northwest of x^{\leftarrow} . But since $\mathcal{F}^! \in x^{\leftarrow}$ in T , T has a $\bullet_{\mathcal{G}^+}$ northwest of x^{\leftarrow} . Hence by Lemma A.3, U has a $\bullet_{\mathcal{G}^+}$ or \mathcal{G} northwest of x^{\leftarrow} , a contradiction.

Use H1 or H6 for the second output. Since x^{\rightarrow} is not in any ladder of U , $\text{label}_U(x^{\rightarrow}) = \text{label}_T(x^{\rightarrow})$. Moreover by U 's (G.3), $\text{label}_U(x^{\rightarrow}) \succ \mathcal{G}$, so $\text{label}_T(x^{\rightarrow}) \succ \mathcal{G}$. If $\text{label}_T(x^{\rightarrow}) = \mathcal{G}^+$, H6 applies. Otherwise, H1 applies.

L2.3: Use B1 for the first output; applicability is as for the first output of L2.2. Use H2 or H7 for the second output. Since x^{\rightarrow} is not in any ladder of U , $\text{label}_U(x^{\rightarrow}) = \text{label}_T(x^{\rightarrow})$. Moreover by U 's (G.3), $\text{label}_U(x^{\rightarrow}) \succ \mathcal{G}$, so $\text{label}_T(x^{\rightarrow}) \succ \mathcal{G}$. If $\text{label}_T(x^{\rightarrow}) = \mathcal{G}^+$, H7 applies. Otherwise, H2 applies.

L3: Use H1. By U 's (G.12), $\text{label}_U(x^{\rightarrow}) \notin \{\mathcal{G}, \mathcal{G}^+\}$. Moreover by U 's (G.13) and (G.12), $\text{label}_U(x^{\rightarrow})$ is not marked, so $\text{label}_U(x^{\rightarrow}) \succeq \mathcal{G}^+$. Thus $\text{label}_U(x^{\rightarrow}) \succ \mathcal{G}^+$. Since x^{\rightarrow} is not in any ladder of U , $\text{label}_T(x^{\rightarrow}) \succ \mathcal{G}^+$.

L4.1: Use T3. By inspection of the reverse miniswaps, T has $\mathcal{G} \in x^{\downarrow}$. Hence x 's snake in T has at least two rows. Hence x is part of its snake's tail.

L4.2: Use H6.

L4.3: Use H7.

L4.4: If $Z \neq \emptyset$, use T4.2 or T4.3. Otherwise use H5.3, B2, B3 or T2. If $Z \neq \emptyset$, some T4 applies. If it is T4.1, T has $\mathcal{H} \in x^{\rightarrow}$ with $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$. Hence U also has $\mathcal{H} \in x^{\rightarrow}$, contradicting Lemma 5.6 for U . If $Z = \emptyset$, there is nothing to check.

L4.5: If $Z \neq \emptyset$, use T5. Otherwise use H5.3, B2, B3 or T2. \square

The following proposition characterizes good tableaux in terms of forward swapping.

Proposition 9.12. *A tableau U is \mathcal{G} -good if and only if $U \in \text{swap}_{\mathcal{G}^-} \circ \cdots \circ \text{swap}_{1_2} \circ \text{swap}_{1_1}(T^{(1_1)})$ for some bundled tableau T and choice of inner corners of T to initially place \bullet_{1_1} 's in.*

Proof. (\Rightarrow) Given a \mathcal{G} -good tableau U , let $T^{(1_1)}$ be any tableau appearing in $\text{revswap}_{1_2} \circ \cdots \circ \text{revswap}_{\mathcal{G}^-} \circ \text{revswap}_{\mathcal{G}}(U)$. By Proposition 9.5, $T^{(1_1)}$ is a 1_1 -good tableau. By $T^{(1_1)}$'s (G.2) and (G.9), the \bullet_{1_1} 's of $T^{(1_1)}$ are at inner corners and there is no genetic label northwest of a \bullet_{1_1} . Let T be obtained by removing the \bullet_{1_1} 's of $T^{(1_1)}$. Then it is clear T is a bundled tableau. Now $U \in \text{swap}_{\mathcal{G}^-} \circ \cdots \circ \text{swap}_{1_2} \circ \text{swap}_{1_1}(T^{(1_1)})$ holds by Propositions 7.7 and 9.8.

(\Leftarrow) Immediate from Lemma 7.6 and Proposition 7.7. \square

10. THE REVERSAL TREE

10.1. **Walkways.** An i -walkway W in an $(i+1)_1$ -good tableau T is an edge-connected component of the collection of boxes x in T such that:


(W.1) $\bullet_{(i+1)_1} \in x$; or

(W.2) $i_k \in x$ and x is not southeast of a $\bullet_{(i+1)_1}$ (equivalently, $i_k \in x$ is not marked).

Lemma 10.1 (Structure of an i -walkway). *Let W be an i -walkway.*

- (I) Each column c of W has at most two boxes; if c has two boxes, the southern box contains $\bullet_{(i+1)_1}$.
- (II) W has no 2×2 subsquare.
- (III) W is an edge-connected skew shape.
- (IV) The $\bullet_{(i+1)_1}$'s are at outer corners of W .
- (V) The box and upper edge labels of family i form a \prec -interval in the set of genes.

Therefore, each i -walkway looks like:



where each \star is a genetic label and the blank box contains either $\bullet_{(i+1)_1}$ or a genetic label.

Proof of Lemma 10.1: (I): By (G.2), at most one box of c comes from (W.1). By (G.4), at most one box of c comes from (W.2). Thus the first assertion of (I) holds. The second assertion holds by (W.2).

(II): Suppose W contains a 2×2 subsquare. Then the two southern boxes of the subsquare contain $\bullet_{(i+1)_1}$'s by (I), contradicting (G.2).

(III): W is edge-connected by definition. In view of (II), it remains to show there are no two boxes y, z of W with y NorthWest of z . Suppose otherwise. By (G.2), at least one of y, z contains a genetic label. If $\bullet_{(i+1)_1} \in y$ and $i_k \in z$, we violate (W.2). If $\bullet_{(i+1)_1} \in z$ and $i_k \in y$, consider the box b in y 's column and z 's row. By (G.2), b contains a genetic label. By (G.4), $\text{label}(b) > i_k$. Since $\bullet_{(i+1)_1} \in z$, this contradicts (G.9). Finally, if $i_k \in y$ and $i_h \in z$, then we contradict (G.12).

(IV): Immediate from (W.2) and (G.2).

(V): By the edge-connectedness of W we know that W occupies consecutive columns. Thus we are done by (G.4)–(G.6). \square

10.2. Walkway reversal. Let $U \in B_{\lambda, \mu}^\alpha$ for some $\alpha \in \{\nu\} \cup \nu^-$. Obtain $U^{(0)}$ from U by placing $\bullet_{(\ell(\mu)+1)_1}$ in each box of ν/α . The root of the **reversal tree** \mathfrak{T}_U is $U^{(0)}$. The children $\{U^{(1)}\}$ of $U^{(0)}$ are the tableaux in the formal sum $\text{revswap}_{\ell(\mu)_1^+} \circ \cdots \circ \text{revswap}_{(\ell(\mu)+1)_1}(U^{(0)})$. By Proposition 9.5, each $U^{(1)}$ is $\ell(\mu)_1$ -good. We define the children $\{U^{(2)}\}$ of a $U^{(1)}$ by reverse swapping successively through labels of family $\ell(\mu) - 1$, etc. Similarly, all tableaux thus obtained are also good. (A tableau may have a copy of itself as a child; this occurs only if $U^{(0)}$ has no $\bullet_{(\ell(\mu)+1)_1}$'s.) After $\ell(\mu) - i$ steps, a descendant $U' = U^{(\ell(\mu)-i)}$ is an $(i+1)_1$ -good tableau.

Lemma 10.2. *Let U' be an $(i+1)_1$ -good tableau. If ℓ is a box or edge label that is not in an i -walkway, then ℓ appears in the same location in every $T \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$.*

Proof. The case analysis is as follows:

Case 1: ($\ell \in x$ is a box label in U'):

Subcase 1.1: ($\text{family}(\ell) \neq i$): During the reversal process $\text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}$, the label ℓ is never part of any ladder consisting of \mathcal{H} and $\bullet_{\mathcal{H}^+}$ where $\mathcal{H} \in \{i_1, \dots, i_{\mu_i}\}$. Thus $\text{revswap}_{\mathcal{H}^+}$ does not move ℓ .

Subcase 1.2: ($\text{family}(\ell) = i$): Since x is not part of an i -walkway, by (W.2) it is southeast of a $\bullet_{(i+1)_1}$ in U' . By inspection of the reverse miniswaps, this remains true for each tableau

V appearing in the reversal process $\text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}$. The box x is never part of a ladder during this process, for when we apply $\text{revswap}_{\mathcal{H}^+}$, where \mathcal{H} is ℓ 's gene, $\bullet_{\mathcal{H}^+}$ is northwest of x and so $\ell^! \in x$. The case then follows.

Case 2: (ℓ is an edge label in U'): Let x and x^\downarrow be the boxes adjacent to the edge.

Subcase 2.1: (x and x^\downarrow do not contain a label of family i in U'): As above, x and x^\downarrow are not part of a ladder consisting of \mathcal{H} and $\bullet_{\mathcal{H}^+}$, where $\mathcal{H} \in \{i_1, \dots, i_{\mu_i}\}$. Hence neither is the ℓ in question, and so this ℓ remains fixed throughout the reversal process.

Subcase 2.2: (x or x^\downarrow contains a label \mathcal{H} of family i in U'): By (G.4), at most one of x or x^\downarrow contains such a label. Without loss of generality, suppose it is x (the argument in the other case is the same). Since $\ell \in \underline{x}$ is not part of an i -walkway, neither is x . By the arguments of Subcase 1.2, x is never part of a ladder, since $\mathcal{H}^! \in x$. Thus \underline{x} is unchanged. \square

Consider an i -walkway W of U' . By Lemma 10.1(V), the genes of family i in W form an interval; let it be (w_1, \dots, w_n) in increasing \prec -order.

Lemma 10.3 (Characterization of one-row walkway reversals). *Let W be a 1-row i -walkway in an $(i+1)_1$ -good tableau U' . Let a and z be the westmost and eastmost boxes of W , respectively. Consider the region \mathcal{R} occupied by W .*

- (I) *Suppose U' has $\bullet_{(i+1)_1} \in z$ and no label of family i in \bar{z} . Then there exists a filling R of \mathcal{R} with $\bullet_{i_1} \in a$ and $w_1 \notin \underline{a}$ such that for any $V \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$, $V|_{\mathcal{R}} = R$.*
- (II) *Suppose U' has $\bullet_{(i+1)_1} \in z$ and a label of family i in \bar{z} . Then there exists a filling R of \mathcal{R} with $\bullet_{i_1} \in a$ and either $w_1 \in \underline{a}$ or $(w_1) \in \underline{a}$ such that for any $V \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$, $V|_{\mathcal{R}} = R$.*
- (III) *Suppose U' has a label of family i in z . Then there exist two fillings R, R' of \mathcal{R} such that*
 - (i) R has $w_1 \in a$;
 - (ii) R' has $\bullet_{i_1} \in a$ and either $w_1 \in \underline{a}$ or $(w_1) \in \underline{a}$;
 - (iii) R and R' are otherwise identical; and
 - (iv) for any $V \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$, $V|_{\mathcal{R}} \in \{R, R'\}$.

Proof. We argue (I)–(III) separately, by induction on the number of boxes of W . The base cases (where W consists of a single box $a = z$) are clear by Lemma 8.2 and inspection of the reverse miniswaps. Assume W has at least two boxes and let \overline{W} be W with a removed.

(I): By induction, \overline{W} reverses uniquely to some \overline{R} , which has $\bullet_{w_2} \in a^\rightarrow$ and $w_2 \notin \underline{a}^\rightarrow$. (By a technical modification of the hypotheses, we may apply the inductive hypothesis to this partial walkway here and below.) This extends uniquely by L4.4 or L4.5 (followed by some number of applications of L1.2) to an R with the claimed properties.

(II): The unique reversal \overline{R} of \overline{W} has $\bullet_{w_2} \in a^\rightarrow$ and $w_2 \in \underline{a}^\rightarrow$. (By (V.2), $(w_2) \notin \underline{a}^\rightarrow$.) We obtain the desired unique reversal R by applying L4.2 or L4.3 to $\{a, a^\rightarrow\}$ in $\overline{R} \cup \{a\}$.

(III): There are precisely two reversals of \overline{W} : \overline{R} and \overline{R}' . The former reversal has $w_2 \in a^\rightarrow$, while the latter has $\bullet_{w_2} \in a^\rightarrow$ and $w_2 \in \underline{a}^\rightarrow$. (By (V.2), $(w_2) \notin \underline{a}^\rightarrow$.) Applying L4.2 or L4.3 (as appropriate) to $\{a, a^\rightarrow\}$ in $\overline{R}' \cup \{a\}$ returns R' as described. Applying L2.2 or L2.3 (as appropriate) to a in $\overline{R} \cup \{a\}$ returns precisely R and R' . (We apply L4.2 to $\overline{R}' \cup \{a\}$ exactly when we apply L2.2 to $\overline{R} \cup \{a\}$.) \square

Lemma 10.4 (Characterization of multirow walkway reversals). *Let W be an i -walkway with at least two rows in an $(i+1)_1$ -good tableau U' . Let a and z be the westmost and eastmost boxes, respectively, in its southmost row. Thus $\bullet_{(i+1)_1} \in z$. Let \mathcal{R} be the region occupied by W .*

- (I) *Suppose $a = z$. Then there exists a filling R of \mathcal{R} with $w_1 \in a$ such that for any $V \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$, $V|_{\mathcal{R}} = R$.*
- (II) *Suppose $a \neq z$ and $\text{label}_W(z^{\leftarrow}) = \text{label}_W(z^{\uparrow})$. Then there exists a filling R of \mathcal{R} with $\bullet_{i_1} \in a$ and no label of family i on \underline{a} such that for any $V \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$, $V|_{\mathcal{R}} = R$.*
- (III) *Suppose $a \neq z$ and $\text{label}_W(z^{\leftarrow}) \neq \text{label}_W(z^{\uparrow})$. Then there exist two fillings R, R' of \mathcal{R} such that*
 - (i) *R has $w_1 \in a$;*
 - (ii) *R' has $\bullet_{i_1} \in a$ and either $w_1 \in \underline{a}$ or $(w_1) \in \underline{a}$;*
 - (iii) *R and R' are otherwise identical; and*
 - (iv) *for any $V \in \text{revswap}_{i_1^+} \circ \cdots \circ \text{revswap}_{(i+1)_1}(U')$, $V|_{\mathcal{R}} \in \{R, R'\}$.*

Proof. (I): Let \overline{W} be W with the two boxes in the westmost column of W removed. If $\overline{W} = \emptyset$, then $W = \{\bullet_{(i+1)_1} \in z, w_1 \in z^{\uparrow}\}$; here we obtain the desired result by use of L1.1 and L2.1. Hence assume $\overline{W} \neq \emptyset$. Clearly,

$$(10.1) \quad \text{label}_W(z^{\uparrow \rightarrow}) \in \{w_2, \bullet_{(i+1)_1}\}.$$

Depending on whether \overline{W} has multiple rows, by induction or by Lemma 10.3, there are at most two reversals of \overline{W} .

Case 1: (\overline{W} has a unique reversal \overline{R}): By (10.1) and induction/Lemma 10.3, we have two scenarios possible:

Subcase 1.1: (\overline{R} has $\bullet_{w_2} \in z^{\uparrow \rightarrow}$ and no labels of family i appear on $\underline{z^{\uparrow \rightarrow}}$): Here we extend to a unique reversal of W by applying L4.4 or L4.5 at z^{\uparrow} and L1.1 at z . This results in $w_1 \in z = a$.

Subcase 1.2: (\overline{R} has $w_2 \in z^{\uparrow \rightarrow}$): We extend to a unique reversal of W by applying L2.1 at z^{\uparrow} and L1.1 at z . This results in $w_1 \in z = a$, as desired.

Case 2: (\overline{W} has two reversals \overline{R} and \overline{R}'): By (10.1) and induction/Lemma 10.3, \overline{R} and \overline{R}' differ only in $z^{\uparrow \rightarrow}$: \overline{R} has $w_2 \in z^{\uparrow \rightarrow}$ whereas \overline{R}' has $\bullet_{w_2} \in z^{\uparrow \rightarrow}$ and $w_2 \in \underline{z^{\uparrow \rightarrow}}$. By L2.1 and L1.1 in the \overline{R} case and by L4.1 and L1.1 in the \overline{R}' case, both extend to the same reversal R of W ; here R has $w_1 \in z = a$, as claimed.

(II): We have some cases.

Case 1: (The southmost row of W has exactly two boxes $\{a = z^{\leftarrow}, z\}$): Let \overline{W} be W with $\{a, z, z^{\uparrow}\}$ removed. If \overline{W} is empty, the result is clear, so we may assume otherwise. Thus (10.1) still holds. Depending on whether \overline{W} has multiple rows or not, either by induction or by Lemma 10.3, it follows there are at most two reversals of \overline{W} .

Subcase 1.1: (\overline{W} has a unique reversal \overline{R}): By (10.1) and induction/Lemma 10.3, two scenarios are possible:

Subcase 1.1.1: (\overline{R} has $\bullet_{w_2} \in z^{\uparrow \rightarrow}$ and no label of family i on $\underline{z^{\uparrow \rightarrow}}$): We extend to a unique reversal R of W by applying L4.5 at $\{z^{\uparrow}, z^{\uparrow \rightarrow}\}$ and either L4.4 or L4.5 (as required) at $\{a, z\}$; R has $\bullet_{w_1} \in a$ and no label of family i on \underline{a} .

Subcase 1.1.2: ($w_2 \in z^{\uparrow\rightarrow}$): We extend to a unique reversal R of W by applying L2.1 at z^\uparrow and either L4.4 or L4.5 (as required) at z . This again results in $\bullet_{w_1} \in a$ and no label of family i on \underline{a} .

Subcase 1.2: (\overline{W} has two reversals \overline{R} and \overline{R}'): By (10.1) and induction/Lemma 10.3, \overline{R} and \overline{R}' differ only in $z^{\uparrow\rightarrow}$: \overline{R} has a $w_2 \in z^{\uparrow\rightarrow}$ whereas \overline{R}' has a $\bullet_{w_2} \in z^{\uparrow\rightarrow}$ and $w_2 \in \underline{z^{\uparrow\rightarrow}}$. By L2.1 and L4.4 or L4.5 in the \overline{R} case and by L4.1 and L4.4 or L4.5 in the \overline{R}' case, both extend to the same reversal R of W . R has $\bullet_{w_1} \in z = a$.

In each of the Subcases above, we are done after applying a sequence of L1.2's at a .

Case 2: (The southmost row of W contains at least three boxes): Let \overline{W} be W with a removed. By induction, \overline{W} has a unique reversal \overline{R} with \bullet_{w_2} in a^\rightarrow and no label of family i on $\underline{a^\rightarrow}$. Now we uniquely extend \overline{R} to a reversal R of W by applying L4.4 or L4.5 at $\{a, a^\rightarrow\}$; R has $\bullet_{w_1} \in a$ and no label of family i on \underline{a} , and the result follows after applying a sequence of L1.2's at a .

(III): Let \overline{W} be W with the southmost row and z^\uparrow removed. Recall $\text{label}_W(z) = \bullet_{(i+1)_1}$ and suppose W has $w_{q-1} \in z^\leftarrow$ and $w_q \in z^\uparrow$. If \overline{W} is empty, we are done by applying L2.1 at z^\uparrow and L1.1 at z , followed by application of Lemma 10.3(III) to the southmost row. Thus assume \overline{W} is not empty. By induction or Lemma 10.3, there are at most two reversals of \overline{W} :

Case 1: (\overline{W} has a unique reversal \overline{R}): Observe that exactly one of the following two cases holds.

Subcase 1.1: (\overline{R} has $\bullet_{w_{q+1}} \in z^{\uparrow\rightarrow}$ and no label of family i on $\underline{z^{\uparrow\rightarrow}}$): Apply L4.4 at z^\uparrow and L1.1 at z .

Subcase 1.2: (\overline{R} has $w_{q+1} \in z^{\uparrow\rightarrow}$): Apply L2.1 at z^\uparrow and L1.1 at z .

Case 2: (\overline{W} has two reversals \overline{R} and \overline{R}'): By induction/Lemma 10.3, \overline{R} and \overline{R}' differ only in $z^{\uparrow\rightarrow}$: \overline{R} has $w_{q+1} \in z^{\uparrow\rightarrow}$ whereas \overline{R}' has a $\bullet_{w_{q+1}} \in z^{\uparrow\rightarrow}$ and $w_{q+1} \in \underline{z^{\uparrow\rightarrow}}$. Apply L2.1 and L1.1 in the \overline{R} case. Apply L4.1 and L1.1 in the \overline{R}' case.

In each of the cases above, the indicated reverse miniswaps leave us with the southmost row having $w_1 \in a$ and $w_q \in z$. We complete the reversal using Lemma 10.3(III), yielding the desired conclusion. \square

Proposition 10.5. *The children of a node U' in \mathfrak{T}_U are obtained by replacing each walkway W with R or R, R' (as defined in Lemmas 10.3 and 10.4) independently in all possible ways.*

Proof. That nothing changes outside the walkways is Lemma 10.2. Independence follows from walkways being edge-disjoint. \square

Proposition 10.6. *\mathfrak{T}_U is a tree.*

Proof. Let U' and U'' be distinct i_1 -good nodes of \mathfrak{T}_U . By induction and Lemmas 10.3 and 10.4, U' and U'' differ in the placement of a label of family strictly larger than i . This label is unaffected by later reverse swaps, so U' and U'' cannot have the same child. \square

Proposition 10.7 (Characterization of reversal tree leaves).

(I) *Let L be a leaf of \mathfrak{T}_U . Then if we ignore the \bullet_{i_1} 's, either $L = U$ or $L \in \Lambda^+$ and has shape ν/ρ for some $\rho \in \lambda^+$. Moreover, $[U]\text{slide}_{\rho/\lambda}(L) \neq 0$.*

(II) If $M \in \Lambda^+$ has shape ν/ρ and $[U]\text{slide}_{\rho/\lambda}(M) \neq 0$, then M appears as a leaf of \mathfrak{T}_U . \square

Proof. (I): By Proposition 9.5, L is 1_1 -good. By (G.9), there are no labels northwest of a \bullet_{1_1} . By (G.2), \bullet_{1_1} 's appear in distinct rows and columns. This proves the second sentence. The third sentence then follows from Proposition 9.8.

(II): Immediate from Proposition 9.8. \square

11. THE RECURRENCE COEFFICIENTS

Given $U \in B_{\lambda,\mu}^\alpha$, where $\alpha \in \{\nu\} \cup \nu^-$, let $\text{leaf}(\mathfrak{T}_U)$ be the collection of leaves of the tree \mathfrak{T}_U defined in Section 10.

Let W be an i -walkway of shape $\bar{\nu}/\bar{\lambda}$ with $\bullet_{(i+1)_1}$'s in $\bar{\nu}/\bar{\alpha}$. Let S be a reversal of W , as defined by Lemmas 10.3 and 10.4. Let \mathbf{a} be the southwestmost box of W , \mathbf{b} be the northeastmost box of W and \mathbf{z} the eastmost box of W 's southmost row. By Lemma 10.1(V), the labels of family i of S form an interval (w_1, \dots, w_n) with respect to \prec . Let $\bar{\alpha}_*$ denote $\bar{\alpha}$ with its southmost row deleted, and set $\bar{\lambda}_* := \bar{\lambda} \cap \bar{\alpha}_*$. Let $\Delta(S, W) := (\#\bullet_{i_1}$'s in $S) - (\#\bullet_{(i+1)_1}$'s in $W)$. For a tableau T , let \tilde{T} denote T excluding boxes containing w_1 and outer corners containing $\bullet_{w_1^+}$.

Claim 11.1.

(I.i) If S has $w_1 \notin \mathbf{a}^{\rightarrow}$ and w_1 or $(w_1) \in \mathbf{a}$, while W has either at least two rows or $w_n \in \mathbf{b}$, then $[W]\text{slide}_{\bar{\rho}/\bar{\lambda}}(S) = (-1)^{\Delta(S,W)-1} (1 - \text{wt } \bar{\alpha}/(\bar{\alpha}_* \cup \bar{\lambda})) \text{wt } \bar{\alpha}_*/\bar{\lambda}_*$.

(I.ii) If S has $w_1 \notin \mathbf{a}^{\rightarrow}$ and w_1 or $(w_1) \in \mathbf{a}$, while W has exactly one row and $w_n \in \bar{\mathbf{b}}$, then $[W]\text{slide}_{\bar{\rho}/\bar{\lambda}}(S) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda}$.

(II) If S has $\bullet_{i_1} \in \mathbf{a}$, $w_1 \in \mathbf{a}^{\rightarrow}$ and $w_1 \notin \mathbf{a}$, then $[W]\text{slide}_{\bar{\rho}/\bar{\lambda}}(S) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda}$.

(III) If S has $w_1 \in \mathbf{a}$, then $[W]\text{slide}_{\bar{\rho}/\bar{\lambda}}(S) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}_*/\bar{\lambda}_*$.

Proof. We simultaneously induct on the number of genes of family i in S . (We gloss over some technical reindexing in the arguments below.) We check the base case of one gene directly from the swapping rules of Section 7. Now let us assume that S has at least two genes of family i and the claims hold for situations with fewer genes of family i .

In the illustrative examples below that accompany the general analysis, we use for simplicity $1, 2, \dots$ to represent w_1, w_2, \dots respectively. Also, for simplicity, our examples assume \mathbf{a} is the southwest corner of $k \times (n-k)$, i.e., $\beta(\mathbf{a}) = 1 - \frac{t_1}{t_2}$.

Case (I.i).1: ($\mathbf{a}^{\rightarrow} \neq \mathbf{z}$): Consider $S = \begin{array}{|c|c|c|} \hline & \bullet & 4 \ 5 \\ \hline \bullet & 2 & 3 \\ \hline \end{array}$ and $W = \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 1 & 2 & \bullet \\ \hline \end{array}$. Then

$$\text{swap}_1(S) = (1 - \frac{t_1}{t_2}) \begin{array}{|c|c|c|} \hline & \bullet & 4 \ 5 \\ \hline 1 & 2 & 3 \\ \hline \end{array} + \frac{t_1}{t_2} \begin{array}{|c|c|c|} \hline & \bullet & 4 \ 5 \\ \hline 1 & \bullet & 3 \\ \hline \end{array} := (1 - \frac{t_1}{t_2})S' + \frac{t_1}{t_2}S''.$$

Inductively by (III), $[W]\text{slide}(\widetilde{S}') = \frac{t_4}{t_7}$. Inductively by (I.i), $[W]\text{slide}(\widetilde{S}'') = (1 - \frac{t_2}{t_3})\frac{t_4}{t_7}$. Hence $[W]\text{slide}(S) = (1 - \frac{t_1}{t_2})\frac{t_4}{t_7} + \frac{t_1}{t_2}(1 - \frac{t_2}{t_3})\frac{t_4}{t_7} = (1 - \frac{t_1}{t_3})\frac{t_4}{t_7}$, as desired. In general,

$$\begin{aligned} [W]\text{slide}(S) &= \left(1 - \hat{\beta}(\mathbf{a})\right) (-1)^{\Delta(S,W)-1} \text{wt } \bar{\alpha}_*/\bar{\lambda}_* + \hat{\beta}(\mathbf{a}) (-1)^{\Delta(S,W)-1} \left(1 - \frac{\text{wt } \bar{\alpha}/(\bar{\alpha}_* \cup \bar{\lambda})}{\hat{\beta}(\mathbf{a})}\right) \text{wt } \bar{\alpha}_*/\bar{\lambda}_* \\ &= (-1)^{\Delta(S,W)-1} (1 - \text{wt } \bar{\alpha}/(\bar{\alpha}_* \cup \bar{\lambda})) \text{wt } \bar{\alpha}_*/\bar{\lambda}_*. \end{aligned}$$

Case (I.i).2: ($\mathbf{a}^\rightarrow = \mathbf{z}$): Let $S = \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array}$ and $W = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array}$. Then

$$\text{swap}_1(S) = \left(1 - \frac{t_1}{t_2}\right) \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} := \left(1 - \frac{t_1}{t_2}\right) S'.$$

By (III), $[W]\text{slide}(\widetilde{S}') = \frac{t_3 t_6}{t_5 t_7}$. Hence $[W]\text{slide}(S) = (1 - \frac{t_1}{t_2})\frac{t_3 t_6}{t_5 t_7}$, as desired. In general,

$$[W]\text{slide}(S) = \left(1 - \hat{\beta}(\mathbf{a})\right) (-1)^{\Delta(S,W)-1} \text{wt } \bar{\alpha}_*/\bar{\lambda}_* = (-1)^{\Delta(S,W)-1} (1 - \text{wt } \bar{\alpha}/(\bar{\alpha}_* \cup \bar{\lambda})) \text{wt } \bar{\alpha}_*/\bar{\lambda}_*.$$

Case (I.ii): Let $S = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array}$ and $W = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array}$. Then $\text{swap}_1(S) = (1 - \frac{t_1}{t_2})\begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} + \frac{t_1}{t_2}\begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} := (1 - \frac{t_1}{t_2})S' + \frac{t_1}{t_2}S''$. By Lemma 10.3, $[W]\text{slide}(\widetilde{S}') = 0$. By (I.ii), $[W]\text{slide}(\widetilde{S}'') = \frac{t_2}{t_3}$. Hence $[W]\text{slide}(S) = \frac{t_1 t_2}{t_2 t_3} = \frac{t_1}{t_3}$, as desired. In general,

$$[W]\text{slide}(S) = \hat{\beta}(\mathbf{a}) (-1)^{\Delta(S,W)} \frac{1}{\hat{\beta}(\mathbf{a})} \text{wt } \bar{\alpha}/\bar{\lambda} = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda}.$$

Case (II).1: ($\mathbf{a}^\rightarrow \neq \mathbf{z}$): Let $S = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} \begin{array}{c} \bullet \\ \boxed{4} \end{array} \begin{array}{c} \bullet \\ \boxed{5} \end{array}$ and $W = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} \begin{array}{c} \bullet \\ \boxed{4} \end{array} \begin{array}{c} \bullet \\ \boxed{5} \end{array}$. Then $\text{swap}_1(S) = \frac{t_1}{t_2}\begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} \begin{array}{c} \bullet \\ \boxed{4} \end{array} \begin{array}{c} \bullet \\ \boxed{5} \end{array} := \frac{t_1}{t_2}S'$. By (II), $[W]\text{slide}(\widetilde{S}') = \frac{t_2 t_5}{t_4 t_8}$. Hence $[W]\text{slide}(S) = \frac{t_1}{t_2} \cdot \frac{t_2 t_5}{t_4 t_8} = \frac{t_1 t_5}{t_4 t_8}$, as desired. In general,

$$[W]\text{slide}(S) = \hat{\beta}(\mathbf{a}) \cdot (-1)^{\Delta(S,W)} \frac{1}{\hat{\beta}(\mathbf{a})} \text{wt } \bar{\alpha}/\bar{\lambda} = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda}.$$

Case (II).2: ($\mathbf{a}^\rightarrow = \mathbf{z}$ and the northmost $w_1 \in S$ is *not* immediately below \bullet_{i_1}): Let $S = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} \begin{array}{c} \bullet \\ \boxed{4} \end{array} \begin{array}{c} \bullet \\ \boxed{5} \end{array}$ and $W = \begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} \begin{array}{c} \bullet \\ \boxed{4} \end{array} \begin{array}{c} \bullet \\ \boxed{5} \end{array}$. Then $\text{swap}_1(S) = \frac{t_1 t_3}{t_2 t_4}\begin{array}{c} \bullet \\ \boxed{1} \end{array} \begin{array}{c} \bullet \\ \boxed{2} \end{array} \begin{array}{c} \bullet \\ \boxed{3} \end{array} \begin{array}{c} \bullet \\ \boxed{4} \end{array} \begin{array}{c} \bullet \\ \boxed{5} \end{array} := \frac{t_1 t_3}{t_2 t_4}S'$. By (II), $[W]\text{slide}(\widetilde{S}') = -\frac{t_4 t_7}{t_6 t_{10}}$. Hence $[W]\text{slide}(S) = \frac{t_1 t_3}{t_2 t_4} \cdot \left(-\frac{t_4 t_7}{t_6 t_{10}}\right) = -\frac{t_1 t_3 t_7}{t_2 t_6 t_{10}}$, as desired. In general,

$$[W]\text{slide}(S) = \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot (-1)^{\Delta(S,W)} \prod_{y:\text{label}_W(y)>1} \hat{\beta}(y) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda}.$$

Case (II).3: ($a^\rightarrow = z$ and the northmost $w_1 \in S$ is immediately below \bullet_{i_1}): Let $S = \begin{array}{cccc} & & & \bullet \\ & & & 5 \\ & & & 6 \\ \bullet & 2 & 3 & 4 \\ \bullet & 1 & & \end{array}$

and $W = \begin{array}{cccc} & & & 4 \\ & & & 5 \\ & & & 6 \\ 1 & 2 & 3 & \bullet \\ 1 & \bullet & & \end{array}$. Then

$$\text{swap}_1(S) = -\frac{t_1 t_3}{t_2 t_4} \begin{array}{cccc} & & & \bullet \\ & & & 5 \\ & & & 6 \\ 1 & 2 & 3 & 4 \\ 1 & \bullet & & \end{array} + \frac{t_1 t_3}{t_2 t_4} \begin{array}{cccc} & & & \bullet \\ & & & 5 \\ & & & 6 \\ 1 & \bullet & 3 & 4 \\ 1 & \bullet & 2 & \end{array} := -\frac{t_1 t_3}{t_2 t_4} S' + \frac{t_1 t_3}{t_2 t_4} S''.$$

By (III), $[W]\text{slide}(\tilde{S}') = \frac{t_7}{t_{10}}$. By (I.i), $[W]\text{slide}(\tilde{S}'') = \left(1 - \frac{t_4}{t_6}\right) \frac{t_7}{t_{10}}$. Hence $[W]\text{slide}(S) = -\frac{t_1 t_3 t_7}{t_2 t_4 t_{10}} + \frac{t_1 t_3}{t_2 t_4} \left(1 - \frac{t_4}{t_6}\right) \frac{t_7}{t_{10}} = -\frac{t_1 t_3 t_7}{t_2 t_6 t_{10}}$, as desired. Depending whether (I.i) or (I.ii) applies inductively, we have in general respectively

$$\begin{aligned} [W]\text{slide}(S) &= - \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot Y(-1)^{\Delta(S,W)-1} + \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot (1-Z)Y(-1)^{\Delta(S,W)-1} \\ &= (-1)^{\Delta(S,W)} Y Z \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda} \end{aligned}$$

or

$$\begin{aligned} [W]\text{slide}(S) &= \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot Z(-1)^{\Delta(S,W)} \\ &= (-1)^{\Delta(S,W)} Z \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}/\bar{\lambda}, \end{aligned}$$

where Y is the weight of the boxes of W that contain genetic labels and are North of all w_1 's and Z is the weight of the boxes of W that contain genetic labels greater than w_1 and are not North of all w_1 's.

Case (III).1: ($a \neq z$): Let $S = \begin{array}{cccc} & & & \bullet \\ & & & 5 \\ & & & 6 \\ \bullet & 2 & 3 & 4 \\ 1 & 2 & & \end{array}$ and $W = \begin{array}{cccc} & & & 4 \\ & & & 5 \\ & & & 6 \\ 2 & 3 & 4 & \bullet \\ 1 & \bullet & & \end{array}$. Then $\text{swap}_1(S) = \begin{array}{cccc} & & & \bullet \\ & & & 5 \\ & & & 6 \\ \bullet & 2 & 3 & 4 \\ 1 & 2 & & \end{array} := S'$. By (III), $[W]\text{slide}(\tilde{S}') = \frac{t_3 t_7}{t_6 t_{10}}$. Hence $[W]\text{slide}(S) = \frac{t_3 t_7}{t_6 t_{10}}$, as

desired. In general, $[W]\text{slide}(S) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}_*/\bar{\lambda}_*$.

Case (III).2: ($a = z$ and the northmost $w_1 \in S$ is *not* immediately below \bullet_{i_1}): Let $S = \begin{array}{cccc} & & & \bullet \\ & & & 4 \\ & & & 5 \\ \bullet & 1 & 2 & 3 \\ 1 & & & \end{array}$ and $W = \begin{array}{cccc} & & & 3 \\ & & & 4 \\ & & & 5 \\ 1 & 2 & 3 & \bullet \\ \bullet & & & \end{array}$. Then $\text{swap}_1(S) = -\frac{t_2}{t_3} \begin{array}{cccc} & & & \bullet \\ & & & 4 \\ & & & 5 \\ 1 & \bullet & 2 & 3 \\ \bullet & & & \end{array} := -\frac{t_2}{t_3} S'$. By (II),

$[W]\text{slide}(\tilde{S}') = -\frac{t_3 t_6}{t_5 t_9}$. Hence $[W]\text{slide}(S) = -\frac{t_2}{t_3} \cdot \left(-\frac{t_3 t_6}{t_5 t_9}\right) = \frac{t_2 t_6}{t_5 t_9}$, as desired. In general,

$$[W]\text{slide}(S) = - \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot (-1)^{\Delta(S,W)-1} \prod_{y:\text{label}_W(y)>1} \hat{\beta}(y) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}_*/\bar{\lambda}_*.$$

Case (III).3: ($a = z$ and the northmost $w_1 \in S$ is immediately below \bullet_{i_1}): Let $S = \begin{array}{cccc} & & & \bullet \\ & & & 4 \\ & & & 5 \\ \bullet & 2 & 3 & 4 \\ 1 & & & \end{array}$

and $W = \begin{array}{cccc} & & & 4 \\ & & & 5 \\ & & & \bullet \\ 1 & 2 & 3 & \bullet \\ \bullet & & & \end{array}$. Then $\text{swap}_1(S) = \frac{t_2}{t_3} \begin{array}{cccc} & & & \bullet \\ & & & 4 \\ & & & 5 \\ 1 & 2 & 3 & 4 \\ \bullet & & & \end{array} - \frac{t_2}{t_3} \begin{array}{cccc} & & & \bullet \\ & & & 4 \\ & & & 5 \\ 1 & \bullet & 3 & 4 \\ \bullet & 2 & & \end{array} := \frac{t_2}{t_3} S' - \frac{t_2}{t_3} S''$

By (III), $[W]\text{slide}(\tilde{S}') = -\frac{t_6}{t_8}$. By (I.i), $[W]\text{slide}(\tilde{S}'') = -(1 - \frac{t_3}{t_5})\frac{t_6}{t_8}$. Hence $[W]\text{slide}(S) = \frac{t_2}{t_3} \cdot \left(-\frac{t_6}{t_8}\right) - \frac{t_2}{t_3} \cdot \left(-\left(1 - \frac{t_3}{t_5}\right)\frac{t_6}{t_8}\right) = -\frac{t_2}{t_5}\frac{t_6}{t_8}$, as desired. Depending whether (I.i) or (I.ii) applies inductively, we have in general respectively


$$\begin{aligned} [W]\text{slide}(S) &= \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot (-1)^{\Delta(S,W)} Y - \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot (-1)^{\Delta(S,W)} (1-Z) Y \\ &= (-1)^{\Delta(S,W)} Y Z \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}_* / \bar{\lambda}_* \end{aligned}$$

or

$$\begin{aligned} [W]\text{slide}(S) &= - \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) \cdot (-1)^{\Delta(S,W)-1} Z \\ &= (-1)^{\Delta(S,W)} Z \prod_{x:\text{label}_W(x)=1} \hat{\beta}(x) = (-1)^{\Delta(S,W)} \text{wt } \bar{\alpha}_* / \bar{\lambda}_*, \end{aligned}$$

where Y is the weight of the boxes of W containing genetic labels and are North of all w_1 's and Z is the weight of the boxes of W containing genetic labels greater than w_1 and are not North of all w_1 's. \square

Example 11.2. Let $\lambda = (1)$, $\nu = (3, 2)$ and $\mu = (2, 1)$. Consider $U = \begin{array}{|c|c|} \hline \square & \begin{array}{|c|c|} \hline 1_1 & 1_2 \\ \hline \end{array} \\ \hline \begin{array}{|c|c|} \hline 1_1 & 2_1 \\ \hline \end{array} & \square \\ \hline \end{array} \in \Lambda$. Below, we give the reversal tree \mathfrak{T}_U .



Each edge is labeled (in blue) by $[U']\text{swap}_{i_{\mu_i}} \circ \dots \circ \text{swap}_{i_1}(V')$ where U' is the parent of the i_1 -good tableau V' . This label agrees with the application of Claim 11.1 to each i -walkway of V' . Below each leaf (in red) is the coefficient in Λ^+ (i.e., $(-1)^{|\rho/\lambda|+1}$ if nonzero). \square

Lemma 11.3. *Suppose U' is an $(i+1)_1$ -good node of \mathfrak{T}_U . Let Γ be the boxes of U containing labels of family i . Then $\sum_{V'} (-1)^{1 + \#\bullet\text{'s in } V'} [U']\text{swap}_{i_{\mu_i}} \circ \dots \circ \text{swap}_{i_1}(V') = (-1)^{1 + \#\bullet\text{'s in } U'} \text{wt } \Gamma$, where the sum is over all children V' of U' in \mathfrak{T}_U .*

Proof. Consider boxes of U' containing unmarked labels of family i or $\bullet_{(i+1)_1}$. By (W.1) and (W.2), these boxes decompose into an edge-disjoint union of i -walkways W_1, W_2, \dots, W_t .

Let Γ_j be the boxes of W_j in U containing labels of family i ; thus $\Gamma = \Gamma_1 \sqcup \Gamma_2 \sqcup \cdots \sqcup \Gamma_t$. Let R_j and R'_j (if it exists) be the reversal(s) defined by Lemmas 10.3 and 10.4 with respect to the walkway W_j . As computed by Claim 11.1, let a_j be the coefficient of W_j obtained by sliding R_j . Let b_j be the coefficient of W_j obtained by sliding R'_j if it exists; set $b_j := 0$ if R'_j does not exist. We now assert that

$$(11.1) \quad (-1)^{\#\bullet\text{'s in } R_j} a_j + (-1)^{\#\bullet\text{'s in } R'_j} b_j = (-1)^{\#\bullet\text{'s in } W_j} \text{wt } \Gamma_j.$$

Suppose there is a unique reversal (i.e., $b_j = 0$). This occurs under Lemma 10.3(I,II) and Lemma 10.4(I,II). In these four cases, R_j is respectively the S from (II), (I.ii), (III) and (II) of Claim 11.1. Hence in each of these cases, (11.1) is immediate from the apposite case of Claim 11.1 (note that for Lemma 10.4(I), the southmost row of W_j has a single box and $\bar{\alpha}_*/\bar{\lambda}_* = \bar{\alpha}/\bar{\lambda} = \Gamma_j$). Suppose there are two reversals. This occurs under Lemma 10.3(III) and Lemma 10.4(III), which show that R_j is the S from Claim 11.1(III) and R'_j is the S from Claim 11.1(I.i). Hence (11.1) also follows in these cases, by adding the two apposite coefficients given by Claim 11.1.

Since by Proposition 10.5 all V' are obtained by independent replacements of W_j by R_j and R'_j (if it exists),

$$\begin{aligned} \sum_{V'} (-1)^{1 + \#\bullet\text{'s in } V'} [U'] \text{swap}_{i_{\mu_i}} \circ \cdots \circ \text{swap}_{i_1}(V') &= - \prod_{j=1}^t \left((-1)^{\#\bullet\text{'s in } R_j} a_j + (-1)^{\#\bullet\text{'s in } R'_j} b_j \right) \\ &= - \prod_{j=1}^t (-1)^{\#\bullet\text{'s in } W_j} \text{wt } \Gamma_j \\ &= (-1)^{1 + \#\bullet\text{'s in } U'} \text{wt } \Gamma. \end{aligned}$$

□

Lemma 11.4. *Let U' be an $(i+1)_1$ -good node of \mathfrak{T}_U . Let $\Gamma^{(i)}$ be the set of boxes $\{x \in \alpha/\lambda : \text{family}(\text{label}_U(x)) \leq i\}$. Then*

$$\sum_T (-1)^{1 + \#\bullet\text{'s in } T} [U'] \text{swap}_{i_{\mu_i}} \circ \text{swap}_{i_{\mu_i}^-} \circ \cdots \circ \text{swap}_{1_1^+} \circ \text{swap}_{1_1}(T) = \text{wt}(\Gamma^{(i)}) (-1)^{1 + \#\bullet\text{'s in } U'},$$

where the sum is over all $T \in \text{leaf}(\mathfrak{T}_{U'})$ that are descendants of U' .

Proof. We induct on $i \geq 0$. In the base case $i = 0$, $U' = T$ for $T \in \text{leaf}(\mathfrak{T}_{U'})$ and the lefthand side equals $(-1)^{1 + \#\bullet\text{'s in } T}$. This equals the righthand side since $\Gamma^{(0)} = \emptyset$ so $\text{wt } \Gamma^{(0)} = 1$.

Now let $i > 0$. We have $\sum_T (-1)^{1 + \#\bullet\text{'s in } T} [U'] \text{swap}_{i_{\mu_i}} \circ \text{swap}_{i_{\mu_i}^-} \circ \cdots \circ \text{swap}_{1_1^+} \circ \text{swap}_{1_1}(T)$

$$\begin{aligned} &= \sum_{V' \text{ a child of } U'} \sum_{T \in \text{leaf}(\mathfrak{T}_{V'}) \text{ below } V'} (-1)^{1 + \#\bullet\text{'s in } T} [U'] \text{swap}_{i_{\mu_i}} \circ \cdots \circ \text{swap}_{i_1} \circ \\ &\quad \text{swap}_{(i-1)_{\mu_{i-1}}} \circ \cdots \circ \text{swap}_{1_1}(T) \\ &= \sum_{V' \text{ a child of } U'} \sum_{T \in \text{leaf}(\mathfrak{T}_{V'}) \text{ below } V'} (-1)^{1 + \#\bullet\text{'s in } T} [U'] \text{swap}_{i_{\mu_i}} \circ \cdots \circ \text{swap}_{i_1}(V') \cdot \\ &\quad [V'] \text{swap}_{(i-1)_{\mu_{i-1}}} \circ \cdots \circ \text{swap}_{1_1}(T). \end{aligned}$$

The previous equality is since \mathfrak{T}_U is a tree (Proposition 10.6) and V' is the unique child of U' that is an ancestor of T . The previous summation equals

$$\begin{aligned}
& \sum_{V' \text{ a child of } U'} [U'] \text{swap}_{i_{\mu_i}} \circ \cdots \circ \text{swap}_{i_1}(V') \sum_{T \in \text{leaf}(\mathfrak{T}_{U'}) \text{ below } V'} (-1)^{1 + \#\bullet\text{'s in } T} [V'] \text{swap}_{(i-1)\mu_{i-1}} \circ \cdots \circ \text{swap}_{1_1}(T) \\
&= \sum_{V' \text{ a child of } U'} [U'] \text{swap}_{i_{\mu_i}} \circ \cdots \circ \text{swap}_{i_1}(V') \cdot \text{wt}(\Gamma^{(i-1)}) (-1)^{1 + \#\bullet\text{'s in } V'} \quad (\text{by induction}) \\
&= \text{wt}(\Gamma^{(i-1)}) \sum_{V' \text{ a child of } U'} (-1)^{1 + \#\bullet\text{'s in } V'} [U'] \text{swap}_{i_{\mu_i}} \circ \cdots \circ \text{swap}_{i_1}(V') \\
&= \text{wt}(\Gamma^{(i-1)}) \cdot (-1)^{1 + \#\bullet\text{'s in } U'} \text{wt}(\Gamma) \quad (\text{by Lemma 11.3}) \\
&= (-1)^{1 + \#\bullet\text{'s in } U'} \text{wt}(\Gamma^{(i)}),
\end{aligned}$$

since by definition $\text{wt}(\Gamma^{(i)}) = \text{wt}(\Gamma) \cdot \text{wt}(\Gamma^{(i-1)})$. \square

Proposition 11.5. For $U \in B_{\lambda, \mu}^\alpha$

$$(11.2) \quad \sum_{T \in \text{leaf}(\mathfrak{T}_U)} (-1)^{|\rho(T)/\lambda|+1} [U] \text{slide}_{\rho(T)/\lambda}(T) = \text{wt}(\alpha/\lambda) (-1)^{|\nu/\alpha|+1}$$

where $\rho(T) \in \{\lambda\} \cup \lambda^+$ is the “inner shape” of T , i.e., T has shape $\nu/\rho(T)$.

Proof. Take $U' = U$ in Lemma 11.4. \square

Now assume $U \in B_{\lambda, \mu}^\nu$. The root of \mathfrak{T}_U contains no $\bullet_{(\ell(\mu)+1)_1}$'s. One leaf of \mathfrak{T}_U is U itself. This is the unique leaf not in Λ^+ . Let $\text{leaf}^*(\mathfrak{T}_U)$ be the collection of all other leaves.

Proposition 11.6. For $U \in B_{\lambda, \mu}^\nu$

$$(11.3) \quad \sum_{T \in \text{leaf}^*(\mathfrak{T}_U)} (-1)^{|\rho(T)/\lambda|+1} [U] \text{slide}_{\rho(T)/\lambda}(T) = 1 - \text{wt}(\nu/\lambda)$$

where $\rho(T) \in \lambda^+$ is the “inner shape” of T , i.e., T has shape $\nu/\rho(T)$.

Proof. This is immediate from Proposition 11.5, since $\nu = \alpha$ and the contribution from the excluded leaf is 1. \square

Example 11.7. In Example 11.2, summing the weights below the left child of U gives $1 - \frac{t_1 t_3}{t_2 t_5}$, in agreement with Lemma 11.3. Proposition 11.6 asserts in this case that

$$1 - \text{wt}(\nu/\lambda) = 1 - \frac{t_1}{t_5} = \left(1 - \frac{t_1}{t_2}\right) + \left(1 - \frac{t_3}{t_5}\right) - \left(1 - \frac{t_1}{t_2}\right) \left(1 - \frac{t_3}{t_5}\right) + \frac{t_1 t_3}{t_2 t_5} \cdot \left(1 - \frac{t_2}{t_3}\right),$$

as the reader may verify. \square

Recall $\Lambda^+ = \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|+1} \sum_{T \in B_{\rho, \mu}^\nu} T$. For $T \in B_{\rho, \mu}^\nu$, write $T^{(1_1)}$ (cf. Section 7.2) for T with \bullet_{1_1} in each box of ρ/λ .

Now set

$$(11.4) \quad P_{\mathcal{G}} := \sum_{\rho \in \lambda^+} (-1)^{|\rho/\lambda|-1} \sum_{T \in B_{\rho, \mu}^\nu} \text{swap}_{g^-} \circ \text{swap}_{(g^-)^-} \circ \cdots \circ \text{swap}_{1_1}(T^{(1_1)}).$$

In particular, P_{1_1} is Λ^+ where each T is replaced by $T^{(1_1)}$. By Lemma 7.6 and Proposition 7.7, each $P_{\mathcal{G}}$ is a formal sum of \mathcal{G} -good tableaux.

The main conclusion of this section is

Proposition 11.8. $P_{\mathcal{G}_{\max}^+}$ with all $\bullet_{\mathcal{G}_{\max}^+}$'s removed equals $\Lambda + \Lambda^-$.

Proof. By Corollary 7.11 each tableau appearing in $P_{\mathcal{G}_{\max}^+}$ (with $\bullet_{\mathcal{G}_{\max}^+}$'s removed) is a tableau in $\Lambda + \Lambda^-$. On the other hand, given any U appearing in $\Lambda + \Lambda^-$, we constructed the tree \mathfrak{T}_U in Section 10. By Proposition 10.7, the leaves of \mathfrak{T}_U are exactly those tableaux $T \in \Lambda^+$ such that $U \in \text{slide}_{\rho/\lambda}(T)$. It remains to show that $[U]P_{\mathcal{G}_{\max}^+} = 1 - \text{wt}(\nu/\lambda)$ if $U \in \Lambda$ and $[U]P_{\mathcal{G}_{\max}^+} = (-1)^{|\nu/\delta|+1} \text{wt}(\delta/\lambda)$ if $U \in \Lambda^-$ and the shape of U is δ/λ . These are precisely the statements of Propositions 11.6 and 11.5, respectively. \square

12. WEIGHT PRESERVATION

12.1. Fine tableaux and their weights. A tableau is **fine** if it is good or can be obtained from a good tableau by swapping some subset of its snakes, i.e. it appears in the formal sum of tableaux resulting from this partial swap.

Let T be fine and fix $x \in T$. Suppose $\ell \in \underline{x}$. Define $\text{edgefactor}(\ell)$ as in Section 1.4; see (1.1). The **edge weight** $\text{edgewt}(T) := \prod_{\ell} \text{edgefactor}(\ell)$, where the product is over all (non-virtual) edge labels of T .

Suppose T is obtained by swapping some of the snakes of the good tableau S and U is obtained from T by swapping the remaining snakes. We define the positions in T of a **virtual label** (\mathcal{H}) as follows. Consider a box x in column c . If c intersects a snake in S that has been swapped in T , and that snake is not the upper snake described in Lemma 6.6(III), then $(\mathcal{H}) \in \underline{x}$ (in T) if and only if $(\mathcal{H}) \in \underline{x}$ (in U). Otherwise, $(\mathcal{H}) \in \underline{x}$ (in T) if and only if $(\mathcal{H}) \in \underline{x}$ (in S). Observe that if T is indeed good, this definition is clearly consistent with the definition of virtual labels in a good tableau.

Suppose $(\mathcal{H}) \in \underline{x}$. If $\text{label}_T(x)$ is marked and each $\mathcal{F} \in \underline{x}$ with $\mathcal{F} \prec \mathcal{G}$ is marked, then

$$(12.1) \quad \text{virtualfactor}_{\underline{x} \in T}((\mathcal{H})) := -\text{edgefactor}_{\underline{x} \in T}(\mathcal{H}) = \frac{t_{\text{Man}(x)}}{t_{r+N_{\mathcal{H}}+1-\text{family}(\mathcal{H})+\text{Man}(x)}} - 1.$$

Otherwise

$$(12.2) \quad \text{virtualfactor}_{\underline{x} \in T}((\mathcal{H})) := 1 - \text{edgefactor}_{\underline{x} \in T}(\mathcal{H}) = \frac{t_{\text{Man}(x)}}{t_{r+N_{\mathcal{H}}+1-\text{family}(\mathcal{H})+\text{Man}(x)}}.$$

The **virtual weight** $\text{virtualwt}(T)$ is $\prod_{(\mathcal{L})} \text{virtualfactor}((\mathcal{L}))$, where the product is over all instances of virtual labels.

Call $x \in T$ **productive** if any of the following hold:

- (P.1) $\text{label}_T(x) < \text{label}_T(x^{\rightarrow})$ or $x^{\rightarrow} \notin T$;
- (P.2) $\bullet_{i_{k+1}} \in x, i_k \in x^{\leftarrow}, i_{k+1} \in \underline{x}$, and either $\text{family}(\text{label}(x^{\rightarrow})) \neq i$ or $x^{\rightarrow} \notin T$;
- (P.3) $\mathcal{H} \in x, \bullet_{\mathcal{G}} \in x^{\rightarrow}$, and $\underline{x^{\rightarrow}}$ does not contain a label of the same family as \mathcal{H} ; or
- (P.4) $i_k \in x, i_{k+1} \in x^{\rightarrow}$ and $\bullet_{i_{k+1}} \in x^{\rightarrow\uparrow}$, with x not SouthEast of a $\bullet_{i_{k+1}}$.

Define $\text{boxfactor}(x)$ and **box weight** $\text{boxwt}(T) = \prod_x \text{boxfactor}(x)$ as in Section 1.4, specifically (1.2), with the addendum that $\bullet_{\mathcal{H}} \in x$ is evaluated like $\mathcal{H} \in x$.

Example 12.1.

- The right two boxes of $\boxed{1_1 1_2 2_1}$ are productive by (P.1). The left box is not productive.
- The left box of $\boxed{1_1 \bullet_{1_2}}_{1_2}$ is not productive. The right box is productive by (P.2).
- The first and third boxes of $\boxed{1_1 \bullet_{1_2} 2_1}$ are productive by (P.3) and (P.1) respectively. The middle box is not productive; a box with \bullet_{1_2} is productive only if (P.2) holds.
- The right box of the second row in both $\begin{array}{c} \bullet_{1_2} \\ \boxed{1_1 1_2} \end{array}$ and $\begin{array}{c} \bullet_{1_1} \\ \boxed{1_1 1_2} \end{array}$ is productive by (P.1). The left box in the second row is productive only in the first case, by (P.4). \square

Finally the **weight** is

$$\text{wt}(T) := (-1)^{d(T)} \text{edgewt}(T) \cdot \text{virtualwt}(T) \cdot \text{boxwt}(T),$$

where $d(T) = \sum_{\mathcal{G}} (|\mathcal{G}| - 1)$, the sum is over genes \mathcal{G} , and $|\mathcal{G}|$ is the (multiset) cardinality of \mathcal{G} (not including virtual labels). We will view wt as a $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm n}]$ -linear operator of formal sums of tableaux.

By Lemma 5.3, bundled tableaux are good and hence also fine. Hence for a bundled tableau B , we have two *a priori* distinct notions of $\text{wt } B$. The following lemma justifies our failure to distinguish these notationally:

Lemma 12.2. *For B a bundled tableau, $\text{wt } B$ as a fine tableau equals $\text{wt } B$ as a bundled tableau.*

Proof. By definition, the two notions of $\text{edgewt}(B)$ coincide, as do the two notions of $d(B)$. Since B has no \bullet 's, only (P.1) is available to effect productivity. Hence the two notions of productive boxes coincide, and thus, by definition, so too do the two notions of $\text{boxwt}(B)$. As remarked above, the locations of virtual labels are the same, whether we think of B as bundled or fine. By Lemma 2.4, $\text{wt } B$ as a bundled tableau is

$$(-1)^{d(B)} \text{edgewt}(B) \text{boxwt}(B) \prod_{\textcircled{\ell}} (1 - \text{edgefactor}(\ell)),$$

where the product is over all instances of virtual labels and $\text{edgefactor}(\ell)$ means the factor that would be given by ℓ in $\textcircled{\ell}$'s place. Since B is bundled, it has no marked labels. Hence $\text{virtualwt}(B)$ is calculated using only (12.2), not (12.1). Thus $\text{virtualwt}(B) = \prod_{\textcircled{\ell}} (1 - \text{edgefactor}(\ell))$, and the lemma follows. \square

12.2. Main claim about weight preservation.

Proposition 12.3.

- (I) $\text{wt } P_{1_1} = \text{wt } \Lambda^+$.
- (II) For every \mathcal{G} , $\text{wt } P_{\mathcal{G}} = \text{wt } P_{1_1}$.
- (III) $\text{wt } P_{\mathcal{G}_{\max}^+} = \text{wt } \Lambda + \text{wt } \Lambda^-$.

Proof. We will first prove the easier statements (I) and (III).

(I): Suppose $T \in B_{\rho, \mu}^{\nu}$ for some $\rho \in \lambda^+$. It is enough to show $\text{wt } T = \text{wt } T^{(1_1)}$. Certainly $\text{edgewt}(T) = \text{edgewt}(T^{(1_1)})$ and $d(T) = d(T^{(1_1)})$. Adding \bullet_{1_1} 's preserves the virtual labels' locations, so $\text{virtualwt}(T) = \text{virtualwt}(T^{(1_1)})$.

A productive box in T is also productive in $T^{(1_1)}$ and has the same boxfactor. Suppose x is a productive box of $T^{(1_1)}$ that is not productive in T . It satisfies one of (P.1)–(P.4). If x satisfies (P.1), it is productive in T . If it satisfies (P.2), then $\bullet_{1_1} \in x$ and x^{\leftarrow} contains a label, contradicting $x^{\leftarrow} \in \rho$. If it satisfies (P.3), then $x^{\rightarrow} \in \rho$, contradicting that x contains a label. Finally if x satisfies (P.4), then $\bullet_{i_{k+1}} \in x^{\rightarrow\uparrow}$ and $i_k \in x$. But every \bullet in $T^{(1_1)}$ is \bullet_{1_1} . Hence $i_{k+1} = 1_1$, which is impossible since 1_0 is not a label in our alphabet. Thus the productive boxes of T and $T^{(1_1)}$ are the same, and with the same respective boxfactors. Therefore, $\text{wt } T = \text{wt } T^{(1_1)}$.

(III): Suppose $U \in P_{\mathcal{G}_{\max}^+}$ and let \tilde{U} be given by deleting each $\bullet_{\mathcal{G}_{\max}^+}$. Proposition 11.8 states $P_{\mathcal{G}_{\max}^+}$ with all $\bullet_{\mathcal{G}_{\max}^+}$'s removed equals $\Lambda + \Lambda^-$. Thus, it suffices to show $\text{wt } U = \text{wt } \tilde{U}$. Clearly, $\text{edgewt}(U) = \text{edgewt}(\tilde{U})$ and $d(U) = d(\tilde{U})$. One checks that the virtual labels of U and the virtual labels of \tilde{U} appear in the same places. Hence $\text{virtualwt}(U) = \text{virtualwt}(\tilde{U})$.

Suppose x is productive in U . Then it satisfies one of (P.1)–(P.4). If x satisfies (P.1) in U , then it satisfies (P.1) in \tilde{U} . Now x cannot satisfy (P.2) in U , since if it did, $\bullet_{\mathcal{G}_{\max}^+} \in x$ and x contains a label, contradicting Lemma 7.9. If x satisfies (P.3) in U , then it satisfies (P.1) in \tilde{U} . If x satisfies (P.4) in U , then $\bullet_{\mathcal{G}_{\max}^+} \in x^{\rightarrow\uparrow}$ but is not an outer corner, again contradicting Lemma 7.9. Thus if x is productive in U , it is productive in \tilde{U} . Conversely, if x is productive in \tilde{U} , it satisfies (P.1), since there are no $\bullet_{\mathcal{G}_{\max}^+}$'s in \tilde{U} . Hence x satisfies (P.1) or (P.3) in U . Thus the productive boxes of U and \tilde{U} are the same. These boxes have the same boxfactors. Thus $\text{boxwt}(U) = \text{boxwt}(\tilde{U})$.

(II): We induct on \mathcal{G} with respect to \prec . The base case $\mathcal{G} = 1_1$ is trivial. The inductive hypothesis is that $\text{wt } P_{\mathcal{G}} = \text{wt } P_{1_1}$. Our inductive step is to show $\text{wt } P_{\mathcal{G}^+} = \text{wt } P_{\mathcal{G}}$.

Consider the set

$$\text{Snakes}_{\mathcal{G}} = \{S \text{ is a snake in } T: [T]P_{\mathcal{G}} \neq 0\}.$$

We emphasize that each $S \in \text{Snakes}_{\mathcal{G}}$ refers to a particular instance of a snake in a specific tableau $T \in P_{\mathcal{G}}$. In particular, $\text{Snakes}_{\mathcal{G}}$ is not a multiset.

For $\mathcal{B} \subseteq \text{Snakes}_{\mathcal{G}}$ define $\text{swapset}_{\mathcal{B}}(T)$ to be the formal sum of fine tableaux obtained by swapping each snake of \mathcal{B} that appears in T (done in any order, as permitted by Lemma 7.5).

We will construct m subsets \mathcal{B}_i such that (D.1) and (D.2) below hold:

(D.1) We have a disjoint union $\text{Snakes}_{\mathcal{G}} = \bigsqcup_{1 \leq i \leq m} \mathcal{B}_i$.

(D.2) For every $1 \leq i \leq m$ and $J \subseteq \{1, \dots, \hat{i}, \dots, m\}$, let $\mathcal{B}_J := \cup_{j \in J} \mathcal{B}_j$. Then

$$(12.3) \quad \sum_{T \in \Gamma_i} [T]P_{\mathcal{G}} \cdot \text{wt}(\text{swapset}_{\mathcal{B}_J}(T)) = \sum_{T \in \Gamma_i} [T]P_{\mathcal{G}} \cdot \text{wt}(\text{swapset}_{\mathcal{B}_i} \circ \text{swapset}_{\mathcal{B}_J}(T)),$$

where $\Gamma_i := \{T \in P_{\mathcal{G}}: T \text{ contains a snake from } \mathcal{B}_i\}$.

Claim 12.4. *The existence of $\{\mathcal{B}_i\}$ satisfying (D.1) and (D.2) implies $\text{wt}(P_{\mathcal{G}^+}) = \text{wt}(P_{\mathcal{G}})$.*

Proof. By Lemma 7.5, snakes may be swapped in any order, so choose an arbitrary ordering of the blocks \mathcal{B}_i . By (D.1), $P_{\mathcal{G}^+} := \text{swap}_{\mathcal{G}}(P_{\mathcal{G}}) = \text{swapset}_{\mathcal{B}_m} \circ \dots \circ \text{swapset}_{\mathcal{B}_1}(P_{\mathcal{G}})$.

Thus

$$\begin{aligned} \text{wt}(P_{\mathcal{G}^+}) &= \text{wt}(\text{swap}_{\mathcal{G}}(P_{\mathcal{G}})) \\ &= \text{wt}(\text{swapset}_{\mathcal{B}_m} \circ \cdots \circ \text{swapset}_{\mathcal{B}_1}(P_{\mathcal{G}})) \\ &= \text{wt}(\text{swapset}_{\mathcal{B}_{m-1}} \circ \cdots \circ \text{swapset}_{\mathcal{B}_1}(P_{\mathcal{G}})) \end{aligned}$$

Here we have just used (12.3) from (D.2) together with linearity of wt and $\text{swapset}_{\mathcal{B}_i}$ and the triviality $\text{swapset}_{\mathcal{B}_j}(T) = \text{swapset}_{\mathcal{B}_i} \circ \text{swapset}_{\mathcal{B}_j}(T)$ for $T \notin \Gamma_i$. Repeating this argument $m - 1$ further times, we obtain the desired equality with $\text{wt}(P_{\mathcal{G}})$. \square

In order to provide the desired decomposition, we need to first construct certain “pairing” maps. These are given in Section 12.3. Given these, the description of the decomposition satisfying (D.1) and (D.2) is relatively straightforward and is found in Appendix C. \square

12.3. Pairing maps. Let $G_{\lambda, \mu}^{\nu}(\mathcal{G})$ be the set of \mathcal{G} -good tableaux of shape ν/λ and content μ . For $\mathcal{Q} \prec \mathcal{G}$ and $T \in G_{\lambda, \mu}^{\nu}(\mathcal{G})$, let $\mathfrak{R}_{\mathcal{Q}}(T) := \{V \in \text{revswap}_{\mathcal{Q}^+} \circ \cdots \circ \text{revswap}_{\mathcal{G}}(T)\}$.

Lemma 12.5. *For any genes $\mathcal{Q} \prec \mathcal{G}$ and any tableau $T \in G_{\lambda, \mu}^{\nu}(\mathcal{G})$*

$$\mathfrak{R}_{\mathcal{Q}}(T) = \{W \in G_{\lambda, \mu}^{\nu}(\mathcal{Q}) : T \in \text{swap}_{\mathcal{G}^-} \circ \cdots \circ \text{swap}_{\mathcal{Q}}(W)\}.$$

Proof. This is immediate from Proposition 9.8, noting that, by Lemmas 7.8 and 9.4 and Propositions 7.7 and 9.5, both forward and reverse swaps preserve goodness and content. \square

Let \mathcal{S}_1 be the subset of tableaux in $G_{\lambda, \mu}^{\nu}(i_k)$ with a box x such that for some $\ell \geq k$, $\bullet_{i_k} \in x$,

$\bullet_{i_k} \in x^{\rightarrow\uparrow}$, $i_{\ell+1} \in x^{\rightarrow}$ and $(i_{\ell}) \in \underline{x}$, i.e. locally the tableau is $\mathcal{C}_1 = \begin{array}{c} \bullet_{i_k} \\ \bullet_{i_k} \ i_{\ell+1} \\ (i_{\ell}) \end{array}$ (with possibly additional edge labels), where x southwestmost depicted box. Let \mathcal{S}'_1 be the subset of tableaux in $G_{\lambda, \mu}^{\nu}(i_k)$ with a box x such that $i_{\ell} \in x$, $i_{\ell+1} \in x^{\rightarrow}$, $\bullet_{i_k} \in x^{\rightarrow\uparrow}$, i_{ℓ} appears outside of

x and no \bullet_{i_k} appears West of x in x 's row. Locally the tableau is $\mathcal{C}'_1 = \begin{array}{c} \bullet_{i_k} \\ i_{\ell} \ i_{\ell+1} \end{array}$ (with possibly additional edge labels).

Lemma 12.6. *If $T \in \mathcal{S}_1$ (respectively, \mathcal{S}'_1), there is a unique \mathcal{C}_1 (respectively, \mathcal{C}'_1) that it contains.*

Proof. Let x be the lower-left box of any fixed choice of \mathcal{C}_1 in T . Since $(i_{\ell}) \in \underline{x}$, the $i_{\ell+1} \in x^{\rightarrow}$ is westmost in T by (G.6). Hence this configuration is unique. The argument for the other claim is the same, except we replace “ $(i_{\ell}) \in \underline{x}$ ” with “ $i_{\ell} \in x$ ”. \square

For $T \in \mathcal{S}_1$, let $\phi_1(T)$ to be the same tableau with the unique \mathcal{C}_1 replaced by \mathcal{C}'_1 . (By this we mean that we delete the labels specified in \mathcal{C}_1 and add the labels specified in \mathcal{C}'_1 ; any additional edge labels in T are unchanged.)

Lemma 12.7. $\phi_1 : \mathcal{S}_1 \rightarrow \mathcal{S}'_1$ is a bijection.

Proof. Let $\phi_1^{-1} : \mathcal{S}'_1 \rightarrow \mathcal{S}_1$ be the putative inverse of ϕ_1 , defined by replacing \mathcal{C}'_1 in a $T \in \mathcal{S}'_1$ by \mathcal{C}_1 . We are done once we show that ϕ_1 and ϕ_1^{-1} are well-defined since the maps are clearly injective and are mutually inverse.

Let x be the southwestmost box in the unique (by Lemma 12.6) \mathcal{C}_1 in T .

(ϕ_1 is well-defined): Let $T \in \mathcal{S}_1$. We only need that $\phi_1(T)$ is good. Conditions (G.1) and (G.2) hold trivially in $\phi_1(T)$. (G.3) holds if x^\leftarrow is empty. Suppose $\mathcal{F} \in x^\leftarrow$. By T 's (G.9), $\mathcal{F} \prec i_k$. Hence $\mathcal{F} \prec i_\ell$, and (G.3) holds in $\phi_1(T)$. The $(i_\ell) \in \underline{x}$ in T shows that $\phi_1(T)$ satisfies (G.4), (G.6) and (G.8). (G.5), (G.7), (G.9), (G.11) hold trivially. Since $i_{\ell+1} \in x^\rightarrow$ is not marked, by Lemma 5.4(II) there is no marked label in T in x 's row, so (G.10) and (G.13) hold for $\phi_1(T)$. For (G.12), suppose T has labels ℓ, ℓ' that violate (G.12) in $\phi_1(T)$. Since ℓ must be northWest of x , by T 's (G.9), $\ell \prec i_k$. Since ℓ' must be southeast of x , by T 's (G.3), (G.4) and (G.11), $\ell' \succ i_{\ell+1}$. Hence $\text{family}(\ell) = \text{family}(\ell') = i$. If ℓ is North of x , then by (G.4) the box of x 's row directly below ℓ contains a label that violates T 's (G.9). By T 's (G.4), ℓ' is not South of x^\rightarrow . Hence ℓ, ℓ' are box labels in the row of x , and no violation of $\phi_1(T)$'s (G.12) occurs.

(ϕ_1^{-1} is well-defined): Let $T' \in \mathcal{S}'_1$. We must show that (G.7) and (G.13) hold in $\phi_1^{-1}(T')$ and that (G.1)–(G.6) and (G.8)–(G.12) hold even if the virtual label is replaced by a non-virtual one (cf. (V.1)–(V.3)). (G.1), (G.3)–(G.10), (G.12) and (G.13) are trivial to verify. To verify (G.2) for $\phi_1^{-1}(T')$, it suffices to show T' has no \bullet_{i_k} South of x in the same column, or West of x in the same row. (G.9) for T' rules out the possibility of \bullet_{i_k} South of x in the same column of T' . By definition, there is no \bullet_{i_k} West of x in the same row. To see (G.11) for $\phi_1^{-1}(T')$, we check there is no marked label \mathcal{F}^\dagger in the column of x . Such a label cannot appear North of x in T' by Lemma 5.4 and (G.2), considering the $\bullet_{i_k} \in x^{\rightarrow\dagger}$. By (G.4), it cannot appear South of x in T' either. \square

Proposition 12.8. *For each $T \in \mathcal{S}_1$, $[T]P_{i_k} = -[\phi_1(T)]P_{i_k}$.*

Proof. Let $T^\dagger := \phi_1(T)$.

Special case $k = 1$: Let \tilde{T} be the tableau obtained from T by deleting:

- all labels of family i and greater;
- all marked labels; and
- all boxes containing a deleted box label.

Notice that any label SouthEast of a deleted label or a \bullet_{i_1} will have been deleted.

As well we reindex the genes so that the subscripts of each family form an initial segment of $\mathbb{Z}_{>0}$. (This reindexing is only possibly needed if T contained a marked label.) We leave \bullet_{i_1} 's in place. In the same way, produce \tilde{T}^\dagger from T^\dagger . By definition of ϕ_1 , \tilde{T} has one more \bullet_{i_1} than \tilde{T}^\dagger and otherwise the two tableaux are exactly the same (the family i labels of \mathcal{C}_1 and \mathcal{C}'_1 having been deleted).

Ignoring \bullet_{i_1} 's, $\tilde{T}, \tilde{T}^\dagger$ are of some common skew shape θ/λ . If we include the \bullet_{i_1} 's, their respective total shapes are some ω/λ and ω^\dagger/λ where $\omega, \omega^\dagger \in \theta^+$.

Claim 12.9. *$\tilde{T} \in G_{\lambda, \tilde{\mu}}^\omega(i_1)$ and $\tilde{T}^\dagger \in G_{\lambda, \tilde{\mu}}^{\omega^\dagger}(i_1)$ where $\tilde{\mu}$ is a partition (e.g., if T has no marked labels then $\tilde{\mu} := (\mu_1, \mu_2, \dots, \mu_{i-1})$). Thus, \tilde{T} and \tilde{T}^\dagger (with \bullet_{i_1} 's removed) are in $B_{\lambda, \tilde{\mu}}^\theta$.*

Proof. We prove the claim for \tilde{T} ; the proof for \tilde{T}^\dagger is essentially the same.

Clearly, (G.1)–(G.7), (G.9) and (G.12) for \tilde{T} are inherited from the assumption T is good. (G.10), (G.11) and (G.13) are vacuous for \tilde{T} . It remains to show (G.8) holds for \tilde{T} (which moreover implies $\tilde{\mu}$ is a partition).

Suppose \tilde{T} fails (G.8). Then there is a least q such that \tilde{T} has a ballotness violation between families q and $q + 1$. That is, in some genotype G of \tilde{T} there are more labels of family $q + 1$ than of family q in some initial segment of $\text{word}(G)$. Since we have deleted all labels of family i and greater, $q < i - 1$. By failure of (G.8), either there exist labels q_r and $(q + 1)_s$ of \tilde{T} with $N_{q_r} = N_{(q+1)_s}$ such that $(q + 1)_s$ appears before q_r in $\text{word}(G)$, or else there is a label $(q + 1)_s$ of \tilde{T} with $N_{(q+1)_s} > N_{q_v}$ for all v . Let $q_{r'}$ (if q_r exists) and $(q + 1)_{s'}$ be the corresponding labels of T . We assert in the former case that $N_{q_{r'}} \leq N_{(q+1)_{s'}}$ in T . In the latter case, we assert $N_{(q+1)_{s'}} > N_{q_{v'}}$ in T for all v' . Either of these inequalities contradicts T 's (G.8).

To see these assertions, suppose that q_h is a gene of T that is entirely deleted in the construction of \tilde{T} (i.e. every instance of q_h in T is marked). Consider an instance of q_h in T in \underline{x} or \underline{x} . Since this q_h is marked and $q < i - 1$, by Lemma 5.5 we know T has some nonvirtual and marked $(q + 1)_z \in \underline{x}$ with $N_{q_h} = N_{(q+1)_z}$. By T 's (G.7), there is no $(q + 1)_z$ West of \underline{x} in T . By T 's Lemma 5.6, there is no $(q + 1)_z$ East of \underline{x} in T . Hence the $(q + 1)_z \in \underline{x}$ is the only $(q + 1)_z$ in T . Since it is marked, the gene $(q + 1)_z$ is entirely deleted in \tilde{T} . By this argument, if $q_{\tilde{h}}$ is any other gene of T that is entirely deleted in the construction of \tilde{T} , there is a distinct $(q + 1)_{\tilde{z}}$ with $N_{q_{\tilde{h}}} = N_{(q+1)_{\tilde{z}}}$ that is also entirely deleted in \tilde{T} . Hence $N_{q_{r'}} \geq N_{(q+1)_{s'}}$ or $N_{(q+1)_{s'}} > N_{q_{v'}}$ in T for all v' , as asserted.

The last sentence of the claim follows from the first by Lemma 7.10, since no genetic label is southeast of a \bullet_{i_1} . \square

In view of Claim 12.9, it makes sense to speak of $\mathfrak{S}_{\tilde{T}}$ and of $\mathfrak{S}_{\tilde{T}^\dagger}$. By Proposition 11.5,

$$(12.4) \quad \sum_{L \in \text{leaf}(\mathfrak{S}_{\tilde{T}})} (-1)^{|\rho(L)/\lambda|+1} [\tilde{T}] \text{slide}_{\rho(L)/\lambda}(L) = (-1)^{1+\#\text{ of } \bullet\text{'s in } \tilde{T}} \cdot \text{wt}(\theta/\lambda).$$

Similarly,

$$(12.5) \quad \sum_{L \in \text{leaf}(\mathfrak{S}_{\tilde{T}^\dagger})} (-1)^{|\rho(L)/\lambda|+1} [\phi_1(\tilde{T})] \text{slide}_{\rho(L)/\lambda}(L) = (-1)^{1+\#\text{ of } \bullet\text{'s in } \tilde{T}^\dagger} \cdot \text{wt}(\theta/\lambda).$$

In particular, these quantities differ by a factor of -1 .

By inspection of the reverse miniswaps, revswap_{a_q} for $1 \leq a \leq i - 1$ does not affect any labels of family i or greater or any labels that are marked in T . Hence one sees that $\text{revswap}_{1_2} \circ \cdots \circ \text{revswap}_{(i-1)_{\mu_{i-1}}} \circ \text{revswap}_{i_1}(T)$ (respectively T^\dagger) is the same as $\text{revswap}_{1_2} \circ \cdots \circ \text{revswap}_{(i-1)_{\mu_{i-1}}} \circ \text{revswap}_{i_1}(\tilde{T})$ (respectively \tilde{T}^\dagger) followed by adding back the labels of $T \setminus \tilde{T}$ (respectively $T^\dagger \setminus \tilde{T}^\dagger$). Therefore, by our comparison of (12.4) and (12.5) above,

$$[T]P_{i_1} = (-1)^{1+\#\text{ of } \bullet\text{'s in } \tilde{T}} \cdot \text{wt}(\theta/\lambda) = -[T^\dagger]P_{i_1},$$

as desired.

Reduction to the $k = 1$ case: In the calculation of $\text{revswap}_{i_2} \circ \text{revswap}_{i_3} \circ \cdots \circ \text{revswap}_{i_k}(T)$ and $\text{revswap}_{i_2} \circ \text{revswap}_{i_3} \circ \cdots \circ \text{revswap}_{i_k}(T^\dagger)$, it is straightforward by inspection that each

reverse miniswap involving either \bullet of \mathcal{C}_1 or the \bullet of \mathcal{C}'_1 is L1.2. Therefore there exists an instance of \mathcal{C}_1 in each $W \in \mathfrak{R}_{i_1}(T)$ and an instance of \mathcal{C}'_1 in each $W' \in \mathfrak{R}_{i_1}(T^\dagger)$. By Lemma 12.6, these instances are unique. Extending ϕ_1 linearly, since T and T^\dagger are the same outside of the regions \mathcal{C}_1 and \mathcal{C}'_1 , it is easy to see inductively that for all $2 \leq q \leq k$,

$$\phi_1(\text{revswap}_{i_q} \circ \cdots \circ \text{revswap}_{i_k}(T)) = \text{revswap}_{i_q} \circ \cdots \circ \text{revswap}_{i_k}(T^\dagger).$$

In particular, ϕ_1 bijects $\mathfrak{R}_{i_1}(T)$ with $\mathfrak{R}_{i_1}(T^\dagger)$.

Let $V \in \mathfrak{R}_{i_1}(T)$. By the $k = 1$ case above, $[V]P_{i_1} = -[\phi_1(V)]P_{i_1}$. Moreover, when we apply $\text{swap}_{i_k^-} \circ \cdots \circ \text{swap}_{i_1}$ to V and $\phi_1(V)$, each miniswap involving a \bullet of \mathcal{C}_1 or \mathcal{C}'_1 is H3. Hence, $[T] \text{swap}_{i_k^-} \circ \cdots \circ \text{swap}_{i_1}(V) = [T^\dagger] \text{swap}_{i_k^-} \circ \cdots \circ \text{swap}_{i_1}(\phi_1(V))$. Thus by Lemma 12.5, $[T]P_{i_k} = -[T^\dagger]P_{i_k}$. \square

Let \mathcal{S}_2 be the subset of tableaux in $G_{\lambda,\mu}^\nu(i_k)$ with a box x such that $\bullet_{i_k} \in x$, $\bullet_{i_k} \in x^{\rightarrow\uparrow}$,

$i_{k+1} \in x^{\rightarrow}$ and $i_k \in \underline{x}$, i.e. locally the tableau is $\mathcal{C}_2 = \begin{array}{|c|c|} \hline & \bullet_{i_k} \\ \hline \bullet_{i_k} & i_{k+1} \\ \hline \end{array}$ (with possibly additional edge labels). Let \mathcal{S}'_2 be the subset of tableaux in $G_{\lambda,\mu}^\nu(i_k)$ with a box x such that $i_k \in x$, $i_{k+1} \in x^{\rightarrow}$, $\bullet_{i_k} \in x^{\rightarrow\uparrow}$, no i_k appears outside of x and no \bullet_{i_k} appears West of x in x 's row. Locally the

tableau is $\mathcal{C}'_2 = \begin{array}{|c|c|} \hline & \bullet_{i_k} \\ \hline i_k & i_{k+1} \\ \hline \end{array}$ (with possibly additional edge labels).

Lemma 12.10. *If $T \in \mathcal{S}_2$ (respectively, \mathcal{S}'_2), there is a unique \mathcal{C}_2 (respectively, \mathcal{C}'_2) that it contains.*

Proof. Let x be the southwestmost box of a \mathcal{C}_2 in T . By (G.7), the $i_k \in \underline{x}$ is the westmost i_k in T ; hence this configuration is unique. The claim about \mathcal{C}'_2 is clear since the i_k is unique. \square

For $T \in \mathcal{S}_2$, let $\phi_2(T)$ be T with the unique \mathcal{C}_2 replaced by \mathcal{C}'_2 .

Lemma 12.11. *$\phi_2 : \mathcal{S}_2 \rightarrow \mathcal{S}'_2$ is a bijection.*

Proof. This may be proved almost exactly as Lemma 12.7. \square

Proposition 12.12. *For each $T \in \mathcal{S}_2$, $[T]P_{i_k} = -[\phi_2(T)]P_{i_k}$.*

Proof. Let $T^\dagger := \phi_2(T)$. Let x be the southwestmost box of \mathcal{C}_2 in T . Then x is also the southwestmost box of \mathcal{C}'_1 in T^\dagger .

Special case $k = 1$: The proof is *verbatim* the argument for the $k = 1$ case of Proposition 12.8.

Reduction to the $k = 1$ case: Suppose $k > 1$. Let \mathcal{Z} be the set of boxes in an i_k -good tableau that either (1) contain \bullet_{i_k} or (2) contain a label \mathcal{F} with $i_1 \preceq \mathcal{F} \preceq i_{k-1}$ and are not southeast of a \bullet_{i_k} . Call an edge connected component of \mathcal{Z} an i_k -**walkway**. We will now apply the development of i -walkways, from Sections 10 and 11, in slightly modified form to the i_k -**walkways**. To be more precise, Lemmas 10.1, 10.2, 10.3 and 10.4 are true after replacing “ $(i+1)_1$ ” with “ i_k ” and “ i -walkway” with “ i_k -walkway”. In addition, Claim 11.1 holds *verbatim*. The proofs are trivial modifications of those given.

Let W be the i_k -walkway of T containing x (W includes all edges of boxes in W). Let W^\dagger be the analogous i_k -walkway of T^\dagger . Note that W and W^\dagger have the same skew shape.

Claim 12.13. *Let \mathbb{S}, \mathbb{S}' and \mathbb{T} be respectively the set of reversals of W , W^\dagger and W^c (the complement of W) under $\text{revswap}_{i_2} \circ \cdots \circ \text{revswap}_{i_k}$. Then:*

- (I) $\mathfrak{R}_{i_1}(T) = \{V \in G_{\lambda, \mu}^\nu(i_1) : V|_W \in \mathbb{S}, V|_{W^c} \in \mathbb{T}\}$
 (II) $\mathfrak{R}_{i_1}(T^\dagger) = \{V' \in G_{\lambda, \mu}^\nu(i_1) : V'|_{W^\dagger} \in \mathbb{S}', V'|_{(W^\dagger)^c} \in \mathbb{T}\}$

Proof. We prove only (I), as the proof of (II) is similar (using $T|_{W^c} = T^\dagger|_{W^c}$). Fix $2 \leq h \leq k$ and let L be a ladder of $A \in \mathfrak{R}_{i_h}(T)$. L contains only \bullet_{i_h} and unmarked i_{h-1} . Each of the boxes x of L is in \mathcal{Z} : This is clear if $h = k$ and follows for smaller h by induction. Thus $L \subseteq \mathcal{Z}$. Therefore, since L is edge connected, it sits inside an edge connected component of \mathcal{Z} . Thus, since W is one such component, reverse swapping acts independently on W and W^c . \square

Case 1: (W (and hence W^\dagger) has a single row): By construction, x is the eastmost box of W and W^\dagger . By Lemma 10.3(II), for every $V \in \mathfrak{R}_{i_1}(T)$, $V|_W = R'$. By Lemma 10.3(III), for every $V' \in \mathfrak{R}_{i_1}(T^\dagger)$, $V'|_{W^\dagger} \in \{R, R'\}$ where this R' is the same as in the previous sentence.

Since R' is the unique reversal of W and is a reversal of W^\dagger , we have $\mathfrak{R}_{i_1}(T) \subseteq \mathfrak{R}_{i_1}(T^\dagger)$ by Claim 12.13. Let $\iota : \mathfrak{R}_{i_1}(T) \rightarrow \mathfrak{R}_{i_1}(T^\dagger)$ be the inclusion map. Let $f : \mathfrak{R}_{i_1}(T) \rightarrow \mathfrak{R}_{i_1}(T^\dagger)$ be the map given by replacing the R' occupying the region W with R . Again appealing to Claim 12.13 we see that these maps are well-defined, injective and $\mathfrak{R}_{i_1}(T^\dagger) = \text{im } \iota \sqcup \text{im } f$.

By Claim 11.1(III), forward swapping R produces W^\dagger with coefficient 1. By Claim 11.1 (part (I.i) or (I.ii), as appropriate) forward swapping R' produces $\beta W + (1 - \beta)W^\dagger$ for some β . Moreover, when applying $\text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}$ to $V \in \mathfrak{R}_{i_1}(T)$ or $V' \in \mathfrak{R}_{i_1}(T^\dagger)$, every snake lies entirely inside some edge-connected component of \mathcal{Z} . W is one of these components. Thus, for each $V \in \mathfrak{R}_{i_1}(T)$, $[T] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(V)$ factors as a contribution from the region W times a contribution from $\mathcal{Z} \setminus W$. That is, for the same α ,

$$\sum_{V \in \mathfrak{R}_{i_1}(T)} [T] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(V) = \alpha\beta, \quad \sum_{V' \in \mathfrak{R}_{i_1}(T^\dagger)} [T^\dagger] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(V') = \alpha$$

$$\sum_{V' \in \mathfrak{R}_{i_1}(T^\dagger)} [T^\dagger] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(V') = \alpha(1 - \beta).$$

Therefore,

$$[T]P_{i_k} = \sum_{V \in \mathfrak{R}_{i_1}(T)} [V]P_{i_1} \cdot [T] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(V) = [V]P_{i_1} \alpha\beta,$$

while

$$\begin{aligned} [T^\dagger]P_{i_k} &= \sum_{V' \in \mathfrak{R}_{i_1}(T^\dagger)} [V']P_{i_1} \cdot [T^\dagger] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(V') \\ &= \sum_{V \in \mathfrak{R}_{i_1}(T)} [\iota(V)]P_{i_1} \cdot [T^\dagger] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(\iota(V)) \\ &+ \sum_{V \in \mathfrak{R}_{i_1}(T)} [f(V)]P_{i_1} \cdot [T^\dagger] \text{swap}_{i_{k-1}} \circ \dots \circ \text{swap}_{i_1}(f(V)) \end{aligned}$$

$$\begin{aligned}
&= \sum_{V \in \mathfrak{R}_{i_1}(T)} [V]P_{i_1} \cdot [T^\dagger] \text{swap}_{i_{k-1}} \circ \cdots \circ \text{swap}_{i_1}(V) \\
&- \sum_{V \in \mathfrak{R}_{i_1}(T)} [V]P_{i_1} \text{swap}_{i_{k-1}} \circ \cdots \circ \text{swap}_{i_1}(f(V)) \\
&= [V]P_{i_k}(\alpha(1 - \beta) - \alpha).
\end{aligned}$$

Now, $[T]P_{i_k} = -[T^\dagger]P_{i_k}$ follows.

Case 2: (W (and hence W^\dagger) has at least two rows): There are three cases to consider, corresponding to the case of Lemma 10.4.

In Cases (I) and (II) of Lemma 10.4, W and W^\dagger have a unique reversal R . By Claim 11.1(III) or Claim 11.1(II) respectively, forward swapping R produces $\beta W - \beta W^\dagger$ for some β . In Case (III) of Lemma 10.4, W and W^\dagger share the same pair of reversals R, R' . By Claim 11.1(III) and (I.i), forward swapping R produces $\beta W - \beta W^\dagger$ for some β , while forward swapping R' produces $\beta' W - \beta' W^\dagger$ for some β' . Using these facts, one may argue similarly to Case 1 to deduce $[T]P_{i_k} = -[T^\dagger]P_{i_k}$. \square

Let \mathcal{S}_3 be the subset of tableaux in $G_{\lambda, \mu}^\nu(i_k)$ with a box x such that $\bullet_{i_k} \in x$, $i_k \in x^\rightarrow$ and $i_k \in \underline{x}$, i.e. locally the tableau is $\mathcal{C}_3 = \begin{array}{|c|c|} \bullet_{i_k} & i_k \\ \hline & \end{array}$ (with possibly additional edge labels). Let \mathcal{S}'_3 be the subset of tableaux in $G_{\lambda, \mu}^\nu(i_k)$ with a box x such that $\bullet_{i_k} \in x$, $i_k \in x^\rightarrow$, no i_k appears West of x^\rightarrow , $i_{k-1} \notin x^\leftarrow$, and $(i+1)_h \notin \underline{x^\rightarrow}$ where $N_{i_k} = N_{(i+1)_h}$. Locally the tableau is $\mathcal{C}'_3 = \begin{array}{|c|c|} \bullet_{i_k} & i_k \\ \hline & \end{array}$ (with possibly additional edge labels).

Lemma 12.14. *If $T \in \mathcal{S}_3$ (respectively, \mathcal{S}'_3), there is a unique \mathcal{C}_3 (respectively, \mathcal{C}'_3) that it contains.*

Proof. If \mathcal{C}_3 occurs in a good tableau, it is unique since the edge i_k is westmost in its gene by (G.7). Similarly \mathcal{C}'_3 is unique since the $i_k \in x^\rightarrow$ is westmost by assumption. \square

Define $\phi_3(T)$ to be T with the unique \mathcal{C}_3 replaced by \mathcal{C}'_3 .

Lemma 12.15. $\phi_3 : \mathcal{S}_3 \rightarrow \mathcal{S}'_3$ is a bijection.

Proof. Define the (putative) inverse ϕ_3^{-1} by replacing \mathcal{C}'_3 with \mathcal{C}_3 . Once we establish that ϕ_3 and ϕ_3^{-1} are well-defined, we are done, since ϕ_3 and ϕ_3^{-1} are clearly mutually inverse.

Let $T \in \mathcal{S}_3$. Trivially, each (G.n) holds for $\phi_3(T)$. By T 's (G.12), $i_{k-1} \notin x^\leftarrow$. If $(i+1)_h \in \underline{x^\rightarrow}$ in $\phi_3(T)$ with $N_{i_k} = N_{(i+1)_h}$, then T would violate Lemma 5.6. By T 's (G.4) and (G.7), the $i_k \in x^\rightarrow$ is westmost in $\phi_3(T)$.

Now let $T \in \mathcal{S}'_3$. We check the goodness conditions for $\phi_3^{-1}(T)$.

Claim 12.16. *No label of family i appears in x 's column in T .*

Proof. By T 's (G.12), there are no labels of family i North of x and in its column. By T 's (G.11), a label ℓ South of x and in its column is not marked, i.e., $\ell \succeq i_k$. Since we assumed the $i_k \in x^\rightarrow$ is westmost, $\ell \neq i_k$. By T 's (G.6), $\ell \neq i_l$ for $l > k$. Hence $i_k < \ell$. \square

(G.4) and (G.5): By T 's (G.9), all labels North of x and in its column are of family at most i . By T 's (G.11), all labels South of x and in its column are of family at least i . Hence by Claim 12.16, $\phi_3^{-1}(T)$'s (G.4) and (G.5) follow.

(G.8): If there is a genotype G of $\phi_3^{-1}(T)$ that is not ballot, then it uses the $i_k \in \underline{x}$. Furthermore, since T is ballot, some $(i+1)_h$ with $N_{i_k} = N_{(i+1)_h}$ appears in $\text{word}(G)$ before the $i_k \in \underline{x}$. By Lemma 5.6 applied to T , this $(i+1)_h$ can only be South of x^\rightarrow and in x^\rightarrow 's column or North of x and in x 's column. By T 's (G.9), it cannot be North of x and in its column. Suppose it appears South of x^\rightarrow and in its column. By assumption, $(i+1)_h \notin \underline{x^\rightarrow}$. Hence suppose it appears south of x^\rightarrow , and consider $\text{label}(x^\downarrow)$. By (G.11) $\text{family}(\text{label}(x^\downarrow)) \geq i$. By Claim 12.16, $\text{family}(\text{label}(x^\downarrow)) \neq i$. By T 's (G.3) and (G.4), $\text{label}(x^\downarrow) \prec (i+1)_h$. Hence $\text{family}(\text{label}(x^\downarrow)) = i+1$. But by Lemma 5.10, $\text{label}(x^\downarrow) \neq (i+1)_h$. Hence by T 's (G.6), $(i+1)_{h-1} \in x^\downarrow$. This creates a (G.8) violation in T , as this label is read before any i_{k-1} .

(G.12): Since T is good, if $\phi_3^{-1}(T)$ violates (G.12), the violation involves the $i_k \in \underline{x}$. Since by assumption $i_{k-1} \notin x^\leftarrow$, the last sentence of (G.12) does not apply. Suppose i_j is SouthEast of \underline{x} , then it is also SouthEast of $i_k \in x^\rightarrow$, which will lead to a violation of T 's (G.12). Suppose i_j is NorthWest of \underline{x} , then to avoid a violation of T 's (G.12) with the $i_k \in x^\rightarrow$, i_j must be either in x 's row or in an upper edge of that row. Since we have $\bullet_{i_k} \in x$, this avoids violating $\phi_3^{-1}(T)$'s (G.12).

All of the remaining (G. n)-conditions are trivial to verify. \square

Proposition 12.17. For $T \in \mathcal{S}_3$, $[T]P_{i_k} = [\phi_3(T)]P_{i_k}$.

Proof. Let $T^\dagger := \phi_3(T)$. By inspection of the reverse miniswaps, and downward induction on \mathcal{Q} , there is a bijection $f_{\mathcal{Q}} : \mathfrak{R}_{\mathcal{Q}}(T) \rightarrow \mathfrak{R}_{\mathcal{Q}}(T^\dagger)$ given by deletion of the $i_k \in \underline{x}$. If $L \in \mathfrak{R}_{1_1}(T)$, then L and $f_{1_1}(L)$ have the same number of \bullet_{1_1} 's. Hence, $[L]P_{1_1} = [f_{1_1}(L)]P_{1_1}$; cf. (11.4).

Extend $f_{\mathcal{Q}}$ linearly. By inspection of the miniswaps,

$$f_{i_k}(\text{swap}_{i_k^-} \circ \cdots \circ \text{swap}_{1_1}(L)) = \text{swap}_{i_k^-} \circ \cdots \circ \text{swap}_{1_1}(f_{1_1}(L)).$$

Hence by Lemma 12.5, $[T]P_{i_k} = [T^\dagger]P_{i_k}$. \square

Let \mathcal{S}_4 be the subset of tableaux in $G_{\lambda, \mu}^\nu(i_k)$ with a box x such that $\bullet_{i_k} \in x$, $\mathcal{F}^1 \in x^\rightarrow$, $i_k \in \underline{x}$ and $(i_k) \in \underline{x^\rightarrow}$, i.e. locally the tableau is $\mathcal{C}_4 = \begin{array}{|c|c|} \bullet_{i_k} & \mathcal{F}^1 \\ \hline i_k & (i_k) \end{array}$ (with possibly additional edge labels).

Let \mathcal{S}'_4 be the subset of tableaux in $G_{\lambda, \mu}^\nu(i_k)$ with a box x such that $\bullet_{i_k} \in x$, $\mathcal{F}^1 \in x^\rightarrow$, $i_k \in \underline{x^\rightarrow}$,

$(i+1)_h \notin \underline{x^\rightarrow}$ if $N_{(i+1)_h} = N_{i_k}$, and $i_{k-1} \notin x^\leftarrow$. Locally the tableau is $\mathcal{C}'_4 = \begin{array}{|c|c|} \bullet_{i_k} & \mathcal{F}^1 \\ \hline i_k & \end{array}$ (with possibly additional edge labels).

Lemma 12.18. If $T \in \mathcal{S}_4$ (respectively, \mathcal{S}'_4), there is a unique \mathcal{C}_4 (respectively, \mathcal{C}'_4) that it contains.

Proof. This follows since by (G.7), T contains at most one edge label i_k . \square

Set $\phi_4 : \mathcal{S}_4 \rightarrow \mathcal{S}'_4$ by replacing \mathcal{C}_4 with \mathcal{C}'_4 .

Lemma 12.19. $\phi_4 : \mathcal{S}_4 \rightarrow \mathcal{S}'_4$ is a bijection.

Proof. Define a putative inverse $\phi_4^{-1} : \mathcal{S}'_4 \rightarrow \mathcal{S}_4$ by replacing \mathcal{C}'_4 with \mathcal{C}_4 . Clearly, ϕ_4 and ϕ_4^{-1} are mutually inverse. It remains to check well-definedness. Indeed, it is trivial to check each goodness condition holds for $\phi_4(T)$. By Lemma 5.6 for T , there is not $(i+1)_h \in \underline{x^\rightarrow}$ with $N_{(i+1)_h} = N_{i_k}$. By T 's (G.12), $i_{k-1} \notin x^\leftarrow$. Thus ϕ_4 is well-defined.

Claim 12.20. No label of family i appears in x 's column in T .

Proof. By T 's (G.12), i_ℓ cannot appear North of x and in its column. If i_ℓ is South of x and in its column, then by T 's (G.6) and (G.7), $\ell < k$, so this i_ℓ is marked, contradicting T 's (G.11). \square

Now let $T \in \mathcal{S}'_4$. We check the goodness conditions for $\phi_4^{-1}(T)$:

(G.4) and (G.5): By T 's (G.9), every label North of x and in its column has family at most i . By T 's (G.11), every label South of x and in its column has family at least i . Moreover, by Claim 12.20, no label of family i appears in x 's column in T . Hence (G.4) and (G.5) hold in $\phi_4^{-1}(T)$.

(G.8): Suppose $\phi_4^{-1}(T)$ has a nonballot genotype G . By T 's (G.8), G must use the $i_k \in \underline{x}$. Also by T 's (G.8), some $(i+1)_h$ with $N_{(i+1)_h} = N_{i_k}$ appears in $\text{word}(G)$ before this $i_k \in \underline{x}$. By T 's (G.9) and (G.8), this $(i+1)_h$ appears South of x^\rightarrow and in x^\rightarrow 's column. By T 's (G.4) and the first hypothesis on \mathcal{S}'_4 , in fact $(i+1)_h \in x^{\rightarrow\downarrow}$. By T 's (G.3), $\text{family}(\text{label}(x^\downarrow)) \leq i+1$. By (G.11) and the $\bullet_{i_k} \in x$, $\text{family}(\text{label}(x^\downarrow)) \geq i$. By Claim 12.20, no label of family i appears in x 's column in T . Thus $\text{family}(\text{label}(x^\downarrow)) = i+1$. Then by T 's (G.3) and (G.6), $(i+1)_{h-1} \in x^\downarrow$. Hence by Claim 12.20, this contradicts Lemma 5.6 for T .

(G.12): If there is an i_ℓ SouthEast of the $i_k \in \underline{x}$ in $\phi_4^{-1}(T)$, then we either violate T 's (G.2), (G.4) or (G.12). Now suppose there is an i_ℓ NorthWest of $i_k \in \underline{x}$ in $\phi_4^{-1}(T)$. By T 's (G.12), this i_ℓ is West and either in x 's row or on the upper edge of that row. If $i_\ell \in x^\leftarrow$, then $\ell = k-1$ by T 's (G.6). However then we contradict the last hypothesis on \mathcal{S}'_4 . So the i_ℓ and i_k satisfy (G.12).

The remaining goodness conditions are trivial to verify. \square

Proposition 12.21. *For each $T \in \mathcal{S}_4$, $[T]P_{i_k} = [\phi_4(T)]P_{i_k}$.*

Proof. Let $T^\dagger = \phi_4(T)$. Let $f_Q : \mathfrak{R}_Q(T) \rightarrow \mathfrak{R}_Q(T^\dagger)$ be defined by deleting the $i_k \in \underline{x}$ and replacing the $(i_k) \in \underline{x}^\rightarrow$ by i_k . Now the proof proceeds exactly as that for Proposition 12.17. \square

13. PROOF OF THE CONJECTURAL K_T RULE FROM [ThYo13]

We briefly recap the conjectural rule for $K_{\lambda, \mu}^\nu$ from [ThYo13, Section 8]. An **equivariant increasing tableau** is an edge-labeled filling of ν/λ using the labels $1, 2, \dots, |\mu|$ such that each label is strictly smaller than any label below it in its column and each *box* label is strictly smaller than the box label immediately to its right. Any subset of the boxes of ν/λ may be marked by \star 's, except that if i and $i+1$ are box labels in the same row, then the box containing i may *not* be \star -ed. Let $\text{EqInc}(\nu/\lambda, |\mu|)$ denote the set of all such equivariant increasing tableaux.

An **alternating ribbon** R is a filling of a short ribbon by two symbols such that adjacent boxes are filled differently; all edges except the southwestmost edge are empty; and if this edge is filled, it is filled with the other symbol than in the box above it. Let $\text{switch}(R)$ be the alternating ribbon of the same shape where each box is instead filled with the other symbol. If the southwestmost edge was filled by one of these symbols, that symbol is deleted. If R consists of a single box with only one symbol used, then switch does nothing to it. Define switch to act on an edge-disjoint union of alternating ribbons, by acting on each independently.

Example 13.1. Let $R = \begin{array}{|c|c|c|} \hline & \heartsuit & \spadesuit \\ \hline \heartsuit & \spadesuit & \\ \hline \spadesuit & & \\ \hline \end{array}$. Then $\text{switch}(R) = \begin{array}{|c|c|c|} \hline & \spadesuit & \heartsuit \\ \hline \spadesuit & \heartsuit & \\ \hline & & \\ \hline \end{array}$. □

Given $T \in \text{EqInc}(\nu/\lambda, |\mu|)$ and an inner corner $x \in \lambda$, label x with \bullet and erase all \star 's. Call this tableau V_1 . Consider the alternating ribbons $\{R_1\}$ made of \bullet and 1. V_2 is obtained by applying switch to each R_1 . Now let $\{R_2\}$ be the collection of ribbons consisting of \bullet and 2, and produce V_3 by applying switch to each R_2 . Repeat until the \bullet 's have been switched past all the numerical labels in T ; the final placement of numerical labels gives $\text{KEqjdt}_x(T)$, the **slide** of T into x . The sequence V_1, V_2, \dots is the **switch sequence** of (T, x) . Finally, define $\text{KEqrect}(T)$ by successively applying KEqjdt_x in **column rectification order**, i.e., successively pick x to be the eastmost inner corner.

Lemma 13.2. For V_j in the switch sequence of (T, x) :

- (I) The numerical box labels strictly increase along rows from left to right (ignoring \bullet 's).
- (II) The numerical labels strictly increase down columns (ignoring \bullet 's and reading labels of a given edge in increasing order).
- (III) Every numerical label southeast of a \bullet is at least j .
- (IV) Every numerical label northwest of a \bullet is strictly less than j .

Proof. These are proved by simultaneous induction on j . In the inductive step, one considers any 2×2 local piece of V_j and studies the possible cases that can arise as one transitions from $V_j \rightarrow V_{j+1}$; we leave the straightforward details to the reader. □

A set of labels is a **horizontal strip** if they are arranged in increasing order from southwest to northeast, with no two labels of the set in the same column.

Lemma 13.3. Let $T \in \text{EqInc}(\nu/\lambda, |\mu|)$ and $x \in \lambda$ be an inner corner. Then $\{i, i+1, \dots, j\}$ forms a horizontal strip in V_k of the switch sequence of (T, x) if and only if it does so in V_{k+1} .

Proof. This quickly reduces to consideration of the possibilities in a 2×2 local piece of V_k . Then we proceed by straightforward case analysis using Lemma 13.2. □

A label $s \in T$ is **special** if it is an edge label or lies in a \star -ed box. At most one s appears in a column c . In column rectification order, each slide KEqjdt_x for $x \in c$ moves an s in c at most one step North (and it remains in c). A special label s in c **passes through** x if it occupies x at any point during c 's rectification *and* initially $s \notin x$. Let x_1, \dots, x_s be the boxes s passes through and let y_1, \dots, y_t be the numerically labeled boxes East of x_s in the same row. Set $\text{factor}_K(s) := 1 - \prod_{i=1}^s \hat{\beta}(x_i) \prod_{j=1}^t \hat{\beta}(y_j)$. If s does not move during the rectification of c , then $\text{factor}_K(s) := 0$. Now set $\text{wt}_K(T) := \prod_s \text{factor}_K(s)$, where the product is over all special labels. Lastly, we define $\text{sgn}(T) := (-1)^{|\mu| - \#\star\text{'s in } T - \#\text{labels in } T}$.

Let $\mu[1] = (1, 2, 3, \dots, \mu_1)$, $\mu[2] = (\mu_1 + 1, \mu_1 + 2, \dots, \mu_1 + \mu_2)$, etc. Let T_μ be the **super-standard tableau** of shape μ , i.e., row i is filled by $\mu[i]$. The following is the conjecture of [ThYo13]:

Theorem 13.4. $K_{\lambda, \mu}^\nu = \sum_T \text{sgn}(T) \cdot \text{wt}_K(T)$, where the sum is over

$$\mathcal{A}_{\lambda, \mu}^\nu := \{T \in \text{EqInc}(\nu/\lambda, |\mu|) : \text{KEqrect}(T) = T_\mu\}.$$

We will prove Theorem 13.4 (after some preparation) by connecting to Theorem 1.3.

Let $\mathcal{B}_{\lambda,\mu}^\nu$ be the set of all $T \in \text{BallotGen}(\nu/\lambda)$ that have content μ . We need a **semistandardization map** $\Phi: \mathcal{A}_{\lambda,\mu}^\nu \rightarrow \mathcal{B}_{\lambda,\mu}^\nu$. Given $A \in \mathcal{A}_{\lambda,\mu}^\nu$, erase all \star 's and replace the labels $1, 2, \dots, \mu_1$ with $1_1, 1_2, \dots, 1_{\mu_1}$ respectively. Next, replace $\mu_1 + 1, \mu_1 + 2, \dots, \mu_1 + \mu_2$ by $2_1, 2_2, \dots, 2_{\mu_2}$ respectively, etc. The result is $\Phi(A)$. Note Φ is not bijective. Define a **standardization map** $\Psi: \mathcal{B}_{\lambda,\mu}^\nu \rightarrow \mathcal{A}_{\lambda,\mu}^\nu$ by reversing the above process in the obvious way; $\Psi(B)$ is \star -less.

Lemma 13.5. *For $B \in \mathcal{B}_{\lambda,\mu}^\nu$, $\Psi(B) \in \text{EqInc}(\nu/\lambda, |\mu|)$.*

Proof. That $\Psi(B)$ has the desired shape and content is clear. Row strictness follows from (S.1), and column strictness from (S.2). \square

Lemma 13.6. *For $B \in \mathcal{B}_{\lambda,\mu}^\nu$ and for each i , $\mu[i]$ forms a horizontal strip in $\Psi(B)$ and also in each tableau of any switch sequence during the column rectification of $\Psi(B)$.*

Proof. By (S.2–4), the labels i_1, \dots, i_{μ_i} form a horizontal strip of B . The claim for $\Psi(B)$ is then immediate by definition of Ψ . The claim for the tableaux of the switch sequences then follows by Lemma 13.3. \square

Lemma 13.7. *Let $B \in \mathcal{B}_{\lambda,\mu}^\nu$. Then*

- (I) *after column rectifying the eastmost j columns of $\Psi(B)$, there are no edge labels in these eastmost j columns; and*
- (II) *while rectifying the next column, there is never an edge label north of a \bullet and in the same column, in any tableau of any switch sequence.*

Proof. (I): Suppose there were such an edge label $\ell \in \underline{x}$ after rectifying the eastmost j columns. Then $\ell \in \underline{x}$ in $\Psi(B)$, since rectification never adds a label to any edge. Suppose x is in the i th row from the top of $\Psi(B)$. Then since no label of B is too high, $\ell \in \mu[k]$ where $k \leq i$. Let the boxes North of \underline{x} and in the same column be $x_1, \dots, x_i = x$ from north to south. By Lemma 13.2(II), we have for each e that $\text{label}(x_e) \in \mu[f(e)]$ for some $f(e) \leq k$. But then by Lemma 13.6, $f: \{1, 2, \dots, i\} \rightarrow \{1, 2, \dots, k-1\}$ is injective, a contradiction.

(II): Let c be the column currently being rectified. For the columns East of c , the claim follows from part (I), noting that rectification never adds a label to any edge. For column c itself, the claim is vacuous if there is no \bullet in c . If there is $\bullet \in c$, the claim follows from noting that every label of column c North of this \bullet must have participated in some switch and that switch never outputs any edge labels. \square

An equivariant increasing tableau T is **ballot** if $\Phi(T)$ is ballot in the sense of Section 1.3. That is, for every \tilde{T} obtained by selecting one copy of each label in T , every initial segment of \tilde{T} 's column reading word has, for each $i \geq 1$, at least as many labels from $\mu[i]$ as from $\mu[i+1]$. We extend this definition to tableaux with \bullet 's by ignoring the \bullet 's.

Lemma 13.8. *Let $B \in \mathcal{B}_{\lambda,\mu}^\nu$. Then $\Psi(B)$ is ballot, as is each tableau of any switch sequence during the column rectification of $\Psi(B)$.*

Proof. Let $A = \Psi(B)$. Since B is ballot and $\Phi(A) = B$, A is ballot by definition. Suppose that some V_q is ballot, but V_{q+1} is not. Then there exist i and a \widetilde{V}_{q+1} with a ballotness violation between $\mu[i]$ and $\mu[i+1]$.

If $q \notin \mu[i] \cup \mu[i+1]$, then the labels of $\mu[i]$ and $\mu[i+1]$ appear in the same locations in V_q and V_{q+1} , contradicting that V_q is ballot.

If $q \in \mu[i+1]$, then no $\mu[i]$ -label moves. For each $\ell \in \mu[i+1]$ appearing in \widetilde{V}_{q+1} , there is an ℓ east of that position in V_q . Hence we construct a nonballot \widetilde{V}_q by choosing those corresponding ℓ 's, the same labels from $\mu[i]$ as in \widetilde{V}_{q+1} , and all other labels arbitrarily. This contradicts that V_q is ballot.

Finally if $q \in \mu[i]$, then there is some x in column c of V_q with $\bullet \in x$ and $q \in x^{\rightarrow}$ such that the q moving into x violates ballotness in the columns East of c . That is, locally the switch is

$$V_q \supseteq \begin{array}{|c|c|} \hline a & b \\ \hline \bullet & q \\ \hline d & e \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & b \\ \hline q & \bullet \\ \hline d & e \\ \hline \end{array} \subseteq V_{q+1} \quad \text{or} \quad V_q \supseteq \begin{array}{|c|c|} \hline a & \bullet \\ \hline \bullet & q \\ \hline d & e \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & q \\ \hline q & \bullet \\ \hline d & e \\ \hline \end{array} \subseteq V_{q+1},$$

where the x is the left box of the second row. The $q \in x^{\rightarrow}$ is Westmost in V_q , since otherwise the nonballotness of V_{q+1} contradicts that V_q is ballot. In particular, $q \neq d$. Hence by Lemma 13.2(III), $q < d$.

Since V_q is ballot but V_{q+1} is not, there is a $\bar{q} \in \mu[i+1]$ in c^{\rightarrow} in V_q , and hence in V_{q+1} . By Lemma 13.2(II) applied to V_q , this \bar{q} is below q in c^{\rightarrow} . By Lemma 13.7(I), there are no edge labels East of column c . So in fact e and hence d both exist. Indeed by Lemma 13.2(II) and Lemma 13.6, $e = \bar{q}$. By Lemma 13.6, q is the only label of $\mu[i]$ that appears in c in V_{q+1} . Hence $d \notin \mu[i]$. Thus by Lemma 13.2(I) applied to V_q , we conclude $d \in \mu[i+1]$. However this again contradicts that V_q is ballot. \square

For $A \in \text{EqInc}(\nu/\lambda, |\mu|)$, let $A^{(k)}$ be the ‘‘partial’’ tableau that is the column rectification of the eastmost k columns of A .

Lemma 13.9. *Let $B \in \mathcal{B}_{\lambda, \mu}^{\nu}$ and let $A = \Psi(B)$. For each i , the i th row of $A^{(k)}$ consists of a (possibly empty) final segment from $\mu[i]$.*

Proof. By Lemma 13.2(I, II), $A^{(k)}$ has strictly increasing rows and columns. By Lemma 13.6, the labels $\mu[i]$ form a horizontal strip in $A^{(k)}$ for each i ; moreover the labels of $\mu[i]$ appearing in $A^{(k)}$ are a final segment of $\mu[i]$. By Lemma 13.7(I), there are no edge labels in $A^{(k)}$. By Lemma 13.8, $A^{(k)}$ is ballot. The lemma follows. \square

Corollary 13.10. *A rectifies to T_{μ} .*

Proof. Immediate from Lemma 13.9. \square

Proposition 13.11. *For $B \in \mathcal{B}_{\lambda, \mu}^{\nu}$, $\Psi(B) \in \mathcal{A}_{\lambda, \mu}^{\nu}$.*

Proof. By Lemma 13.5, $\Psi(B) \in \text{EqInc}(\nu/\lambda, |\mu|)$. By Corollary 13.10, $\Psi(B)$ rectifies to T_{μ} . \square

Lemma 13.12. *For $A \in \mathcal{A}_{\lambda, \mu}^{\nu}$, $\mu[i]$ forms a horizontal strip in A and each A' in the column rectification of T .*

Proof. This is true for T_{μ} , and hence true for A and each A' by Lemma 13.3. \square

Lemma 13.13. *For $A \in \mathcal{A}_{\lambda, \mu}^{\nu}$, $\Phi(A)$ is semistandard.*

Proof. Row-strictness of A implies that $\Phi(A)$ satisfies (S.1). Since by Lemma 13.12, $\mu[i]$ is a horizontal strip in A for each i , (S.2)–(S.4) hold in $\Phi(A)$. \square

Lemma 13.14. *For $A \in \mathcal{A}_{\lambda, \mu}^\nu$, $\Phi(A)$ is ballot.*

Proof. Suppose $\Phi(A)$ is not ballot. Then by definition, A is not ballot. We assert that every tableau in every switch sequence in the column rectification of A is also *not* ballot, implying T_μ is not ballot, a contradiction.

Suppose V_ℓ is not ballot, but $V_{\ell+1}$ is. We derive a contradiction. Since V_ℓ is not ballot, we pick a nonballot \tilde{V}_ℓ . Suppose this nonballotness can be blamed on positions a_1, \dots, a_s containing labels of $\mu[i]$ and positions b_1, \dots, b_{s+1} containing labels of $\mu[i+1]$ (for some i). Suppose a_1, \dots, a_s and b_1, \dots, b_{s+1} are left to right in \tilde{V}_ℓ ; no two a_j 's (respectively b_j 's) are in the same column by Lemma 13.12. We may assume b_1 is southwestmost among all these positions, say in column c and that among all offending choices of i and positions, we picked one so that c is eastmost.

Since $V_{\ell+1}$ is supposed ballot, there is a label $\ell \in \mu[i+1]$ in b_1 of V_ℓ that moved to column c^\leftarrow . Locally, the switch is $\begin{array}{|c|c|} \hline x & y \\ \hline \bullet & \ell \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline x & y \\ \hline \ell & \bullet \\ \hline \end{array}$. By Lemma 13.12, $\mu[i+1]$ forms a horizontal strip in V_ℓ . Hence $x, y \notin \mu[i+1]$. Also, no label in column c is in $\mu[i]$ since otherwise we contradict that c is chosen eastmost. Now, there is some label $m \in \mu[i]$ above the \bullet in column c^\leftarrow of V_ℓ since $V_{\ell+1}$ is ballot. Using Lemma 13.7(II), it follows that $m = x$. Now, we have argued $y \notin \mu[i] \cup \mu[i+1]$. However, by Lemma 13.2(I, II) applied to V_ℓ , there are no other possibilities for y , a contradiction. \square

Proposition 13.15. *For $A \in \mathcal{A}_{\lambda, \mu}^\nu$, $\Phi(A) \in \mathcal{B}_{\lambda, \mu}^\nu$.*

Proof. By construction, $\Phi(A)$ is an edge-labeled genomic tableau of shape ν/λ and content μ . By Lemma 13.13, $\Phi(A)$ is semistandard. By Lemma 13.14, $\Phi(A)$ is ballot. Since A rectifies to T_μ , no label of $\Phi(A)$ is too high. \square

Given a label ℓ in $A \in \mathcal{A}_{\lambda, \mu}^\nu$, let $\Phi(\ell)$ be the corresponding label in $\Phi(A) \in \mathcal{B}_{\lambda, \mu}^\nu$. Recall the definitions of Section 1.4.

Lemma 13.16.

- (I) *If ℓ is an edge label, then $\text{factor}_K(\ell) = \text{edgfactor}(\Phi(\ell))$.*
- (II) *If ℓ is in a \star -ed box, then $\text{factor}_K(\ell) = 1 - \text{boxfactor}(\Phi(\ell))$.*

Proof. These follow from the definitions of the factors combined with Lemma 13.9. \square

Lemma 13.17. *If $B \in \mathcal{B}_{\lambda, \mu}^\nu$, then*

$$\text{boxwt}(B) = \sum_{A \in \Phi^{-1}(B)} (-1)^{\#\star\text{'s in } A} \prod_{\text{special box label } \ell \text{ of } A} \text{factor}_K(\ell).$$

Proof. A box x is productive in B if and only if it may be \star -ed in $\Psi(B)$. We are done by Lemma 13.16(II) and the ‘‘inclusion-exclusion’’ identity $\sum_{S \subset [N]} (-1)^{|S|} \prod_{s \in S} (1 - z_s) = z_1 z_2 \cdots z_N$. \square

Proof of Theorem 13.4. Recall Theorem 13.4 asserts $K_{\lambda, \mu}^\nu = \sum_{A \in \mathcal{A}_{\lambda, \mu}^\nu} \text{sgn}(A) \text{wt}_K(A)$. To see this, observe that by Propositions 13.11 and 13.15,

$$\begin{aligned}
\sum_{A \in \mathcal{A}_{\lambda, \mu}^{\nu}} \text{sgn}(A) \text{wt}_K(A) &= \sum_{B \in \mathcal{B}_{\lambda, \mu}^{\nu}} \sum_{A \in \Phi^{-1}(B)} \text{sgn}(A) \text{wt}_K(A) \\
&= \sum_{B \in \mathcal{B}_{\lambda, \mu}^{\nu}} \sum_{A \in \Phi^{-1}(B)} (-1)^{|\mu| - \#\star\text{'s in } A - \#\text{labels in } A} \prod_{\text{edge label } \ell \text{ of } A} \text{factor}_K(\ell) \prod_{\text{special box label } \ell \text{ of } A} \text{factor}_K(\ell) \\
&= \sum_{B \in \mathcal{B}_{\lambda, \mu}^{\nu}} \sum_{A \in \Phi^{-1}(B)} (-1)^{|\mu| - \#\text{labels in } A} \left(\prod_{\text{edge label } \ell \text{ of } A} \text{factor}_K(\ell) \right) (-1)^{\#\star\text{'s in } A} \prod_{\text{special box label } \ell \text{ of } A} \text{factor}_K(\ell).
\end{aligned}$$

The number of labels of A equals the number of labels of B for any $A \in \Phi^{-1}(B)$. Combining this with Lemma 13.16(I) shows the previous expression equals

$$= \sum_{B \in \mathcal{B}_{\lambda, \mu}^{\nu}} (-1)^{|\mu| - \#\text{labels in } B} \left(\prod_{\text{edge label } \ell \text{ of } B} \text{edgefactor}(\ell) \right) \sum_{A \in \Phi^{-1}(B)} (-1)^{\#\star\text{'s in } A} \prod_{\text{special box label } \ell \text{ of } A} \text{factor}_K(\ell).$$

By Lemma 13.17, this equals

$$= \sum_{B \in \mathcal{B}_{\lambda, \mu}^{\nu}} (-1)^{|\mu| - \#\text{labels in } B} \text{edgewt}(B) \text{boxwt}(B) := L_{\lambda, \mu}^{\nu}.$$

Since by Theorem 1.3, $L_{\lambda, \mu}^{\nu} = K_{\lambda, \mu}^{\nu}$, we are done. \square

APPENDIX A. PROOF OF PROPOSITION 7.7

We check (G.1)–(G.13) are preserved. Let $U \in \text{swap}_{\mathcal{G}}(T)$. We prove the conditions in the order: (G.1), (G.2), (G.4), (G.5), (G.6), (G.7), (G.3), (G.8), (G.9), (G.11), (G.10), (G.12), (G.13). In this way, each proof depends only on previously proved conditions. It is also necessary to show that the prescribed virtual labels in H5.2 and T4.2 satisfy the rules (V.1)–(V.3) for virtual labels. This is done in Lemma A.7 as part of the discussion of (G.13).

(G.1): By T 's (G.1), no label of T is too high. Hence if some label \mathcal{P} of U is too high, it must be placed in such a location by some miniswap. We therefore consider all miniswaps that might place a label \mathcal{P} on x or \underline{x} in U , when there is no \mathcal{P} or $\textcircled{\mathcal{P}}$ north of \underline{x} in T . By inspection, the miniswaps that can do so are H1, T1, T3, T4 and T5.

H1: The first output of H1 is not problematic. For if the $\mathcal{G} \in x$ in the first output were too high, the $\mathcal{G} \in \underline{x}$ in T would also have been too high, in violation of T 's (G.1). If the second output of H1 creates a label that is too high, then by definition of γ , that output is produced with coefficient 0. Thus H1 does not create a tableau violating (G.1).

T1: Suppose T has $\bullet_{\mathcal{G}} \in x$ and that, after applying T1, U has $\mathcal{G} \in x$, which is too high. By Lemma 6.9(IV,VII), T has $\mathcal{G} \in x^{\downarrow}$. By T 's (G.1), this $\mathcal{G} \in x^{\downarrow}$ is not too high. Since $\mathcal{G} \in x^{\downarrow}$ is *not* too high, but $\mathcal{G} \in x$ is too high, x is in row $i - 1$ where $i = \text{family}(\mathcal{G})$. In particular, $i > 1$.

By T 's (G.2), $\bullet_{\mathcal{G}} \notin x^{\rightarrow}$ in T , so if x^{\rightarrow} is a box of T , then it contains a genetic label. Consider $\text{label}_T(x^{\rightarrow})$. If $\text{label}_T(x^{\rightarrow}) \prec \mathcal{G}$, then $\text{label}_T(x^{\rightarrow})$ is marked. Hence by Lemma 5.13, T has $\textcircled{\mathcal{G}} \in x^{\rightarrow}$. This contradicts that we are applying T1, for then x^{\rightarrow} is adjoined to the snake containing x by (R.3). If $\text{label}_T(x^{\rightarrow}) = \mathcal{G}$, this again contradicts that we are applying T1. If $\text{label}_T(x^{\rightarrow}) \succ \mathcal{G}$, then $\text{family}(\mathcal{G}) \leq \text{family}(\text{label}_T(x^{\rightarrow}))$. Thus if $\mathcal{G} \in x$ were too high

in U , then $\text{label}_T(x^\rightarrow)$ would already be too high in T , violating T 's (G.1). Thus x^\rightarrow must not be a box of T .

Let c be the column of x . Consider the labels of T of family $i - 1$. Suppose such a label appears in column c of T . It cannot be South of x , for then it would be marked and violate T 's (G.11). It cannot be in x , since $\bullet_{\mathcal{G}} \in x$. It cannot be North of x , for then it would be too high and violate T 's (G.1). Thus there is no label of family $i - 1$ in column c .

Since x^\rightarrow is not a box of T , every column East of c contains at most $i - 2$ boxes. Therefore any label of family $i - 1$ East of column c would be too high, contradicting T 's (G.1). Consider a genotype of T involving the $\mathcal{G} \in x^\downarrow$. No label of family $i - 1$ is read before the $\mathcal{G} \in x^\downarrow$. This contradicts T 's (G.8). Thus this miniswap cannot create a label that is too high.

T3: If the $\mathcal{G} \in x$ in either output of T3 were too high, the $\mathcal{G}^+ \in x^\rightarrow$ in T with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$ would violate T 's (G.1).

T4 and T5: Suppose that either T4 or T5 created a label \mathcal{P} that was too high in U . Then in the notation of those miniswaps, U has \mathcal{P} in the edge \bar{x} and we must have $\mathcal{P} \in \{\mathcal{F}\} \cup Z$. By T 's (G.13), there is a label $\mathcal{Q} \in x^\rightarrow$ (possibly virtual) in T with $\text{family } \mathcal{Q} = \text{family}(\mathcal{P}) + 1$. This $\mathcal{Q} \in x^\rightarrow$ is then too high in T , contradicting T 's (G.1).

(G.2): Consider a snake S in T . Since S is a short ribbon (Lemma 6.5), in the region of U defined by S , (G.2) can only be violated by having two $\bullet_{\mathcal{G}^+}$'s in the same row or column. By inspection of the miniswaps, no two $\bullet'_{\mathcal{G}^+}$ s can appear in the same row. If two $\bullet_{\mathcal{G}^+}$'s appear in the same column, the top $\bullet_{\mathcal{G}^+}$ arose from a T4.1 or T4.2 miniswap. However in those cases the edge label $\mathcal{G} \in \text{tail}(S)$ implies $\text{tail}(S) = S$ by T 's (G.7).

Thus we check U 's (G.2) for pairs of snakes S, S' . By Lemma 6.6, say S is southwest of S' . If S is entirely SouthWest of S' , (G.2) preservation is clear. It remains to consider the situations where S and S' share a row or a column.

Suppose the snakes are in the configuration of Lemma 6.6(II). Here $S = \{x\} = \boxed{\bullet_{\mathcal{G}}}$ ($\mathcal{G}, \mathcal{G}^+ \notin x$). So x takes part in a trivial H3 miniswap. By T 's (G.2), the southmost row r' of S' (assumed to be in x 's row) does not contain $\bullet_{\mathcal{G}}$. Thus r' takes part in a H9, B1 or B3 miniswap. H9 and B1 do not introduce a $\bullet_{\mathcal{G}^+}$, so (G.2) holds here. We claim B3 is not possible. If r' participates in a B3 miniswap, then by definition $r' = \{y\} = \boxed{\mathcal{G}}$. It cannot be that $y^\leftarrow = x$, for then S and S' would be the same snake. Let $\mathcal{F} = \text{label}(y^\leftarrow)$. By (G.3), $\mathcal{F} \prec \mathcal{G}$. Hence the $\bullet_{\mathcal{G}} \in x$ means that the $\mathcal{F} \in y^\leftarrow$ is marked. But then y^\leftarrow was adjoined to S' by (R.3), i.e., $r' = \{y^\leftarrow, y\} = \boxed{\mathcal{F} \mid \mathcal{G}}$, a contradiction.

Finally suppose the snakes are in the configuration of Lemma 6.6(III). The two adjacent rows of S and S' are $\begin{array}{c} \boxed{\bullet_{\mathcal{G}}} \\ \bullet_{\mathcal{G}} \mathcal{G}^+ \end{array}$ or $\begin{array}{c} \bullet_{\mathcal{G}} \mathcal{G}^+ \\ \bullet_{\mathcal{G}} \mathcal{G}^+ \end{array}$. Hence S' takes part in a H3 or H8 trivial miniswap.

Let x be the east box of the northmost row of S . The box x takes part in miniswap H6, H7, H8, or T3. If it is miniswap H8, $\bullet_{\mathcal{G}^+} \notin x$ in U . In the other cases, by definition of α , the tableaux produced with $\bullet_{\mathcal{G}^+} \in x$ appears with coefficient 0.

(G.4): We show that U does not violate (G.4) in a given column c .

Case 1: ($\bullet_{\mathcal{G}} \notin c$ in T): By inspection of the miniswaps, c either has labels removed or else a box label of c is pushed onto a lower edge of the same box (and a $\bullet_{\mathcal{G}^+}$ comes into c).

Subcase 1.1: (c is strictly increasing in T): By the above, c is strictly increasing in U .

Subcase 1.2: (c is not strictly increasing in T): Therefore c contains $\{x, x^\uparrow\} = \boxed{\frac{\mathcal{F}}{\mathcal{F}'}}$ in T . By the

above observation, it suffices to show $\{x, x^\uparrow\}$ is not $\boxed{\frac{\mathcal{F}^!}{\mathcal{F}'}}$ or $\boxed{\frac{\mathcal{F}}{\mathcal{F}'}}$ in U . Since $\mathcal{F}^!$ appears in T ,

$\mathcal{F} \prec \mathcal{G}$. Since $\mathcal{F} \in x^\uparrow$ in T , there is no $\bullet_{\mathcal{G}}$ in T northwest of x^\uparrow . Thus in U no $\bullet_{\mathcal{G}^+}$ can appear northwest of x^\uparrow . Hence $\mathcal{F}^! \notin x^\uparrow$ in U . This rules out the first scenario.

We now rule out the second scenario. By Lemma 5.4, in T there is a $\bullet_{\mathcal{G}}$ in some box y West of x in the same row, and furthermore $\mathcal{E}^! \in y^\rightarrow$ in T . By Lemma 5.5, T has $\mathcal{G}' \in \underline{y^\rightarrow}$ (possibly marked), $\mathcal{G}' \in \underline{y^\rightarrow}$ or $(\mathcal{G}')^! \in y^\rightarrow$, where $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$. Let S be the snake containing y .

Subcase 1.2.1: ($y^\rightarrow \in S$): Since $\mathcal{E}^! \in y^\rightarrow$ we have $\{y, y^\rightarrow\} = \text{tail}(S)$. Thus, U has $\bullet_{\mathcal{G}^+} \in y$ or $\bullet_{\mathcal{G}^+} \in y^\rightarrow$. Hence $\mathcal{F} \notin x$ in U .

Subcase 1.2.2: ($y^\rightarrow \notin S$ and neither \mathcal{G} nor \mathcal{G}' appears in y 's column): Then $S = \{y\}$ undergoes H3 and $\text{label}_U(y) = \bullet_{\mathcal{G}^+}$. Hence $\mathcal{F}^! \in x$ in U .

Subcase 1.2.3: ($y^\rightarrow \notin S$ and either \mathcal{G} or \mathcal{G}' appears in y 's column): By Lemma 5.13 applied to T , $\mathcal{G}' = \mathcal{G}$. Hence $y^\rightarrow \in S$, violating the assumption of Subcase 1.2.3.

Case 2: ($\bullet_{\mathcal{G}} \in x$, where x is a box of c in T): Let S be the snake containing x .

Subcase 2.1: ($x \in \text{head}(S)$): Clearly, there is no (G.4) violation except possibly if we apply H5.2 or H5.3, where $\text{label}(x^\rightarrow) = \mathcal{G}$; thus we assume we are using one of these miniswaps. Let \mathcal{F} be the \prec -greatest label appearing in x^\uparrow or $\underline{x^\uparrow}$. Let \mathcal{H} be the \prec -least label appearing in x^\downarrow or $\overline{x^\downarrow}$. After the miniswap, \mathcal{G} appears in x . We show $\mathcal{F} < \mathcal{G} < \mathcal{H}$. Since in T , \mathcal{F} is northwest of $\bullet_{\mathcal{G}}$, $\mathcal{F} \prec \mathcal{G}$ by T 's (G.9). If $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$, then the $\mathcal{F} \in x^\uparrow$ or $\underline{x^\uparrow}$ and the $\mathcal{G} \in x^\rightarrow$ violate T 's (G.12). Hence $\mathcal{F} < \mathcal{G}$. If $\mathcal{H} \prec \mathcal{G}$, then the $\mathcal{H} \in x^\downarrow$ or $\overline{x^\downarrow}$ is marked in T , violating (G.11). If $\mathcal{H} = \mathcal{G}$, then since we are using H5.2 or H5.3, $\mathcal{H} = \mathcal{G} \in x^\downarrow$ so $x^\downarrow \in S$, contradicting $x \in \text{head}(S)$. Hence $\mathcal{G} \prec \mathcal{H}$. So by T 's (G.6), $\mathcal{G} < \mathcal{H}$.

Subcase 2.2: ($x \in \text{tail}(S)$):

Subcase 2.2.1: ($\text{tail}(S)$ is T1, T2 or T3): By Lemma 6.9(IV,V,VII), S has at least two rows and $\text{label}_T(x^\downarrow) = \mathcal{G}$. Suppose there were a label \mathcal{Q} on $\overline{x^\downarrow}$ in T . By T 's (G.4), $\mathcal{Q} < \mathcal{G}$. But then this $\mathcal{Q} \in \overline{x^\downarrow}$ is marked, violating T 's (G.11). Hence $\overline{x^\downarrow}$ is empty. Let \mathcal{F} be the \prec -greatest label appearing in x^\uparrow or $\underline{x^\uparrow}$. Let \mathcal{H} be the \prec -least label appearing in $x^{\downarrow\leftarrow}$ or $\underline{x^\downarrow}$. Since there is $\mathcal{G} \in x^\downarrow$ in T , by T 's (G.4) we have $\mathcal{F} < \mathcal{G} < \mathcal{H}$.

Each of T1, T2, or T3 puts $\mathcal{G} \in x$. The swap does not affect \mathcal{F} nor \mathcal{H} in column c . Now, if the swap puts $\bullet_{\mathcal{G}^+} \in x^\downarrow$ in U , we are done since $\mathcal{F} < \mathcal{G} < \mathcal{H}$. So assume otherwise. Then x^\downarrow takes part in a miniswap H5.1, H5.2 (choosing the first output), or H9. In these cases, U has $\mathcal{G} \in x$ and $\mathcal{G} \in x^\downarrow$. Since $\mathcal{F} < \mathcal{G} < \mathcal{H}$, to show that (G.4) holds, we need $\mathcal{G} \in x^\downarrow$ in U to be marked. If the miniswap was H5.1 or H5.2, then $\bullet_{\mathcal{G}^+} \in x^{\downarrow\leftarrow}$ in U , so U has $\mathcal{G}^! \in x^\downarrow$. If the miniswap is H9, then there is some marked label $\mathcal{E}^! \in x^{\downarrow\leftarrow}$. By T 's (G.10), there is some $\bullet_{\mathcal{G}}$ West of $x^{\downarrow\leftarrow}$ and in its row of T . By Lemma 6.6, this $\bullet_{\mathcal{G}}$ is part of a single-box snake, which undergoes miniswap H3, the $\bullet_{\mathcal{G}}$ becoming $\bullet_{\mathcal{G}^+}$ in the same position. Hence U has $\mathcal{G}^! \in x^\downarrow$.

Subcase 2.2.2: ($\text{tail}(S)$ is T4 or T5): The output of T4.1 and the first output of T4.2 leave c unaffected, so since no other box of c is part of a snake, we are done. The three remaining possibilities (second output of T4.2, T4.3 and T5) are similar, so we argue them together. In these cases, notice x^\uparrow is not part of a snake. Let \mathcal{F} and \mathcal{Z} be as in the description of

these miniswaps. Each places $\mathcal{G} \in \underline{x}$ and $\mathcal{F} \cup \mathcal{Z} \in \bar{x}$. Let \mathcal{H} be the \prec -least label on \underline{x}^\downarrow or \bar{x}^\downarrow . By T 's (G.11) and since $\text{label}_T(\underline{x}) = \bullet_{\mathcal{G}}$, we have $\mathcal{G} \preceq \mathcal{H}$. If $\mathcal{G} < \mathcal{H}$, this \mathcal{H} is not moved by the swap, and column c of U satisfies (G.4) at least south of \underline{x} . Otherwise $\text{family}(\mathcal{G}) = \text{family}(\mathcal{H})$. Since each of these miniswaps says \mathcal{G} or $\mathcal{G} \in \underline{x}^\rightarrow$, if $\mathcal{G} \prec \mathcal{H}$, then by T 's (G.6), $\mathcal{G} = \mathcal{H}$. If $\mathcal{G} = \mathcal{H} \in \bar{x}^\downarrow$, we are in T6, a contradiction. Hence $\mathcal{G} = \mathcal{H} \in \underline{x}^\downarrow$ (and \bar{x}^\downarrow is empty). Consider what happens to \underline{x}^\downarrow during the swap. The analysis to show (G.4) is satisfied south of \underline{x} is essentially the same as in Subcase 2.2.1, so we omit the details.

Finally, we show (G.4) for U north of \underline{x} . Let \mathcal{E} be the \prec -greatest label appearing in T in \underline{x}^\uparrow or \bar{x}^\uparrow . By T 's (G.9), $\mathcal{E} \prec \mathcal{G}$. Each of the miniswaps of interest asserts \mathcal{G} or $\mathcal{G} \in \underline{x}^\rightarrow$. Hence by T 's (G.12), $\mathcal{E} < \mathcal{G}$. Indeed by T 's (G.12), if $\text{family}(\mathcal{E}) \neq \text{family}(\mathcal{F})$ and $\text{family}(\mathcal{E}) \neq \text{family}(\mathcal{Z})$ for any $\mathcal{Z} \in \mathcal{Z}$. Hence by Lemma 5.5 applied to $\underline{x}^\rightarrow$, $\mathcal{E} < \mathcal{F}$. Therefore (G.4) holds in U in c north of \underline{x} .

Subcase 2.2.3: ($\text{tail}(S)$ is T6): No tableau is produced.

Subcase 2.3: ($\underline{x} \in \text{body}(S)$): By Definition-Lemma 6.8 and Lemma 6.9(III), T has $\mathcal{G} \in \underline{x}^\downarrow$ and $\mathcal{G} \in \underline{x}^\rightarrow$. The swap places $\mathcal{G} \in \underline{x}$, and either replaces the $\mathcal{G} \in \underline{x}^\downarrow$ with a $\bullet_{\mathcal{G}^+}$ or else leaves the \mathcal{G} in place there. For the rest of the analysis, one proceeds exactly in the manner given in Subcase 2.2.1.

(G.5): If U violates (G.5), the violation occurs on a horizontal edge e bounding a box of a snake S in T (here e may possibly be a northern boundary edge of S , although only the edge labels of the southern boundary edges are defined as part of S). First assume that we are not in the case of Lemma 6.6(III), so S does not share a column with any other snake.

We break our analysis based on where e appears in relation to S .

Case 1: (e bounds a box of $\text{body}(S)$ but not a box of $\text{head}(S)$ or $\text{tail}(S)$): There is no change of labels on e between T and U . Hence there is no (G.5) violation on e in U .

Case 2: ($e = \underline{x}$ for some $\underline{x} \in \text{head}(S)$): The only head miniswaps that could introduce new edge labels onto e are H6 and H7. In these cases, T has a $\mathcal{G}^+ \in \underline{x}$ that moves to $e = \underline{x}$ in U . If $\mathcal{G}' \in e$ in T with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$, we violate T 's (G.4). Hence the $\mathcal{G}^+ \in e$ in U is the only label of its family on e , as desired.

Case 3: ($e = \bar{x}$ for some $\underline{x} \in \text{head}(S)$): If \underline{x} is the only box of $\text{head}(S)$, we used H1 to move a \mathcal{G} from \underline{x} to \bar{x} . If there is a label $\mathcal{G}' \in \bar{x}$ in T with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$, we violate T 's (G.4). If $|\text{head}(S)| = 2$, no miniswap introduces edges onto a northern edge.

Case 4: ($e = \underline{x}$ for some $\underline{x} \in \text{tail}(S)$): If $|\text{tail}(S)| = 1$, no new edge labels occur during any miniswap (namely T1), so we are done. So assume $|\text{tail}(S)| = 2$. New edge labels on e can only occur when using T3 (second output). Here, T has $\mathcal{G}^+ \in \underline{x}$, while U has $\mathcal{G}^+ \in \underline{x}$. If U violates (G.5), there is $\mathcal{G}' \in \underline{x}$ in T with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$, but this contradicts T 's (G.4).

Case 5: ($e = \bar{x}$ for some $\underline{x} \in \text{tail}(S)$): The miniswaps that could introduce edge labels onto e are T4.2, T4.3 and T5. In the notation of those miniswaps, T has $\bullet_{\mathcal{G}} \in \underline{x}$, an $\mathcal{F}^\uparrow \in \underline{x}^\rightarrow$, and a set \mathcal{Z} of labels ℓ on $\underline{x}^\rightarrow$ such that $\mathcal{F} \prec \ell \prec \mathcal{G}$. In U , all of these labels $\{\mathcal{F}\} \cup \mathcal{Z}$ may have moved to \bar{x} . U violates (G.5) only if it has a label $\mathcal{Q} \in \bar{x}$ with $\text{family}(\mathcal{Q}) = \text{family}(\mathcal{Z})$ for some $\mathcal{Z} \in \{\mathcal{F}\} \cup \mathcal{Z}$. However this \mathcal{Q} and \mathcal{Z} would violate T 's (G.12).

Finally, suppose we are in the case of Lemma 6.6(III). In T the adjacent rows of the snakes are $\begin{array}{|c|c|} \hline \bullet_{\mathcal{G}} & \mathcal{G} \\ \hline \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \end{array}$ or $\begin{array}{|c|c|} \hline \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \end{array}$. Let x be the east box of the south row in either case. If $i_j, k_\ell \in \bar{x}$ in U , then $i_j, k_\ell \in \bar{x}$ in T , since no miniswap affects \bar{x} . By T 's (G.5), $i \neq k$.

(G.6): Consider $\mathcal{H}, \mathcal{H}'$ in T with $\mathcal{H} \prec \mathcal{H}'$ and $\text{family}(\mathcal{H}) = \text{family}(\mathcal{H}')$. Say the eastmost \mathcal{H} in T appears in column c , while the westmost \mathcal{H}' appears in column d . By T 's (G.4) and (G.6), c is West of d . By the swaps' construction, in any $U \in \text{swap}_{\mathcal{G}}(T)$, the westmost \mathcal{H}' in U appears at most one column west of d , while the eastmost \mathcal{H} in U is no further east than column c . In any case, no \mathcal{H}' can be West of an \mathcal{H} in U .

(G.7): Let e be an edge with $\mathcal{H} \in e$ in U . We must show there is no \mathcal{H} West of e in U .

Claim A.1. *Let \mathcal{R} be the region consisting of the leftmost $c - 1$ columns of T (equivalently U). If U has an \mathcal{H} in \mathcal{R} , then T has an \mathcal{H} either in \mathcal{R} or in column c .*

Proof. By inspection of the miniswaps, if there is an \mathcal{H} in column d of U , then there was an \mathcal{H} or (\mathcal{H}) in either d or d^\rightarrow in T . By definition of virtual labels, the existence of (\mathcal{H}) implies the existence of \mathcal{H} further West. The claim follows. \square

Case 1: ($\mathcal{H} \notin e$ in T): We list the miniswaps that put $\mathcal{H} \in e$ in U : H1, H6, H7, T3, T4.2, T4.3, T5. In what follows, x refers to the notation of the miniswap discussed. For H1, locally

we have $T = \begin{array}{|c|} \hline \bullet_{\mathcal{H}} \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \bullet_{\mathcal{H}} \\ \hline \end{array} = U$ (in fact, $\mathcal{H} = \mathcal{G}$). By T 's (G.7), the $\mathcal{H} \in x$ is westmost in T . If the $\mathcal{H} \in \bar{x}(= e)$ is not westmost in U , then by Claim A.1 there is some \mathcal{H} in e 's column in T that takes part in a miniswap leading to an \mathcal{H} West of e in U . Clearly, this \mathcal{H} is not the $\mathcal{H} \in e$. Thus there are two \mathcal{H} 's in e 's column, violating T 's (G.4). For H6, H7 and T3, we have $\mathcal{H} = \mathcal{G}^+$ and $\mathcal{G} \in x$ in U . Thus by U 's (G.4), (G.5) and (G.6), there is not $\mathcal{H} = \mathcal{G}^+$ West of $e = x^\rightarrow$ in U . For the remaining cases, $\mathcal{H} \in \{\mathcal{F}\} \cup Z$ (in the notation of the miniswaps). These labels in x^\rightarrow and x^\rightarrow of T are marked. Hence by Lemma 5.7, they are all westmost in their respective genes in T . Therefore the same labels of e in U are westmost by Claim A.1.

Case 2: ($\mathcal{H} \in e$ in T): No miniswap involving $\mathcal{H} \in e$ both keeps an $\mathcal{H} \in e$ and puts an \mathcal{H} West of e . Thus, if there is an \mathcal{H} West of e in U , then by T 's (G.7) combined with Claim A.1 there is an \mathcal{H} in the column of e in T other than the $\mathcal{H} \in e$. This contradicts T 's (G.4).

(G.3): Consider a row r of T .

Case 1: (The labels of r strictly \prec -increase from left to right, ignoring $\bullet_{\mathcal{G}}$'s): If there is no $\bullet_{\mathcal{G}}$ in r , then only H9, B1 and B3 miniswaps could involve labels of r . Therefore either r is unchanged by $\text{swap}_{\mathcal{G}}$ (or only some labels became marked), or a \mathcal{G} in r is replaced by $\bullet_{\mathcal{G}^+}$. Thus, U 's (G.3) holds for r in this situation.

Otherwise, T has $\bullet_{\mathcal{G}}$ in r , say in box x . By assumption, the exceptional configuration of (G.3) does not occur here. We consider all miniswaps involving a $\bullet_{\mathcal{G}}$. Note that T and U are identical in r both West of x and East of x^\rightarrow . Hence it suffices to study the affect of a miniswap locally at $\{x^\leftarrow, x, x^\rightarrow\}$.

If H1 or H2 applies at x , then locally at x , T looks like $\begin{array}{|c|c|c|} \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{H} \\ \hline \end{array}$ or $\begin{array}{|c|c|c|} \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{H} \\ \hline \end{array}$, where x is the center box. (If \mathcal{F} or \mathcal{H} does not exist, the argument is simplified.) It remains to show $\mathcal{F} \prec \mathcal{G} \prec \mathcal{H}$. By T 's (G.9), $\mathcal{F} \prec \mathcal{G}$. We cannot have $\mathcal{G} = \mathcal{H}$, for then we would apply

H4 or H5.3, instead of H1 or H2. Suppose $\mathcal{H} \prec \mathcal{G}$. Then the $\mathcal{H} \in x^{\rightarrow}$ is marked, and by Lemma 5.13, $\mathcal{G} \in \underline{x^{\rightarrow}}$. Hence we would apply T5 or T6, instead of H1 or H2.

Applying miniswaps H3–H5, H8, B2, B3, T2 or T6 at x clearly preserves (G.3) for r .

Suppose H6, H7 or T3 applies at x in T . Then locally at x , T looks like $\begin{array}{|c|c|c|} \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \end{array}$ or $\begin{array}{|c|c|c|} \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \end{array}$ respectively. (If there is no \mathcal{F} , there is nothing to show.) By T 's (G.9), $\mathcal{F} \prec \mathcal{G}$. Hence (G.3) holds for r in U .

Suppose T1 applies at x . Then at x , T locally looks like $\begin{array}{|c|c|c|} \hline \mathcal{F} & \bullet_{\mathcal{G}} & \mathcal{H} \\ \hline & & \mathcal{G} \\ \hline \end{array}$ (where the $\mathcal{G} \in x^{\downarrow}$ is guaranteed by Lemma 6.9(IV,VII)); again, if \mathcal{F} or \mathcal{H} does not exist, the argument is only easier). We must show $\mathcal{F} \prec \mathcal{G} \prec \mathcal{H}$. By T 's (G.9), $\mathcal{F} \prec \mathcal{G}$. Since we are applying T1, $\mathcal{H} \neq \mathcal{G}$. Now repeat the above argument for H1, H2 above *verbatim*.

Suppose T4 or T5 applies at x . Locally at x , T looks like $\begin{array}{|c|c|c|} \hline \bullet_{\mathcal{G}} & \mathcal{E}^{\downarrow} & \mathcal{F} \\ \hline \bullet_{\mathcal{G}} & \mathcal{E}^{\downarrow} & \mathcal{F} \\ \hline \end{array}$ or $\begin{array}{|c|c|c|} \hline \bullet_{\mathcal{G}} & \mathcal{E}^{\downarrow} & \mathcal{F} \\ \hline \bullet_{\mathcal{G}} & \mathcal{E}^{\downarrow} & \mathcal{F} \\ \hline \end{array}$, while U looks like $\begin{array}{|c|c|c|} \hline \mathcal{G} & \bullet_{\mathcal{G}^+} & \mathcal{F} \\ \hline \mathcal{G} & \bullet_{\mathcal{G}^+} & \mathcal{F} \\ \hline \end{array}$. By assumption $\text{label}_T(x^{\leftarrow}) \prec \mathcal{E}$. Thus, if $\mathcal{G} \prec \mathcal{F}$ (or there is no \mathcal{F}), U 's (G.3) is satisfied. Otherwise $\mathcal{F} \preceq \mathcal{G}$. Then $\mathcal{F} \in x^{\rightarrow\rightarrow}$ is marked in U . By Lemma 5.5, $N_{\mathcal{G}} = N_{\mathcal{E}}$, so by T 's Lemma 5.6, $\mathcal{F} \neq \mathcal{G}$. Therefore $\mathcal{F} \prec \mathcal{G}$. By T 's (G.6), we have $\mathcal{F} < \mathcal{G}$. This three box configuration $\{x, x^{\rightarrow}, x^{\rightarrow\rightarrow}\}$ of U is the exceptional configuration of (G.3).

Case 2: (The labels of r do not strictly increase): Thus r contains the local configuration $\begin{array}{|c|c|c|} \hline \mathcal{H} & \bullet_{\mathcal{G}} & \mathcal{F}^{\downarrow} \\ \hline \mathcal{H} & \bullet_{\mathcal{G}} & \mathcal{F}^{\downarrow} \\ \hline \end{array}$, where $\mathcal{H} > \mathcal{F}$. Call the middle box x . By Lemma 5.5, there is \mathcal{G}' or $\mathcal{G}' \in \underline{x^{\rightarrow}}$ with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$ and $N_{\mathcal{G}'} = N_{\mathcal{F}}$.

If $\mathcal{G}' \neq \mathcal{G}$, then x^{\rightarrow} is not part of the snake containing x . Further by Lemma 5.13, there is no \mathcal{G} or \mathcal{G} in x^{\downarrow} or \underline{x} in T . Hence x takes part in a H3 miniswap, r is unchanged (except for the subscript on the \bullet) and (G.3) is preserved.

If $\mathcal{G}' = \mathcal{G}$, we apply T4, T5 or T6. Recall T6 produces no tableau. In the case of T4.1 and the first output of T4.2, we make no local changes in row r , so U 's (G.3) follows from T 's. The remaining considerations are the second output of T4.2 and the outputs of T4.3 and T5. By T 's (G.9), $\mathcal{H} \prec \mathcal{G}$. Let $\mathcal{E} := \text{label}_T(x^{\rightarrow\rightarrow})$ (if \mathcal{E} does not exist, the argument is trivialized). Since $N_{\mathcal{E}} = N_{\mathcal{G}'} = N_{\mathcal{F}}$, by T 's Lemma 5.6, $\mathcal{E} \neq \mathcal{G}$. If $\mathcal{G} \prec \mathcal{E}$, then U 's (G.3) holds. Otherwise $\mathcal{E} \prec \mathcal{G}$, and the $\mathcal{E} \in x^{\rightarrow\rightarrow}$ in T is marked. Given T 's \mathcal{G} or $\mathcal{G} \in \underline{x^{\rightarrow}}$, it follows by T 's (G.6) that $\mathcal{E} < \mathcal{G}$. Therefore U has the exceptional (G.3) configuration in r .

(G.8): Consider any two genes \mathcal{E} and \mathcal{F} with $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$ and $N_{\mathcal{E}} = N_{\mathcal{F}}$. It suffices to show that in U , every \mathcal{E} is read before every \mathcal{F} , that is:

Claim A.2. *Given (nonvirtual) instances e, f of \mathcal{E} and \mathcal{F} respectively in U , either e is East of f or else e is north of f in the same column.*

Proof. Suppose the claim fails for some fixed choice of e and f . Thus in U either f is North of e in its column or else f is East of e . The first scenario contradicts U 's (G.4), so assume the second occurs.

If U has a label \mathcal{Q} in column c , then T has \mathcal{Q} or \mathcal{Q} in c or c^{\rightarrow} . Thus since U has f East of e , by T 's Lemma 5.6, T has e and f in the same column. By T 's (G.4), e is north of f in T .

Case 1: ($\text{family}(\mathcal{E}) < \text{family}(\mathcal{G})$): We may assume a miniswap moves e West. Since $\text{family}(\mathcal{E}) < \text{family}(\mathcal{G})$, the only such miniswaps are T4 and T5. Hence T has a box x

with $\mathcal{G} \in \underline{x}$ and either $e \in \underline{x}$ or $e \in x$. By T 's (G.4), $f \in \underline{x}$. These miniswaps may move e to $\overline{x^{\leftarrow}}$, but then by definition, they will also move f to x^{\leftarrow} or $\overline{x^{\leftarrow}}$.

Case 2: ($\text{family}(\mathcal{E}) = \text{family}(\mathcal{G})$): We are done unless $\text{swap}_{\mathcal{G}}$ moves e West. The possible miniswaps are H5, B2, B3, T2, and T4. By inspection of the miniswaps, $\mathcal{E} = \mathcal{G}$.

Case 2.1: ($f \in x$): By T 's (G.4), either $e \in \overline{x}$ or else $e \in x^{\uparrow}$ with \overline{x} empty. In all the miniswaps of interest, $\bullet_{\mathcal{E}} \in x^{\uparrow\leftarrow}$. So by T 's (G.2), x^{\leftarrow} contains a genetic label h of some gene \mathcal{H} . The

local picture is either

$\bullet_{\mathcal{E}}$	e
h	f

 or

$\bullet_{\mathcal{E}}$	\star
h	f

 (where \star is some genetic label). By T 's (G.11), h is not marked; hence $\mathcal{H} \succeq \mathcal{E}$. Thus by T 's (G.6), either $\mathcal{H} > \mathcal{E}$ or $\mathcal{H} = \mathcal{E}$. If $\mathcal{H} = \mathcal{E}$, h and f violate Lemma 5.6 for T . Thus $\mathcal{E} < \mathcal{H}$. By T 's (G.3), $\mathcal{H} \prec \mathcal{F}$. Since $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$, we have $\text{family}(\mathcal{H}) = \text{family}(\mathcal{F})$. Indeed by T 's (G.6), $\mathcal{F} = \mathcal{H}^+$. Now since $N_{\mathcal{E}} = N_{\mathcal{F}}$, we violate T 's (G.8) unless there is an \mathcal{E}' above h in its column with $\text{family}(\mathcal{E}') = \text{family}(\mathcal{E})$. If this $\mathcal{E}' \in \overline{x^{\leftarrow}}$, then by (G.11) and (G.6), $\mathcal{E}' = \mathcal{E}$. But then f and this $\mathcal{E} \in \overline{x^{\leftarrow}}$ violate Lemma 5.6 for T . Otherwise, this \mathcal{E}' is North of $x^{\uparrow\leftarrow}$. But then this violates T 's (G.12) together with the $e \in x^{\uparrow}$ or \overline{x} .

Case 2.2: ($f \in \underline{x}$): By T 's (G.4), either $e \in \underline{x}$ or $e \in x$. In the relevant miniswaps, $\bullet_{\mathcal{G}} \in x^{\leftarrow}$. If B2, B3 or T2 applies, then T has $\mathcal{E} \in x^{\leftarrow\downarrow}$; together with f , this violates Lemma 5.6 for T . Hence H5 or T4 applies. Since $f \in \underline{x}$, any H5 miniswap used is H5.1, while any T4 miniswap used is T4.1; both of these fix e , a contradiction.

Case 3: ($\text{family}(\mathcal{E}) > \text{family}(\mathcal{G})$): Neither e nor f is affected by $\text{swap}_{\mathcal{G}}$, so the Claim holds. \square

(G.9): Suppose \mathcal{F} is northwest of $\bullet_{\mathcal{G}^+} \in y$ in U and $\mathcal{G}^+ \preceq \mathcal{F}$; we seek a contradiction. We may suppose such \mathcal{F} is in x or \underline{x} in U .

Case 1: (\mathcal{F} appears in the same position in T): By inspection of the miniswaps, T has $\bullet_{\mathcal{G}} \in y, y^{\leftarrow}$, or y^{\uparrow} . By T 's (G.9), the $\mathcal{F} \in x$ or \underline{x} is not northwest of this box. Hence one of the following subcases occurs:

Subcase 1.1: (T has $\bullet_{\mathcal{G}} \in y^{\leftarrow}$ and y is South of x in its column): Since $\mathcal{G} \prec \mathcal{F}$, T contains no $\mathcal{F}^!$, so by T 's (G.4), $\mathcal{F} < \text{label}_T(y)$ implying $\mathcal{G}^+ < \text{label}_T(y)$. Hence y is not part of any snake in T , contradicting $\bullet_{\mathcal{G}^+} \in y$ in U .

Subcase 1.2: (T has $\bullet_{\mathcal{G}} \in y^{\uparrow}$ and y is East of x in its row): By T 's (G.3) and (G.9), $\mathcal{F} \prec \text{label}_T(y)$. Then $\mathcal{G}^+ \prec \text{label}_T(y)$, so y is not part of any snake in T , contradicting $\bullet_{\mathcal{G}^+} \in y$ in U .

Case 2: (\mathcal{F} does not appear in the same position in T): By inspection of the miniswaps, no label $\mathcal{H} \succ \mathcal{G}^+$ is affected by $\text{swap}_{\mathcal{G}}$. Furthermore labels \mathcal{G}^+ can only be affected if $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$. Since $\mathcal{G}^+ \preceq \mathcal{F}$, this implies $\mathcal{F} = \mathcal{G}^+$ with $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$. By inspection of the miniswaps that affect \mathcal{G}^+ , T has $\mathcal{F} = \mathcal{G}^+ \in x$, while U has $\bullet_{\mathcal{G}^+} \in x$ and $\mathcal{F} = \mathcal{G}^+ \in \underline{x}$. Since $\mathcal{F} \in \underline{x}$ is Southeast of the $\bullet_{\mathcal{G}^+} \in x$ in U , by U 's (G.2) it cannot also be northwest of a $\bullet_{\mathcal{G}^+}$, contradicting our assumption.

(G.11): Consider a label $\mathcal{F}^! \in x$ or \underline{x} in U . By T 's (G.2) and inspection of the miniswaps, T has either \mathcal{F} or $\mathcal{F}^!$ in the same position.

Case 1: (This \mathcal{F} is marked in T): By definition, there is a $\bullet_{\mathcal{G}^+}$ northwest of $\mathcal{F}^!$ in U . If U has a $\bullet_{\mathcal{G}^+}$ South of x and in its column, this contradicts U 's (G.2).

We now show U has no $\bullet_{\mathcal{G}^+}$ North of x and in its column: By inspection of the miniswaps, if $\bullet_{\mathcal{G}^+} \in y$ in U , then T has a $\bullet_{\mathcal{G}}$ northwest of y . By T 's (G.11), T has no $\bullet_{\mathcal{G}}$ in x 's column. Since $\mathcal{F} \prec \mathcal{G}$, by Lemma 5.11 T has no $\bullet_{\mathcal{G}}$ NorthWest of x .

Case 2: (This \mathcal{F} is unmarked in T): By definition, U has a $\bullet_{\mathcal{G}^+}$ northwest of \mathcal{F}^\dagger . By inspection of the miniswaps, if $\bullet_{\mathcal{G}^+} \in y$ in U , then T has a $\bullet_{\mathcal{G}}$ northwest of y . Therefore T has a $\bullet_{\mathcal{G}}$ northwest of said \mathcal{F} . Since the \mathcal{F} is unmarked, it must be that $\mathcal{G} \preceq \mathcal{F}$. But since \mathcal{F}^\dagger appears in U , $\mathcal{F} \prec \mathcal{G}^+$. Hence $\mathcal{F} = \mathcal{G}$. By Lemma 5.11, T has no $\bullet_{\mathcal{G}}$ NorthWest of x . So T has $\bullet_{\mathcal{G}}$ either West of x and in its row or else North of x and in its column. Since $\mathcal{F} = \mathcal{G}$, only the latter case is a concern. In that case, by T 's (G.4) and (G.11), in fact $\mathcal{F} = \mathcal{G} \in x$ and $\bullet_{\mathcal{G}} \in x^\uparrow$. Hence by inspection of the miniswaps, U has $\mathcal{F} = \mathcal{G} \in x^\uparrow$. (Note T4 does not apply at x^\uparrow by T 's (G.7).) Thus U has no $\bullet_{\mathcal{G}^+}$ in x 's column.

(G.10): Consider a label $\mathcal{F}^\dagger \in x$ or \underline{x} in U . By T 's (G.2) and inspection of the miniswaps, T has either \mathcal{F} or \mathcal{F}^\dagger in the same position.

Case 1: (This \mathcal{F} is marked in T): By inspection of the miniswaps, if $\bullet_{\mathcal{G}^+} \in y$ in U , then T has a $\bullet_{\mathcal{G}}$ northwest of y . By T 's (G.11), T has no $\bullet_{\mathcal{G}}$ in x 's column. Since $\mathcal{F} \prec \mathcal{G}$, by Lemma 5.11, T has no $\bullet_{\mathcal{G}}$ NorthWest of x . Hence U has no $\bullet_{\mathcal{G}^+}$ Northwest of x . But by definition, U has a $\bullet_{\mathcal{G}^+}$ northwest of \mathcal{F}^\dagger , so it must be in x 's row.

Case 2: (This \mathcal{F} is unmarked in T): Since \mathcal{F}^\dagger appears in U , $\mathcal{F} \preceq \mathcal{G}$. By definition, U has $\bullet_{\mathcal{G}^+}$ northwest of said \mathcal{F}^\dagger . By inspection of the miniswaps, if $\bullet_{\mathcal{G}^+} \in y$ in U , then $\bullet_{\mathcal{G}}$ appeared northwest of y in T . Hence T has $\bullet_{\mathcal{G}}$ northwest of this \mathcal{F} . Since this \mathcal{F} is unmarked in T , $\mathcal{F} \succeq \mathcal{G}$. Thus $\mathcal{F} = \mathcal{G}$.

By Lemma 5.11, T has no $\bullet_{\mathcal{G}}$ NorthWest of x . Therefore U has no $\bullet_{\mathcal{G}^+}$ NorthWest of x . But U has some $\bullet_{\mathcal{G}^+}$ northwest of x , so it is either West of x in x 's row or North of x in x 's column. In the former case, we are done; in the latter case, we contradict U 's (G.11).

(G.12): Define the **neighborhood** of a box u to be $\text{Neigh}(u) := \{u, u^\leftarrow, u^\uparrow, \underline{u}, \overline{u}^\leftarrow\}$. For a lower edge \underline{u} , let $\text{Neigh}(\underline{u}) := \{\underline{u}, u, u^\leftarrow, \overline{u}, \overline{u}^\leftarrow\}$. Given a (possibly virtual) instance $q \in u$ or \underline{u} in T of the gene \mathcal{Q} , let the **children** of q be all (nonvirtual) \mathcal{Q} 's in U in $\text{Neigh}(u)$ or $\text{Neigh}(\underline{u})$, respectively. Finally define the children of a $\bullet_{\mathcal{G}} \in u$ in T to be those $\bullet_{\mathcal{G}^+}$ in $u, u^\rightarrow, u^\downarrow$ in U . Clearly,

Lemma A.3. *Every q in $U \in \text{swap}_{\mathcal{G}}(T)$ is a child of at least one (possibly virtual) q in T . Also, every $\bullet_{\mathcal{G}^+}$ in $U \in \text{swap}_{\mathcal{G}}(T)$ is a child of at least one $\bullet_{\mathcal{G}}$ in T . \square*

Suppose \mathcal{H} and \mathcal{H}' are instances in T of genes of the same family. By Lemma A.3, it suffices to confirm U 's (G.12) for ℓ a child of \mathcal{H} and ℓ' a child of \mathcal{H}' . To do this, we break into cases depending on the relative position of \mathcal{H} and \mathcal{H}' . By relabeling, we may assume \mathcal{H} west of \mathcal{H}' . Specifically, Cases 1–3 below concern the situation \mathcal{H} NorthWest of \mathcal{H}' . Cases 4–7 consider the case \mathcal{H} southwest of \mathcal{H}' .

For the first three cases, let x, y be boxes in the same row r of T with x West of y . By T 's (G.12), we may assume $\mathcal{H} \in \bar{x}$ or $\mathcal{H} \in x$, as well as $\mathcal{H}' \in y$ or $\mathcal{H}' \in \underline{y}$. By T 's (G.12), there is a $\bullet_{\mathcal{G}}$ in some box z of row r appearing East of x and west of y .

Case 1: (In T , we have \mathcal{H} or $(\mathcal{H}) \in \bar{x}$ and \mathcal{H}' or $(\mathcal{H}') \in \underline{y}$): By T 's (G.4), $\mathcal{H} < \text{label}_T(x)$. By T 's (G.9), $\text{label}_T(x) \prec \mathcal{G}$. Hence $\mathcal{H} < \mathcal{G}$. Since $\text{family}(\mathcal{H}) = \text{family}(\mathcal{H}')$, also $\mathcal{H}' < \mathcal{G}$. Therefore the $\mathcal{H}' \in \underline{y}$ is marked (and is not virtual). By T 's (G.11), this forces $z \neq y$. For convenience, assume $\mathcal{H} \in \bar{x}$. (The argument where this label is virtual is strictly easier.)

By Lemma 5.4, the box labels in y and in every box strictly between y and z are also marked. Let $\text{label}_T(z^\rightarrow) := \mathcal{E}^!$ and note $\mathcal{E} < \mathcal{H}' < \mathcal{G}$ (the first inequality by a combination of T 's (G.3) and (G.4)). Summarizing, T locally looks like one of the following at r :

$$\begin{array}{c} \boxed{\mathcal{H}' \mid \dots \mid \bullet_{\mathcal{G}} \mid \mathcal{E}^! \mid \dots \mid \mathcal{H}'^!} \\ x \qquad z \qquad \qquad y \end{array} \quad \text{or} \quad \begin{array}{c} \boxed{\mathcal{H}' \mid \dots \mid \bullet_{\mathcal{G}} \mid \mathcal{E}^!} \\ x \qquad z \qquad y \end{array} .$$

By T 's (G.2), $\bullet_{\mathcal{G}} \notin x^\uparrow, x^{\uparrow\leftarrow}$. Therefore \bar{x} is not part of a snake and so the only child of $\mathcal{H} \in \bar{x}$ in U is in \bar{x} .

Suppose z is the only box in its snake section, i.e., we apply H1, H2, H3 or T1. If the miniswap is H1, H2 or T1, then \mathcal{G} or \mathcal{G} appears in z^\downarrow or \underline{z} . Hence by Lemma 5.13, $\mathcal{G} \in z^\rightarrow$, so z^\rightarrow is adjoined by (R.3), contradicting z the only box in its snake section. Thus the miniswap is H3 and the unique child of $\bullet_{\mathcal{G}}$ is in z . Moreover y is not part of a snake, or takes part in a trivial H9 miniswap. Hence the unique child of $\mathcal{H}' \in \underline{y}$ is at \underline{y} in U . Thus U 's (G.12) holds in this scenario.

Otherwise the miniswap at z involves z and z^\rightarrow . Then the miniswap is T4, T5 or T6. In these cases, the child of $\bullet_{\mathcal{G}}$ is in either z or z^\rightarrow in U . If the child of $\mathcal{H}' \in \underline{y}$ is in \underline{y} , we are done. If not, $y = z^\rightarrow$ and $\mathcal{H}' \in Z$ (in the notation of the miniswaps). Then the child of \mathcal{H}' is at \bar{z} in U , so the child of \mathcal{H} is not North of the child of \mathcal{H}' and (G.12) holds vacuously.

Case 2: (In T , we have $\mathcal{H} \in x$ and \mathcal{H}' or $\mathcal{H}' \in \underline{y}$): By T 's (G.9), $\mathcal{H} \prec \mathcal{G}$.

Subcase 2.1: ($z = y$): By T 's (G.11), the \mathcal{H}' or $\mathcal{H}' \in \underline{y}$ is not marked; hence $\mathcal{G} \preceq \mathcal{H}'$. Thus $\mathcal{H} \prec \mathcal{G} \preceq \mathcal{H}'$, so $\text{family}(\mathcal{G}) = \text{family}(\mathcal{H})$. Combined with T 's (G.2), this implies the unique child of $\mathcal{H} \in x$ is in x . By T 's (G.3) and (G.9), T has $\mathcal{G}' \in z^\leftarrow$ with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G}) = \text{family}(\mathcal{H})$ and $\mathcal{G}' \prec \mathcal{G}$. Hence by T 's (G.6), $\mathcal{G}' = \mathcal{G}^-$ and $\mathcal{H}' = \mathcal{G}$; moreover by T 's (V.2), this $\mathcal{H}' = \mathcal{G}$ is not virtual, since it is westmost. Hence, locally at r , T is

$$\begin{array}{c} \boxed{\mathcal{H} \mid \dots \mid \mathcal{G}' \mid \bullet_{\mathcal{G}}} \\ x \qquad \qquad \qquad y = z \end{array} \quad (\text{where } \mathcal{H}' = \mathcal{G}, \mathcal{G}' = \mathcal{G}^-).$$

Thus the miniswap involving z is one of H1, H4, H6 and T6. Now H4 and T6 produce no output. If H1 or H6 applies, the child of $\mathcal{H}' \in \underline{y} = \underline{z}$ is northEast of the child of $\mathcal{H} \in x$, so (G.12) is confirmed vacuously.

Subcase 2.2: ($z \neq y$): By T 's (G.4), $\text{label}_T(y) < \mathcal{H}'$. Hence by $\text{family}(\mathcal{H}) \leq \text{family}(\mathcal{G})$, $\text{label}_T(y)$ is marked. By Lemma 5.4, some $\mathcal{E}^! \in z^\rightarrow$. The remainder of this case is argued exactly as Case 1.

Case 3: (In T we have \mathcal{H} or $\mathcal{H} \in \bar{x}$ and $\mathcal{H}' \in y$): Since $\mathcal{H}' \in y$, $\bullet_{\mathcal{G}} \notin y$, so $z \neq y$. By T 's (G.4) and (G.9), $\mathcal{H} < \text{label}_T(x) \prec \mathcal{G}$. Therefore also $\mathcal{H}' < \mathcal{G}$ and so $\mathcal{H}' \in y$ is marked. Since \bar{x} does not participate in the swap, if $\mathcal{H} \in \bar{x}$, this \mathcal{H} has no children, so the (G.12) confirmation is vacuous here. So assume $\mathcal{H} \in \bar{x}$ in T ; since \bar{x} does not participate in the swap, its only child is in the same position in U . In summary, locally at r , T is

$$\boxed{\mathcal{H}' \mid \dots \mid \bullet_{\mathcal{G}} \mid \dots \mid \mathcal{H}'^!} \\ x \qquad z \qquad y$$

Subcase 3.1: ($z = y^\leftarrow$): Consider the miniswap involving z . First suppose z is not the only box in its snake section. By Definition-Lemma 6.8(I,II), it involves y and is T4, T5 or T6. The last miniswap produces no output. For the first two miniswaps, one possibility is that the child of $\mathcal{H}'^! \in y$ is at \bar{z} in U . Here the (G.12) confirmation is vacuous. Otherwise

U 's unique child of $\mathcal{H}' \in y$ is in y ; here the unique child of $\bullet_{\mathcal{G}} \in z$ is in z and so U 's (G.12) holds.

Otherwise, z is the only box in its snake section. Thus H1, H2, H3 or T1 applies. If H1, H2 or T1 applies, then by definition or Lemma 6.9(IV,VII), T has \mathcal{G} or \mathcal{G} in z^\downarrow or z . Hence by Lemma 5.13, $\mathcal{G} \in \underline{y}$, so y is adjoined by (R.3), contradicting z the only box in its snake section. Thus it is H3, and U 's unique children of $\mathcal{H}' \in y$ and $\bullet_{\mathcal{G}} \in z$ are in y, z respectively; hence (G.12) is confirmed here.

Subcase 3.2: ($z \neq y^\leftarrow$): In this case, $\mathcal{H}' \in y$ is not part of a snake in T or takes part in a trivial H9 miniswap; thus its unique child is in y in U . Hence it suffices to check that U has a $\bullet_{\mathcal{G}^+}$ between x and y . By Lemma 5.4(II), since $\text{label}_T(y)$ is marked, $\text{label}_T(z^\rightarrow)$ is also marked. Consider the miniswap involving z . If T has no \mathcal{G} or \mathcal{G} in z^\downarrow or z , then the miniswap must be one of H3, T4 or T5 (it cannot be T1 by Lemma 6.9(IV,VII)). For each of these, a child of $\bullet_{\mathcal{G}} \in z$ appears in z or z^\rightarrow in U . If T has \mathcal{G} or \mathcal{G} in z^\downarrow or z , then by Lemma 5.13 $\mathcal{G} \in \underline{z^\rightarrow}$. Hence the miniswap is T5 or T6. In the former case, a child of $\bullet_{\mathcal{G}} \in z$ appears in z^\rightarrow in U . In the latter case, U does not exist.

Case 4: (In T , $\mathcal{H} \in a$ is southwest of $\mathcal{H}' \in b$): We will use:

Claim A.4. *In U , each child of $\mathcal{H} \in a$ is west of each child of $\mathcal{H}' \in b$.*

Proof. If a is West of b , then the claim holds by the definition of children. So assume a and b are in the same column. By T 's (G.4), $\mathcal{H} = \mathcal{H}'$ and $\mathcal{H} \in a$ is marked. Hence $\mathcal{H} \prec \mathcal{G}$. By Lemma 5.4(II), T has a $\bullet_{\mathcal{G}}$ in some box z West of a and in its row. By Lemma 5.4(II), every box label strictly between z and a is also marked. Thus by T 's (G.11), T has no $\bullet_{\mathcal{G}}$ in any column East of z and west of a . Furthermore by T 's (G.2), T has no $\bullet_{\mathcal{G}}$ Northwest of z .

Summarizing, T is locally: $z \begin{array}{|c|c|c|} \hline \bullet_{\mathcal{G}} & \dots & \mathcal{H}' \\ \hline \end{array} \begin{array}{|c|} \hline \mathcal{H} \\ \hline \end{array} b$. Hence b is not part of any snake, and so U 's unique child of $\mathcal{H}' \in b$ is in b . \square

No child of $\mathcal{H} \in a$ is North of a^\uparrow and no child of $\mathcal{H}' \in b$ is South of \underline{b} . Hence if a is at least two rows below b , the (G.12) confirmation is vacuous.

Subcase 4.1: (a is exactly one row south of b): By inspection of the miniswaps, a child of $\mathcal{H} \in a$ can only appear North of a child of $\mathcal{H}' \in b$ if $\bullet_{\mathcal{G}} \in a^\uparrow, b^\leftarrow$. Then by T 's (G.2), $a^\uparrow = b^\leftarrow$.

Here, the $\mathcal{H}' \in b$ has a child South of b only if T is locally $\begin{array}{|c|c|} \hline \bullet_{\mathcal{G}} & \mathcal{H}' \\ \hline \mathcal{H} & \\ \hline \end{array} = \begin{array}{|c|c|} \hline \bullet_{\mathcal{G}} & \mathcal{G}^+ \\ \hline \mathcal{G} & \\ \hline \end{array}$ and $\mathcal{H}' = \mathcal{G}^+ \in b$

is part of a T3 miniswap, i.e., $\begin{array}{|c|c|} \hline \bullet_{\mathcal{H}} & \mathcal{H}' \\ \hline \mathcal{H} & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline \mathcal{H} & \bullet_{\mathcal{H}'} \\ \hline * & \mathcal{H}' \\ \hline \end{array}$ (here $*$ $\in \{\mathcal{H}', \bullet_{\mathcal{H}'}\}$; the uncertainty is irrelevant). By T 's (G.3), it follows $\text{label}_U(b^\rightarrow)$ is not marked and $\text{label}_U(b^\rightarrow) \neq \mathcal{H}'$. Thus we confirm (G.12).

Subcase 4.2: (a is in the same row r as b): Suppose $\mathcal{H}' \in b$ has a child South of b . Then b is part of a H6, H7 or T3 miniswap. So U 's unique child of $\mathcal{H}' \in b$ is at \underline{b} and U has $\bullet_{\mathcal{G}} \in b$. By T 's (G.2), T has no $\bullet_{\mathcal{G}} \in a^\uparrow$, so no child of $\mathcal{H} \in a$ is in a^\uparrow . Thus U 's (G.12) is confirmed.

Otherwise no child of $\mathcal{H}' \in b$ is South of b . If a child of $\mathcal{H} \in a$ is in $\overline{a^\leftarrow}$, we used T4 or T5 at a ; thus U has $\bullet_{\mathcal{G}^+} \in a$ and (G.12) is confirmed. Thus it remains to consider the scenario that a child of $\mathcal{H} \in a$ is at a^\uparrow in U . This scenario is impossible: By inspection of the miniswaps, $\mathcal{G} = \mathcal{H}$ and T has $\bullet_{\mathcal{G}} \in a^\uparrow$. Let $\text{label}_T(b^\uparrow) := \mathcal{E}$ (by T 's (G.2), $\bullet_{\mathcal{G}} \notin b^\uparrow$).

$\bullet_{\mathcal{H}}$	\cdots	\mathcal{E}
\mathcal{H}	\cdots	\mathcal{H}'

Locally T is $\begin{array}{|c|c|} \hline \mathbf{a} & \mathbf{b} \\ \hline \end{array}$ (where $\mathcal{H} = \mathcal{G}$). By T 's (G.4), either $\mathcal{E} < \mathcal{H}'$ or $\mathcal{E} = \mathcal{H}'$. If $\mathcal{E} = \mathcal{H}'$, then by T 's (G.4), the $\mathcal{H}' \in \mathbf{b}$ is marked; hence $\mathcal{H}' \prec \mathcal{H}$, contradicting T 's (G.6). Thus $\mathcal{E} < \mathcal{H}'$. Then $\mathcal{E} < \mathcal{H}$, so $\mathcal{E} \in \mathbf{b}^\uparrow$ is marked. By Lemma 5.5, T has a label \mathcal{H}'' or $\mathcal{H}'' \in \mathbf{b}^\uparrow$ with $\text{family}(\mathcal{H}'') = \text{family}(\mathcal{H})$. With T 's $\mathcal{H}' \in \mathbf{b}$, this violates T 's (G.4).

Case 5: (In T , $\mathcal{H} \in \mathbf{a}$ is southwest of $\mathcal{H}' \in \mathbf{b}$): By T 's (G.4), \mathbf{a} is West of \mathbf{b} .

Subcase 5.1: (\mathbf{a} South of \mathbf{b}): By the definition of children, every child of $\mathcal{H} \in \mathbf{a}$ is southwest of every child of $\mathcal{H}' \in \mathbf{b}$, so there is nothing to confirm here.

Subcase 5.2: (\mathbf{a} and \mathbf{b} are in the same row): By the definition of children, every child of $\mathcal{H} \in \mathbf{a}$ is west of every child of $\mathcal{H}' \in \mathbf{b}$. Every child of $\mathcal{H}' \in \mathbf{b}$ is north of \mathbf{b} . Thus we are only concerned with the cases that a child of $\mathcal{H} \in \mathbf{a}$ is north of \mathbf{a} , so we assume this. Moreover, by inspection of the miniswaps, we may assume T has $\bullet_{\mathcal{G}} \in \mathbf{a}$ or $\bullet_{\mathcal{G}} \in \mathbf{a}^\leftarrow$.

If $\bullet_{\mathcal{G}} \in \mathbf{a}^\leftarrow$, then \mathbf{a} is part of a T4 or T5 miniswap. Hence the unique child of $\mathcal{H} \in \mathbf{a}$ is in \mathbf{a}^\leftarrow or $\overline{\mathbf{a}^\leftarrow}$; the unique child of $\mathcal{H}' \in \mathbf{b}$ is in \mathbf{b} ; and U has $\bullet_{\mathcal{G}^+} \in \mathbf{a}$. So (G.12) is confirmed.

Otherwise, T has $\bullet_{\mathcal{G}} \in \mathbf{a}$ and moreover $\mathcal{H} = \mathcal{G}$. Let $\mathcal{E} := \text{label}_T(\mathbf{a}^\rightarrow)$. Locally between \mathbf{a} and \mathbf{b} , T is: $\mathbf{a} \begin{array}{|c|c|c|} \hline \bullet_{\mathcal{H}} & \mathcal{E} & \cdots & \mathcal{H}' \\ \hline \mathcal{H} & & & \end{array} \mathbf{b}$. If $\mathcal{E}^\dagger \in \mathbf{a}^\rightarrow$, then by Lemma 5.13, $\mathcal{H} \in \mathbf{a}^\rightarrow$, and so \mathbf{a} swaps by T6 and there is no tableau U . Thus $\mathcal{E} \in \mathbf{a}^\rightarrow$ is not marked, and by T 's (G.3), $\text{family}(\mathcal{E}) = \text{family}(\mathcal{H})$. By T 's (G.6), either $\mathcal{E} = \mathcal{H}$ or $\mathcal{E} = \mathcal{H}^+$. In the former case, H4 applies at \mathbf{a} , producing no U . In the latter case, H6 applies and one confirms (G.12) by inspection. (If $\mathbf{b} = \mathbf{a}^\rightarrow$, one checks, as was done in Case 4, that the configuration of the final sentence of (G.12) does not occur.)

Case 6: (In T , $\mathcal{H} \in \mathbf{a}$ is southwest of $\mathcal{H}' \in \mathbf{b}$): By T 's (G.4), \mathbf{a} is SouthWest of \mathbf{b} . We may assume \mathbf{a} is one row South of \mathbf{b} and U has a child of $\mathcal{H} \in \mathbf{a}$ in \mathbf{a}^\uparrow (otherwise the (G.12) check is vacuous). The unique child of $\mathcal{H}' \in \mathbf{b}$ is in \mathbf{b} . Hence, $\mathcal{H} = \mathcal{G}$ and T has $\bullet_{\mathcal{G}} \in \mathbf{a}^\uparrow$. Consider the miniswap involving \mathbf{a}^\uparrow . It is B2, B3, T1, T2, T3, T4 or T5. If it is B2, B3, T2 or T3, then locally T is

$$\mathbf{a} \begin{array}{|c|c|c|c|} \hline \bullet_{\mathcal{H}} & \mathcal{H} & \cdots & \star \\ \hline \mathcal{H} & & & \mathcal{H}' \\ \hline \end{array} \mathbf{b} \quad \text{or} \quad \mathbf{a} \begin{array}{|c|c|c|c|} \hline \bullet_{\mathcal{H}} & \mathcal{H}^+ & \cdots & \star \\ \hline \mathcal{H} & & & \mathcal{H}' \\ \hline \end{array} \mathbf{b}.$$

Since by T 's (G.2), the $\bullet_{\mathcal{H}} \in \mathbf{a}^\uparrow$ is the only $\bullet_{\mathcal{H}}$ in its row, this contradicts T 's (G.12). Thus it is T1, T4 or T5. Let $\mathcal{E} := \text{label}_T(\mathbf{a}^\uparrow)$. By T 's (G.3) and (G.4), $\mathcal{E} < \mathcal{H}'$. Hence $\mathcal{E} < \mathcal{H}$, and so $\mathcal{E}^\dagger \in \mathbf{a}^\uparrow$. By Lemma 5.13, T has $\mathcal{H} \in \mathbf{a}^\uparrow$. Therefore the miniswap is T5 and one confirms (G.12) directly.

Case 7: (T has $\mathcal{H} \in \mathbf{a}$ southwest of $\mathcal{H}' \in \mathbf{b}$): By T 's (G.4) and (G.5), \mathbf{a} is West of \mathbf{b} . Hence by the definition of children, U has every child of $\mathcal{H} \in \mathbf{a}$ west of every child of $\mathcal{H}' \in \mathbf{b}$. We may assume that \mathbf{a} and \mathbf{b} are in the same row and that some child of $\mathcal{H} \in \mathbf{a}$ is north of \mathbf{a} , for otherwise the (G.12) confirmation is vacuous. Then T has $\bullet_{\mathcal{G}} \in \mathbf{a}$ or $\bullet_{\mathcal{G}} \in \mathbf{a}^\leftarrow$.

Subcase 7.1: ($\bullet_{\mathcal{G}} \in \mathbf{a}$): Here $\mathcal{G} = \mathcal{H}$. By T 's (G.4), $\text{label}_T(\mathbf{b}) < \mathcal{H}'$, whence $\text{label}_T(\mathbf{b}) < \mathcal{H}$. Therefore, $\text{label}_T(\mathbf{b})$ is marked. By Lemma 5.4(II), $\text{label}_T(\mathbf{a}^\rightarrow)$ is also marked, and so by Lemma 5.13, $\mathcal{H} \in \mathbf{a}^\rightarrow$. Thus the miniswap involving \mathbf{a} is T6, and U does not exist.

Subcase 7.2: ($\bullet_{\mathcal{G}} \in \mathbf{a}^\leftarrow$): The miniswap involving \mathbf{a}^\leftarrow is either T4 or T5. U has $\bullet_{\mathcal{G}^+} \in \mathbf{a}$ and the unique child of $\mathcal{H} \in \mathbf{a}$ is in \mathbf{a}^\leftarrow or $\overline{\mathbf{a}^\leftarrow}$. The (G.12) confirmation is therefore clear.

(G.13): By inspection of the miniswaps, if $\mathcal{E}^! \in x$ or \underline{x} in U , then T has \mathcal{E} or $\mathcal{E}^!$ in the same location. Thus there are two cases:

Case 1: (This \mathcal{E} is marked in T): By T 's (G.13), there is an \mathcal{F} or (\mathcal{F}) in \underline{x} with $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$. If $\mathcal{F} \in \underline{x}$ is nonvirtual, then since it appears in the same place in U , U 's (G.13) holds. Thus suppose T has $(\mathcal{F}) \in \underline{x}$. We check the conditions for (\mathcal{F}) to appear in \underline{x} in U . Let U^* be U with \mathcal{F} added in \underline{x} .

((V.1) holds, i.e., $\mathcal{F} \in \underline{x}$ is not marked in U^*): By Lemma 5.9, T has $\bullet_{\mathcal{G}} \in x^{\leftarrow}$. Note $\mathcal{F} \succeq \mathcal{G}$, since otherwise $(\mathcal{F}) \in \underline{x}$ would be marked in T , a contradiction. If $\mathcal{F} = \mathcal{G}$, then T5 or T6 would apply at $\{x^{\leftarrow}, x\}$, contradicting $\mathcal{E}^! \in x$ or $\mathcal{E}^! \in \underline{x}$ in U . Thus $\mathcal{F} \succ \mathcal{G}$, as desired.

((V.2) holds, i.e. U has an \mathcal{F} West of \underline{x}): By T 's (V.2), T has an \mathcal{F} West of x . This remains true for U since no swap removes a nonvirtual genetic label without putting one further west.

((G.1) holds for U^*): Immediate from T 's $(\mathcal{F}) \in \underline{x}$ and U 's (G.1).

((G.4) holds for U^*): We have $\mathcal{E} < \mathcal{F} < \text{label}_T(x^{\downarrow})$. Since $\mathcal{F} \succ \mathcal{G}$, $\text{label}_T(x^{\downarrow}) = \text{label}_U(x^{\downarrow})$. Hence U^* does not violate (G.4) locally, so by U 's (G.4), we are done.

((G.5) holds for U^*): Since $\mathcal{E}^! \in x$ or \underline{x} in U , by inspection of the miniswaps, T and U have the same set of nonvirtual labels on \underline{x} .

((G.6) holds for U^*): If there is a $\tilde{\mathcal{F}}$ West of x 's column in U , there is one west of x 's column in T . Hence we are done by T 's (V.3).

((G.8) holds for U^*): It suffices to show that in the column reading word of U^* (with \mathcal{F} placed in \underline{x}), no \mathcal{E} is read after the $\mathcal{F} \in \underline{x}$. By T 's (V.3), this is true in T . By inspection of the miniswaps, \mathcal{E} does not appear West of x in U . Hence by U^* 's (G.4), we are done.

((G.9) holds for U^*): Since U has $\mathcal{E}^! \in x$ or \underline{x} , U has a $\bullet_{\mathcal{G}^+}$ northwest of x . So by U 's (G.2), there is no a $\bullet_{\mathcal{G}^+}$ southeast of \underline{x} .

((G.12) holds for U^*): This follows from the following two claims:

Claim A.5. *If $\text{family}(\mathcal{F}) = \text{family}(\tilde{\mathcal{F}})$, then U has no $\tilde{\mathcal{F}}$ SouthEast of \underline{x} .*

Proof. By T 's (G.11) and T 's $\mathcal{E}^!$ in x 's column, T has a $\bullet_{\mathcal{G}}$ northWest of x . Hence by T 's (G.2), T has no $\bullet_{\mathcal{G}}$ SouthEast of x . Thus by the (G.12) condition of T 's (V.3), T has no $\tilde{\mathcal{F}}$ SouthEast of $(\mathcal{F}) \in \underline{x}$. A child of $\tilde{\mathcal{F}}$ will only be South of its parent if T has $\tilde{\mathcal{F}} \in y$ and U has $\tilde{\mathcal{F}} \in y$. Thus U has no $\tilde{\mathcal{F}}$ SouthEast of \underline{x} that is a child of a nonvirtual $\tilde{\mathcal{F}}$ in T .

The remaining concern is $(\tilde{\mathcal{F}}) \in \underline{y}$ in T with a child SouthEast of x in U . By inspection of the miniswaps, this $(\tilde{\mathcal{F}})$ cannot have a child South or East of \underline{y} . Therefore y must be SouthEast of x in T . Again by inspection of the miniswaps, T has a $\bullet_{\mathcal{G}} \in y^{\leftarrow}$ or a $\bullet_{\mathcal{G}} \in y$. This is impossible by T 's (G.2), recalling T 's $\bullet_{\mathcal{G}}$ northWest of x . \square

Claim A.6. *If U has an $\tilde{\mathcal{F}}$ NorthWest of \underline{x} with $\text{family}(\mathcal{F}) = \text{family}(\tilde{\mathcal{F}})$, then this $\tilde{\mathcal{F}}$ and the $\mathcal{F} \in \underline{x}$ satisfy U^* 's (G.12).*

Proof. By assumption, $\mathcal{E}^!$ remains in x or \underline{x} in U . By Lemma 5.9, T has $\bullet_{\mathcal{G}} \in x^{\leftarrow}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$. By T 's (V.1), $\mathcal{G} \preceq \mathcal{F}$. If $\mathcal{G} = \mathcal{F}$, then $\{x^{\leftarrow}, x\}$ is a tail of type T5 or T6; however, these miniswaps do not leave $\mathcal{E}^!$ in place, a contradiction. Hence $\mathcal{G} \prec \mathcal{F}$.

Therefore by T 's (G.5) and Lemma 5.13, T has no \mathcal{G} or \mathcal{G} in x^{\leftarrow} 's column. Hence H3 applies at x^{\leftarrow} , and U has $\bullet_{\mathcal{G}^+} \in x^{\leftarrow}$.

By T 's (G.12), there is no $\tilde{\mathcal{F}}$ NorthWest of \bar{x} . Since no miniswap moves a label more than one box north, it suffices to consider a $\tilde{\mathcal{F}}$ in T that is in either y , \underline{y} , \bar{y} or y^\perp , where y is in x 's row. By T 's (G.2) and $\bullet_{\mathcal{G}} \in x^{\leftarrow}$, T has no $\bullet_{\mathcal{G}}$ strictly northwest of x^{\leftarrow} . Hence no such $\tilde{\mathcal{F}}$ in these four positions can move North. Therefore any $\tilde{\mathcal{F}}$ NorthWest of $\mathcal{F} \in \underline{x}$ in U satisfies (G.12), in view of the $\bullet_{\mathcal{G}^+} \in x^{\leftarrow}$. Note that the forbidden configuration from the final sentence of (G.12) cannot occur since $\mathcal{F} \neq \mathcal{G}$, whereas the forbidden configuration forces $\mathcal{F} = \mathcal{G}$. \square

Case 2: (This \mathcal{E} is unmarked in T): By Lemma A.3, since there is a $\bullet_{\mathcal{G}^+}$ northwest of this \mathcal{E} in U , there is a $\bullet_{\mathcal{G}}$ northwest of x in T . Moreover, by U 's (G.11), x^{\leftarrow} is a box of U , and hence of T . Since this \mathcal{E} is unmarked in T , $\mathcal{E} \succeq \mathcal{G}$. Since this \mathcal{E} is marked in U , $\mathcal{E} \prec \mathcal{G}^+$. Thus, $\mathcal{E} = \mathcal{G}$. Let \mathcal{H} be the gene (if it exists) with $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$. If T has $\mathcal{H} \in \underline{x}$, then since it appears in the same place in U , U 's (G.13) holds. Thus assume $\mathcal{H} \notin \underline{x}$ in T .

Subcase 2.1: (In T , $\bullet_{\mathcal{G}} \in x^{\leftarrow}$): The miniswap applied at x^{\leftarrow} is H5.2 or T4.2. We are done by the following lemma:

Lemma A.7. *If T is a \mathcal{G} -good tableau where H5.2 or T4.2 applies and $U \in \text{swap}_{\mathcal{G}}(T)$, then all prescribed \mathcal{H} 's from the outputs of H5.2 and T4.2 are valid virtual labels in the sense of (V.1)–(V.3).*

Proof. Consider such a miniswap. We may assume that U contains an output with a prescribed \mathcal{H} , say in \underline{x} . Let U^* be U with \mathcal{H} added in \underline{x} .

((V.1) holds, i.e., $\mathcal{H} \in \underline{x}$ is not marked in U^*): By assumption, $\mathcal{G} < \mathcal{H}$ and so $\mathcal{G}^+ \preceq \mathcal{H}$.

((V.2) holds, i.e., some \mathcal{H} appears in U West of x): Since $\mathcal{H} \in \underline{x}$ in T , by T 's (V.2), some \mathcal{H} appears in T West of x . By inspection of the miniswaps, this \mathcal{H} has a child, which is West of x in U .

((G.1) holds in U^*): Immediate from the (G.1) condition of T 's (V.3).

((G.4) holds in U^*): By U 's (G.4), the only concern is an $\tilde{\mathcal{H}}$ in x 's column of U with $\text{family}(\tilde{\mathcal{H}}) = \text{family}(\mathcal{H})$. By Lemma A.3, this $\tilde{\mathcal{H}}$ is a child of an $\tilde{\mathcal{H}}$ in T . Since $\mathcal{G} < \tilde{\mathcal{H}}$, this $\tilde{\mathcal{H}}$ in T is in the same location as its unique child in U , contradicting the (G.4) condition of T 's (V.3).

((G.5) holds in U^*): Neither miniswap in question affects the nonvirtual labels on \underline{x} , so we are done by the (G.5) condition of T 's (V.3).

((G.6) holds in U^*): All labels of this family appear in the same places in T and U , so we are done by the (G.6) condition of T 's (V.3).

((G.8) holds in U^*): First suppose there is a nonballot genotype G_{U^*} of U^* , with the $\mathcal{H} \in \underline{x}$ taken. Since no labels of $\text{family}(\mathcal{H}) + k$ (for $k \geq 0$) are moved by $\text{swap}_{\mathcal{G}}$, by T 's (G.8) there is no violation among those families. By U 's (G.8), it suffices to consider the possibility that in $\text{word}(G_{U^*})$ the selected \mathcal{G} appears after the \mathcal{H} . However, if such a G_{U^*} exists, then by T 's (G.4) and inspection of the miniswaps, it follows there is a \mathcal{G} West of x in T ; this contradicts $\mathcal{H} \in \underline{x}$ in T .

((G.9) holds in U^*): By U 's (G.2), U has no \bullet_{g^+} southeast of x^{\leftarrow} .

((G.12) holds in U^*): By T 's (G.2) and (G.12), T has no $\tilde{\mathcal{H}}$ SouthEast of \underline{x} with $\text{family}(\tilde{\mathcal{H}}) = \text{family}(\mathcal{H})$. The same is true in U , as $\text{swap}_{\mathcal{G}}$ does not affect $\tilde{\mathcal{H}}$. It remains to consider such $\tilde{\mathcal{H}}$ in U that are NorthWest of \underline{x} . Since $\text{swap}_{\mathcal{G}}$ does not affect $\tilde{\mathcal{H}}$, by T 's (G.12), no $\tilde{\mathcal{H}}$ is North of \bar{x} in either T or U . Thus assume $\tilde{\mathcal{H}}$ is West of x and either in its row or on a top edge of its row. By assumption, T 's $\bullet_{\mathcal{G}} \in x^{\leftarrow}$ becomes the desired $\bullet_{g^+} \in x^{\leftarrow}$ in U . \square

Subcase 2.2: (In T , $\mathcal{D}^! \in x^{\leftarrow}$): If $\mathcal{G} \in \underline{x}$, this \mathcal{G} is not in a snake of T . Otherwise $\mathcal{G} \in x$ and H9 applies at x . By Lemma 5.5, T has some $\tilde{\mathcal{G}} \in \underline{x^{\leftarrow}}$ or $(\tilde{\mathcal{G}}) \in \underline{x^{\leftarrow}}$ with $\text{family}(\tilde{\mathcal{G}}) = \text{family}(\mathcal{G})$ and $N_{\tilde{\mathcal{G}}} = N_{\mathcal{D}}$. By Lemma 5.6, $\tilde{\mathcal{G}} \neq \mathcal{G}$. Therefore by T 's (G.6), $\tilde{\mathcal{G}} = \mathcal{G}^-$. By T 's (G.11), T has a $\bullet_{\mathcal{G}}$ northWest of x^{\leftarrow} . Hence this $\tilde{\mathcal{G}} \in \underline{x^{\leftarrow}}$ is nonvirtual and marked. Thus by T 's (G.13), there is some $\tilde{\mathcal{H}} \in \underline{x^{\leftarrow}}$ or $(\tilde{\mathcal{H}}) \in \underline{x^{\leftarrow}}$ with $\text{family}(\tilde{\mathcal{H}}) = \text{family}(\tilde{\mathcal{G}}) + 1$ and $N_{\tilde{\mathcal{H}}} = N_{\tilde{\mathcal{G}}}$. We claim there is an \mathcal{H} South of x and in its same column, with $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1 = \text{family}(\tilde{\mathcal{H}})$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$. Certainly by $N_{\tilde{\mathcal{H}}} = N_{\tilde{\mathcal{G}}} = N_{\mathcal{D}}$, there is such an \mathcal{H} somewhere in T . By Lemma 5.6, it is located as described. Now if $\mathcal{H} \in \underline{x}$, we are done. Otherwise by T 's (G.4), $\mathcal{H} \in x^{\downarrow}$. But now $\text{label}_T(x^{\leftarrow\downarrow}) \prec \mathcal{H}$ by T 's (G.3) and $\text{label}_T(x^{\leftarrow\downarrow}) > \tilde{\mathcal{H}}$ by T 's (G.4), a contradiction. \square

APPENDIX B. PROOF OF PROPOSITION 9.5

We check that (G.1)–(G.13) are preserved. Let $T \in \text{revswap}_{g^+}(U)$, where U is \mathcal{G}^+ -good. Below, the proof of property (G. i) only possibly depends on earlier properties (G. i). We also show that the virtual labels prescribed by reverse miniswaps L2.3, L4.3 and L4.5 are valid virtual labels in the sense of (V.1)–(V.3). This appears as Lemma B.4 in the section ‘Consistency of the prescribed virtual labels,’ located between the arguments for (G.12) and (G.13).

(G.1): Suppose T has $\mathcal{Q} \in x$ or $\mathcal{Q} \in \underline{x}$ that is too high. By U 's (G.1), the label \mathcal{Q} does not appear in the same place in U . Hence \mathcal{Q} is placed in x or \underline{x} in T by some reverse miniswap. We consider which reverse miniswap this might be. By U 's (G.1), \mathcal{Q} does not appear anywhere in U north of \underline{x} . Hence by visual inspection, the only miniswap to consider is L1.1. However to apply L1.1, we have by assumption $\mathcal{Q} \in x^{\uparrow}$ in U . Since this is impossible by U 's (G.1), T cannot have any label too high.

(G.2): By Lemma 8.4, ladders lie in distinct rows and columns; hence T has no $\bullet_{\mathcal{G}}$ NorthWest of another. Since revswap is defined by its action on rows, T has at most one $\bullet_{\mathcal{G}}$ in any row.

Suppose T has $\bullet_{\mathcal{G}} \in x$ North of $\bullet_{\mathcal{G}} \in y$ and in the same column. These boxes are in ladders of U . By Lemma 8.4, x and y are in the same ladder of U . By Lemma 8.3, $x = y^{\uparrow}$ and the two boxes are $\begin{array}{|c|} \hline \mathcal{G} \\ \hline \bullet_{g^+} \\ \hline \end{array}$ in U . Hence, in order to have $\bullet_{\mathcal{G}} \in y$ in T , we must apply L1.2 or L3 to y . By definition, U 's $\mathcal{G} \in x$ means that L1.2 does not apply. L3 requires $\mathcal{G} \in \bar{y}$, contradicting U 's (G.4).

(G.3): It is enough to confirm (G.3) for an arbitrary fixed row \mathcal{R} of T . If \mathcal{R} does not intersect any ladder of U , then T 's (G.3) is confirmed in \mathcal{R} by U 's (G.3).

Otherwise by Lemma 8.4, \mathcal{R} intersects a single ladder L . Let $r := \mathcal{R} \cap L$. Let x be the westmost box of r .

If r is L1, confirmation is trivial unless $\mathcal{G} \in x^\uparrow$. In that case, locally at x , $\text{revswap}_{\mathcal{G}^+}$ results in $\begin{array}{|c|c|c|} \hline & \mathcal{G} & \\ \hline \mathcal{F} & \bullet_{\mathcal{G}^+} & \mathcal{H} \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|} \hline & \star & \\ \hline \mathcal{F} & \mathcal{G} & \mathcal{H} \\ \hline \end{array}$ (here $\star = \bullet_{\mathcal{G}}$, but this is not important to us). (If either \mathcal{F} or \mathcal{H} does not exist, the argument is simpler.) We need $\mathcal{F} \prec \mathcal{G} \prec \mathcal{H}$. By U 's (G.9), $\mathcal{F} \preceq \mathcal{G}$. Since r is L1, $\mathcal{F} \neq \mathcal{G}$, so $\mathcal{F} \prec \mathcal{G}$. If $\mathcal{Q} \in x^{\uparrow\rightarrow}$ in U , by U 's (G.3) and (G.4) then $\mathcal{G} \prec \mathcal{Q} \preceq \mathcal{H}$. Otherwise $\bullet_{\mathcal{G}^+} \in x^{\uparrow\rightarrow}$ in U . By U 's (G.11), $\mathcal{H} \in x^\rightarrow$ is not marked in U . Thus $\mathcal{G}^+ \preceq \mathcal{H}$ and $\mathcal{G} \prec \mathcal{H}$, as desired.

Suppose r is L2. Then $\text{label}_U(x) = \mathcal{G}$, while $\text{label}_T(x) \in \{\mathcal{G}, \bullet_{\mathcal{G}}\}$. Hence, T 's (G.3) is confirmed from U 's (G.3).

If r is L3, r does not change and we are done.

Suppose r is L4.1, L4.2 or L4.3. Locally at r , the reverse miniswap is

$$\begin{array}{|c|c|c|c|} \hline \mathcal{F} & \mathcal{G} & \bullet_{\mathcal{G}^+} & \mathcal{H} \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline \mathcal{F} & \bullet & \mathcal{G}^+ & \mathcal{H} \\ \hline \end{array} \quad (\text{the labels of } \underline{x} \text{ in } T \text{ are not displayed}).$$

We need to show $\mathcal{F} \prec \mathcal{G}^+ \prec \mathcal{H}$. (If \mathcal{F} or \mathcal{H} do not exist, the argument is simpler.) By U 's (G.3), $\mathcal{F} \prec \mathcal{G}$, so $\mathcal{F} \prec \mathcal{G}^+$. By U 's (G.12) (final sentence), $\mathcal{H} \neq \mathcal{G}^+$ and $\mathcal{H} \in x^{\rightarrow\rightarrow}$ is not marked in U . Thus $\mathcal{G}^+ \prec \mathcal{H}$.

Lastly, suppose r is L4.4 or L4.5. Locally at r , $\begin{array}{|c|c|c|c|} \hline \mathcal{E} & \mathcal{G} & \bullet & \mathcal{H} \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline \mathcal{E} & \bullet & \mathcal{F} & \mathcal{H} \\ \hline \end{array}$, where \mathcal{F} is as in the definitions of L4.4 and L4.5.

First we check T has no (G.3) violation between \mathcal{E} and \mathcal{F} . If $\mathcal{E} < \mathcal{F}$, this is obvious. If $\mathcal{E} > \mathcal{F}$, \mathcal{E} and \mathcal{F} form the exceptional configuration of (G.3) in T . Suppose $\text{family}(\mathcal{E}) = \text{family}(\mathcal{F})$. By U 's (G.3) and (G.7), $\mathcal{E} \neq \mathcal{F}$. Hence by U 's (G.6), $\mathcal{E} \prec \mathcal{F}$, as desired.

Now we check $\mathcal{F} \prec \mathcal{H}$.

Case 1: ($\mathcal{G} \prec \mathcal{H}$): Since $\mathcal{F} \preceq \mathcal{G}$, $\mathcal{F} \prec \mathcal{H}$ follows.

Case 2: ($\mathcal{G} \succeq \mathcal{H}$): By U 's (G.3), $\mathcal{G} > \mathcal{H}$ and $\mathcal{H}^! \in x^{\rightarrow\rightarrow}$ in U .

Subcase 2.1: ($\text{family}(\mathcal{F}) < \text{family}(\mathcal{H})$): Then $\mathcal{F} \prec \mathcal{H}$, and we are done.

Subcase 2.2: ($\text{family}(\mathcal{F}) = \text{family}(\mathcal{H})$): By Lemma 5.10 applied to U , $\mathcal{F} \neq \mathcal{H}$. Hence by U 's (G.6), $\mathcal{F} \prec \mathcal{H}$.

Subcase 2.3: ($\text{family}(\mathcal{F}) > \text{family}(\mathcal{H})$): We derive a contradiction. By Lemma 5.5, U has $\mathcal{G}' \in x^{\rightarrow\rightarrow}$ or $\mathcal{G}' \in x^{\rightarrow\rightarrow}$ with $N_{\mathcal{H}} = N_{\mathcal{G}'}$ and $\text{family}(\mathcal{G}) = \text{family}(\mathcal{G}')$. If $\mathcal{G} = \mathcal{G}'$, then by U 's (G.7), $\mathcal{G} \in x^{\rightarrow\rightarrow}$ in U . However $x^{\rightarrow\rightarrow}$ is southeast of a $\bullet_{\mathcal{G}^+}$ in x^\rightarrow in U , contradicting U 's (V.1). Hence $\mathcal{G} \neq \mathcal{G}'$, so by U 's (G.6) $\mathcal{G}' \succeq \mathcal{G}^+$. If $\mathcal{G}' \succ \mathcal{G}^+$, then by U 's (G.6) there must be a \mathcal{G}^+ in the column of x^\rightarrow . This \mathcal{G}^+ is not South of x^\rightarrow by U 's (G.12). It is also not north of x^\rightarrow by U 's (G.9). Thus, $\mathcal{G}' = \mathcal{G}^+$.

Since $N_{\mathcal{G}^+} = N_{\mathcal{H}^-}$, $N_{\mathcal{G}} = N_{\mathcal{H}^-}$. By U 's (G.8), since U has $\mathcal{G} \in x$, every \mathcal{H}^- appears before x in column reading order. By U 's (G.4) and (G.6), \mathcal{H}^- does not appear east of $x^{\rightarrow\rightarrow}$'s column. By U 's (G.12), \mathcal{H}^- does not appear North of x^\rightarrow and in its column. Since $\bullet_{\mathcal{G}^+} \in x^\rightarrow$, \mathcal{H}^- is not in x^\rightarrow . By U 's (G.11), \mathcal{H}^- is not South of x^\rightarrow and in its column. Thus \mathcal{H}^- appears only in x 's column and is north of x . By U 's (G.12), it then follows that $\mathcal{H}^- \in \bar{x}$ in U . But then since $N_{\mathcal{H}^-} = N_{\mathcal{G}}$ and $\mathcal{H}^- < \mathcal{F}$, this contradicts the definition of \mathcal{F} .

(G.4): Consider an arbitrary column c of U ; we show that (G.4) holds for c in T .

Case 1: (U has no $\bullet_{\mathcal{G}^+}$ in c):

Subcase 1.1: (c is $<$ -increasing in U): By inspection of the reverse miniswaps, it is clear that c is $<$ -increasing in T .

Subcase 1.2: (c contains $\begin{array}{|c|} \hline \mathcal{F} \\ \hline \mathcal{F}^\dagger \\ \hline \end{array}$): Suppose the depicted \mathcal{F} is in x in U . Since $\mathcal{F}^\dagger \in x^\downarrow$ in U , $\mathcal{F} \preceq \mathcal{G}$. So there are two possibilities:

Subcase 1.2.1: ($\mathcal{F} \prec \mathcal{G}$): Since $\mathcal{F} \prec \mathcal{G}$, x and x^\downarrow are not in ladders of U . Therefore, $\mathcal{F} \in x$ and $\mathcal{F} \in x^\downarrow$ in T ; however, we do not know *a priori* which of these \mathcal{F} 's are marked. We must show T has unmarked $\mathcal{F} \in x$ and $\mathcal{F}^\dagger \in x^\downarrow$. We will need the following

Lemma B.1. *Let U be \mathcal{G}^+ -good and $T \in \text{revswap}_{\mathcal{G}^+}(U)$. Suppose $\bullet_{\mathcal{G}} \in y$ in T . Then in U , either $\mathcal{G} \in y$ or $\bullet_{\mathcal{G}^+} \in y$.*

Proof. Since $\bullet_{\mathcal{G}} \in y$ in T , y is in a ladder of U . Thus the lemma is immediate from the definition of ladders. \square

Since unmarked $\mathcal{F} \in x$ in U , U has no $\bullet_{\mathcal{G}^+}$ northwest of x . By U 's (G.3) or (G.4), U has no \mathcal{G} northwest of x . Hence by Lemma B.1, T has no $\bullet_{\mathcal{G}}$ northwest of x ; hence unmarked $\mathcal{F} \in x$ in T .

By definition of marked labels, U has $\bullet_{\mathcal{G}^+}$ in some box z northwest of x^\downarrow in U . By inspection of the miniswaps, there is a $\bullet_{\mathcal{G}}$ northwest of z in T . Hence $\mathcal{F}^\dagger \in x^\downarrow$ in T .

Subcase 1.2.2: ($\mathcal{F} = \mathcal{G}$): By inspection of the miniswaps, $\bullet_{\mathcal{G}} \in x$ in T and no other box or edge of the column was changed. Hence we are done by U 's (G.4).

Case 2: (U has a $\bullet_{\mathcal{G}^+}$ in c): By U 's (G.11), U has no marked labels in c . Hence by U 's (G.4), the labels of U strictly $<$ -increase down c , ignoring the $\bullet_{\mathcal{G}^+}$. Suppose the $\bullet_{\mathcal{G}^+}$ in c is in x . By inspection, a (G.4) violation in c can only occur from L1.1, L4.4 or L4.5 applied at x . If we apply L1.1 at x , then locally at c , $\begin{array}{|c|} \hline \mathcal{G} \\ \hline \bullet \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \bullet \\ \hline \mathcal{G} \\ \hline \end{array}$, while the rest of the column is unchanged, so c satisfies (G.4) in T .

Suppose we apply L4.4 or L4.5 at x . By definition U has $\mathcal{G} \in x^\leftarrow$ and there is no $\mathcal{G}^+ \notin \underline{x}$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$. Let \mathcal{E} be the \prec -greatest gene in c north of \bar{x} and let \mathcal{H} be the \prec -least gene in c south of \underline{x} . (If either of these fails to exist, the argument is simplified.) Let \mathcal{F} be the \prec -least element of $Z^\#$ in the notation of L4.4 and L4.5. It remains to show

Claim B.2.

- (I) $\mathcal{G} < \mathcal{H}$, and
- (II) either $\mathcal{E} < \mathcal{F}$, or $\mathcal{E} = \mathcal{F}$ with \mathcal{E} an unmarked label in x^\uparrow and $\mathcal{F} < \mathcal{G}$.

Proof. (I): By U 's (G.11), the \mathcal{H} in c is not marked in U , so $\mathcal{G} \prec \mathcal{H}$. If $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G})$, then by U 's (G.12) $\mathcal{H} \in \underline{x}$ rather than x^\downarrow . Then by U 's (G.7), $\mathcal{H} \neq \mathcal{G}$. But then by U 's (G.6), $\mathcal{H} = \mathcal{G}^+ \in \underline{x}$, contradicting our assumption. Thus $\mathcal{G} < \mathcal{H}$.

(II): By U 's (G.9), $\mathcal{E} \preceq \mathcal{G}$. Hence by U 's (G.6), either $\mathcal{E} = \mathcal{G}$ or else $\mathcal{E} < \mathcal{G}$.

Case A: ($\mathcal{E} = \mathcal{G}$): By Lemma 5.6, $Z = \emptyset$ (in the notation of L4.4 and L4.5). Further by U 's (G.7), $\mathcal{E} \in x^\uparrow$ (rather than \bar{x}). Hence locally c changes as $\begin{array}{|c|} \hline \mathcal{G} \\ \hline \bullet \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \bullet \\ \hline \mathcal{G} \\ \hline \end{array}$, while the rest of c is unchanged. Then c satisfies (G.4) in T .

Case B: ($\mathcal{E} < \mathcal{G}$): If $\mathcal{F} = \mathcal{G}$, then $\mathcal{E} < \mathcal{F}$, as desired. So by U 's (G.4), we may assume $\mathcal{F} < \mathcal{G}$ and furthermore that $\text{family}(\mathcal{E}) \geq \text{family}(\mathcal{F})$. Then by Lemma 5.8, there is an $\tilde{\mathcal{E}} \in Z$ with $\text{family}(\tilde{\mathcal{E}}) = \text{family}(\mathcal{E})$.

If $\tilde{\mathcal{E}} \neq \mathcal{E}$, then by U 's (G.6) $\mathcal{E} = \tilde{\mathcal{E}}^+$. Since $N_{\tilde{\mathcal{E}}} = N_{\mathcal{G}} \geq 1$, $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$ and $N_{\mathcal{E}} = N_{\mathcal{G}^+}$. Hence by U 's (G.6) and (G.8), U has \mathcal{G}^+ in c south of \bar{x} . By U 's (G.9), it is south of \underline{x} . By U 's (G.12), it is not South of \underline{x} . Hence $\mathcal{G}^+ \in \underline{x}$ in U , contradicting the assumptions of L4.4 and L4.5.

Thus $\tilde{\mathcal{E}} = \mathcal{E}$. By Lemma 5.6, $\tilde{\mathcal{E}} = \min Z$, so $\tilde{\mathcal{E}} = \mathcal{F}$. By U 's (G.7), the \mathcal{E} in c in U is in x^\uparrow . Since in T , $\bullet_{\mathcal{G}} \in x^{\leftarrow}$ and $\mathcal{E} < \mathcal{G}$, $\mathcal{E}^\dagger \in x$ in T .

We must show that unmarked $\mathcal{E} \in x^\uparrow$ in T . By U 's (G.2), U has no $\bullet_{\mathcal{G}^+}$ northwest of x . By U 's (G.3) and (G.4), U has no \mathcal{G} northwest of x^\uparrow . Hence by Lemma B.1, T has no $\bullet_{\mathcal{G}}$ northwest of x^\uparrow , so T has unmarked $\mathcal{E} \in x^\uparrow$. \square

(G.5): We are only concerned with reverse miniswaps that produce or relocate edge labels. That is L2.2, L3, L4.2, L4.4 and L4.5. In L2.2, we create an edge label \mathcal{G} on \underline{x} , while U has $\mathcal{G} \in x$. Hence by U 's (G.4), there is no other label of \mathcal{G} 's family on \underline{x} . This verifies T 's (G.5) in this situation. A similar argument applies for L3 and L4.2.

The arguments for L4.4 and L4.5 are similar. Consider any $Q \in A''$ or \tilde{A}'' . If (G.5) fails in T , we may assume U has $Q' \in \underline{x}^{\rightarrow}$ with $\text{family}(Q) = \text{family}(Q')$. If $Q \neq \mathcal{G}$, then by U 's (G.4), $Q < \mathcal{G}$, so $Q' \in \underline{x}^{\rightarrow}$ is marked in U , violating U 's (G.11). Thus $Q = \mathcal{G}$. By U 's (G.7), $Q' \neq \mathcal{G}$. Hence by U 's (G.6), $Q' = \mathcal{G}^+$, contradicting that L4.4 or L4.5 applies.

(G.6): We will use

Lemma B.3. *Let U be \mathcal{G}^+ -good. If \mathcal{H} appears in column c of $T \in \text{revswap}_{\mathcal{G}^+}(U)$, then U has \mathcal{H} in column c or column c^{\leftarrow} .*

Proof. By inspection of the reverse miniswaps. \square

Suppose i_b appears West of i_a in T . Then by Lemma B.3, i_b appears west of i_a in U . Therefore by U 's (G.4), either $b = a$ or i_b is West of i_a in U . Thus by U 's (G.6), $b \leq a$.

(G.7): Let $Q \in \underline{x}$ in column c of T . First suppose U has $Q \in \underline{x}$. By U 's (G.7), U has no Q West of column c . Hence by Lemma B.3, T has no Q West of column c , as desired.

So suppose U has $Q \notin \underline{x}$. Thus T 's $Q \in \underline{x}$ was created by one of the reverse miniswaps L2.2, L3, L4.2, L4.4 or L4.5. In each case, we are done by Lemma B.3, provided that we know U has no Q West of c . This is by assumption in L2.2 and L4.2. This holds for L3 by U 's (G.7). For L4.4, $Q \in A''$ (in the terminology of L4.4). If $Q = \mathcal{G}$, we are done by assumption; otherwise we are done by U 's (G.7), since $Q \in A$ (in the terminology of L4.4). For L4.5, $Q \in \tilde{A}''$ (in the terminology of L4.5), and we are done by U 's (G.7), since $Q \in A$ (in the terminology of L4.5).

(G.8): Suppose $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$. By T 's (G.4), to show T 's (G.8), it suffices to show no \mathcal{F} is East of any \mathcal{E} in T .

Let e be the westmost instance of \mathcal{E} and f the eastmost instance of \mathcal{F} in U . By U 's (G.8), e is east of f in U . Swapping does not move e West and moves f at most one column East. We may therefore assume e and f are in the same column c of U . We may also assume the swap moves f East, i.e., the swap involving f is L4.4 or L4.5 (say at $\{x, x^{\rightarrow}\}$). If $e \in \bar{x}$, then

T 's westmost \mathcal{E} is in x^\rightarrow or $\underline{x}^\rightarrow$, and there is no (G.8) violation. Hence by U 's (G.4), we may assume $e \in x^\uparrow$. Note that $N_{\mathcal{F}} = N_{\mathcal{G}}$, U has $\mathcal{G} \in x$, and either $\mathcal{F} = \mathcal{G}$ or $\mathcal{F} \in \bar{x}$.

By U 's (G.3), $\mathcal{E} \prec \mathcal{Q} := \text{label}_U(x^{\uparrow\rightarrow})$. By U 's (G.9), $\mathcal{Q} \preceq \mathcal{G}$. By Lemma 5.6, $\mathcal{Q} \neq \mathcal{G}$. Hence $\mathcal{Q} \prec \mathcal{G}$, so by U 's (G.6), $\mathcal{Q} < \mathcal{G}$. By U 's (G.4) and Lemma 5.8, for every family $(\mathcal{F}) < i < \text{family}(\mathcal{G})$, there is a label $\mathcal{H}^i \in \bar{x}$ with $\text{family}(\mathcal{H}^i) = i$ and $N_{\mathcal{H}^i} = N_{\mathcal{E}} = N_{\mathcal{F}} = N_{\mathcal{G}}$. By Lemma 5.6, $\mathcal{Q} \neq \mathcal{F}$ and $\mathcal{Q} \neq \mathcal{H}^i$. Hence by U 's (G.6), \mathcal{Q} is one of \mathcal{E}^+ , \mathcal{F}^+ , $(\mathcal{H}^i)^+$. Hence, U has a \mathcal{G}^+ in c^\rightarrow with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$. By U 's (G.9) and (G.12), it follows that $\mathcal{G}^+ \in \underline{x}^\rightarrow$; this contradicts the assumptions of L4.4 or L4.5.

(G.9): Let $\mathcal{F} \succeq \mathcal{G}$. Suppose $\mathcal{F} \in x$ (or $\mathcal{F} \in \underline{x}$) is northwest of $\bullet_{\mathcal{G}} \in y$ in T . By inspection of the reverse miniswaps, U has an \mathcal{F} northwest of \underline{x} .

Suppose $\mathcal{F} \succ \mathcal{G}$. By U 's (G.9), U has no $\bullet_{\mathcal{G}^+}$ southeast of x . Hence by Lemma B.1, $\mathcal{G} \in y$ in U . Thus by U 's (G.3) and (G.4), $\mathcal{F} \preceq \mathcal{G}$, contradicting $\mathcal{F} \succ \mathcal{G}$.

Thus suppose $\mathcal{F} = \mathcal{G}$. By Lemma B.1, in U either $\bullet_{\mathcal{G}^+} \in y$ or $\mathcal{G} \in y$. Suppose $\mathcal{G} \in y$. By Lemma 5.10, y is not southEast of x , i.e., x and y are in the same column. Hence by U 's (G.4), U has $\mathcal{G}^! \in y$. Hence y is not in a ladder of U , contradicting T 's $\bullet_{\mathcal{G}} \in y$. Thus, U has $\bullet_{\mathcal{G}^+} \in y$. By Lemma 5.12, it follows that x and y are in the same row or column. By U 's (G.9), in fact $y = x^\downarrow$ or $y = x^\rightarrow$. Now, by inspection of the reverse miniswaps, we conclude that $\mathcal{F} \notin x$ (respectively, $\mathcal{F} \notin \underline{x}$) or $\bullet_{\mathcal{G}} \notin y$ in T , contradicting our initial assumptions.

(G.10): Consider $\mathcal{F}^! \in x$ or $\mathcal{F}^! \in \underline{x}$ in T .

Case 1: (\mathcal{F} is not in the same location in U): By assumption, $\mathcal{F} \prec \mathcal{G}$. By inspection of the reverse miniswaps, the $\mathcal{F}^!$ in question appears in T as a result of L4.4 or L4.5. By definition $\bullet_{\mathcal{G}} \in x^\leftarrow$ in T and we are done.

Case 2: (\mathcal{F} is in the same location in U): By U 's (G.3) and (G.4), U has no \mathcal{G} northwest of x . However, T has a $\bullet_{\mathcal{G}} \in z$ northwest of x . Hence by Lemma B.1, U has a $\bullet_{\mathcal{G}^+}$ northwest of x . Since $\mathcal{F} \prec \mathcal{G} \prec \mathcal{G}^+$, this means U 's $\mathcal{F} \in x$ or \underline{x} is marked

By Lemma 5.4, U has a $\bullet_{\mathcal{G}^+} \in w$ West of x and in the same row. If T has $\bullet_{\mathcal{G}} \in w$ or w^\leftarrow , we are done. Otherwise, L1.1 applies at w . Then U has $\mathcal{G} \in w^\uparrow$, and by Lemma 5.5 U has a $\tilde{\mathcal{G}} \in \underline{x}$ (possibly virtual) in U such that $\text{family}(\mathcal{G}) = \text{family}(\tilde{\mathcal{G}})$; this contradicts U 's (G.12).

(G.11): Consider $\mathcal{F}^! \in x$ or $\mathcal{F}^! \in \underline{x}$ in T .

Case 1: (\mathcal{F} is not in the same location in U): By assumption, $\mathcal{F} \prec \mathcal{G}$. By inspection of the miniswaps, the $\mathcal{F}^!$ in question appears in T as a result of L4.4 or L4.5. Therefore we have $\mathcal{F} < \mathcal{G}$ and $N_{\mathcal{F}} = N_{\mathcal{G}}$.

Suppose $\bullet_{\mathcal{G}} \in y$ in T , where y is in x 's column. By U 's (G.2), $\bullet_{\mathcal{G}^+} \notin y$ in U . Hence by Lemma B.1, U has $\mathcal{G} \in y$. This contradicts Lemma 5.6 (applied to U), since U has $\mathcal{G} \in y$ and $\mathcal{F} \in \bar{x}^\leftarrow$, where $\mathcal{F} < \mathcal{G}$ and $N_{\mathcal{F}} = N_{\mathcal{G}}$.

Case 2: (\mathcal{F} is in the same location in U): By the first paragraph of the argument of (G.10) Case 2, this \mathcal{F} is marked in U . Suppose T has $\bullet_{\mathcal{G}} \in y$, where y is in x 's column. By U 's (G.11), U has no $\bullet_{\mathcal{G}^+}$ in x 's column. Hence by Lemma B.1, U has $\mathcal{G} \in y$. By U 's (G.4), since $\mathcal{F} \prec \mathcal{G}$, y is South of x and in its column. By assumption, T has some $\bullet_{\mathcal{G}}$ northwest of x . With $\bullet_{\mathcal{G}} \in y$, this contradicts T 's (G.2).

(G.12): Let x, z be boxes in row r with x West of z . Suppose $\text{family}(\mathcal{F}) = \text{family}(\mathcal{F}')$. Cases 1–3 consider the case U has \mathcal{F} NorthWest of \mathcal{F}' . By U 's (G.12), we may assume U

has $\mathcal{F} \in \bar{x}$ or $\mathcal{F} \in x$ as well as $\mathcal{F}' \in z$ or $\mathcal{F}' \in \underline{z}$. By U 's (G.12), U has a $\bullet_{\mathcal{G}^+}$ in some box y of row r that is East of x and west of z . Cases 4–7 consider the case U has \mathcal{F} southwest of \mathcal{F}' .

Case 1: (U has \mathcal{F} or $(\mathcal{F}) \in \bar{x}$ and \mathcal{F}' or $(\mathcal{F}') \in \underline{z}$): By U 's (G.4), $\mathcal{F} < \text{label}_U(x)$. By U 's (G.9), $\text{label}_U(x) \prec \mathcal{G}^+$. Hence $\mathcal{F} < \mathcal{G}^+$. Since $\text{family}(\mathcal{F}) = \text{family}(\mathcal{F}')$, also $\mathcal{F}' < \mathcal{G}^+$. Therefore U 's $\mathcal{F}' \in \underline{z}$ is marked (and is not virtual). By U 's (G.11), it follows $y \neq z$. Moreover by U 's Lemma 5.4, $\text{label}_U(y^{\rightarrow})$ is marked. By U 's (G.4), $\mathcal{G} \notin z$. Hence z is not in a ladder, and so T has $\mathcal{F}' \in \underline{z}$. For convenience, assume $\mathcal{F} \in \bar{x}$. (The argument where this label is virtual is strictly easier).

Case 1.1: (x and y are *not* in the same ladder of U): By Lemma 8.4, x is not in any ladder. Hence T has $\mathcal{F} \in \bar{x}$. By inspection, unless the reverse miniswap applied at y is L1.1, T has $\bullet_{\mathcal{G}} \in y$ or $\bullet_{\mathcal{G}} \in y^{\leftarrow} \neq x$. Hence although T has \mathcal{F} NorthWest of \mathcal{F}' , there is no (G.12) violation. If the reverse miniswap is L1.1, then U has $\mathcal{G} \in y^{\uparrow}$. By Lemma 5.13, U has $(\mathcal{G}) \in \underline{y^{\rightarrow}}$; with the $\mathcal{G} \in y^{\uparrow}$, this violates the (G.12) condition of U 's (V.3).

Case 1.2: (x and y are in the same ladder of U): Then $y = x^{\rightarrow}$ and the ladder row is type L4. If it is L4.1, L4.2 or L4.3, then U has $\mathcal{G} \in y^{\leftarrow}$, $\bullet_{\mathcal{G}^+} \in y$ and $\mathcal{G}^+ \in \underline{y}$ with $\text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$. Since $\text{label}_U(y^{\rightarrow})$ is marked, this contradicts U 's (G.12).

Hence it is L4.4 or L4.5. By Lemma 5.5, $\underline{y^{\rightarrow}}$ contains a label \mathcal{F}'' with $\text{family}(\mathcal{F}'') = \text{family}(\mathcal{F})$ and a (possibly virtual) label \mathcal{G}'' with $\text{family}(\mathcal{G}'') = \text{family}(\mathcal{G})$. By Lemma 5.5, $N_{\mathcal{F}''} = N_{\mathcal{G}''}$. By U 's (G.12), there is no label of \mathcal{F} 's family North of y and in y 's column. By U 's (G.11), there is no label of that family South of y and in y 's column. Hence there is no label of \mathcal{F} 's family in y 's column. Hence by U 's (G.6) and (G.7), $\mathcal{F}'' = \mathcal{F}^+$. By U 's (G.12), there is no label in y 's column of the same family as \mathcal{G} . Hence by U 's (G.6) and Lemma 5.10, $\mathcal{G}'' = \mathcal{G}^+$. So $N_{\mathcal{F}} = N_{\mathcal{G}}$ and $\mathcal{F} \in Z$ (in the notation of L4.4/L4.5). Thus T has $\mathcal{F} \in \underline{y}$ or $\mathcal{F} \in y$. Therefore by T 's (G.3) and (G.4), T has $\mathcal{F} \in \underline{y}$, and so has \mathcal{F} southWest of \mathcal{F}' , in agreement with (G.12).

Case 2: (U has $\mathcal{F} \in x$ and \mathcal{F}' or $(\mathcal{F}') \in \underline{z}$): By U 's (G.9), $\mathcal{F} \prec \mathcal{G}^+$.

Subcase 2.1: ($y = z$): By U 's (G.11), $(\mathcal{F}')^{\downarrow} \notin \underline{z}$; hence $\mathcal{G}^+ \preceq \mathcal{F}'$. Thus $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G}^+) = \text{family}(\mathcal{G})$. Hence by U 's (G.9), $\text{family}(\text{label}_U(y^{\leftarrow})) = \text{family}(\mathcal{F})$. So by U 's (G.6) and (G.9), $\text{label}_U(y^{\leftarrow}) = (\mathcal{F}')^{\leftarrow}$ and $\mathcal{F}' = \mathcal{G}^+$. Hence by U 's (G.6) and (G.7), U has (nonvirtual) $\mathcal{G}^+ \in \underline{y}$. The applicable reverse miniswap is then L4.1, L4.2 or L4.3. Hence in T , \mathcal{F} is not NorthWest of \mathcal{F}' , and (G.12) holds.

Subcase 2.2: ($y \neq z$): By U 's (G.4), $\text{label}_U(z) < \mathcal{F}'$. Therefore $\text{label}_U(z)$ is marked; by Lemma 5.4, $\text{label}_U(y^{\rightarrow})$ is marked. The reverse swap does not affect \underline{z} . Hence we may assume T has $\mathcal{F}' \in \underline{z}$.

If the reverse swap moves the $\mathcal{F} \in x$ South, then T has \mathcal{F} southWest of \mathcal{F}' , in accordance with (G.12). No reverse swap can move the $\mathcal{F} \in x$ North. Hence we may assume T has $\mathcal{F} \in x$ or $\mathcal{F} \in x^{\rightarrow}$. If $\text{label}_T(x^{\rightarrow}) = \mathcal{F}$, then $y = x^{\rightarrow}$ and $\mathcal{F} = \mathcal{G}$, so T has $y \ni \mathcal{G} \succeq \text{label}_T(y^{\rightarrow})$, contradicting T 's (G.3). Thus $\text{label}_T(x) = \mathcal{F}$. By Lemma 5.5, U has a label of the same family as \mathcal{G} on $\underline{y^{\rightarrow}}$. Hence the reverse miniswap involving y is not L1.1, since $\mathcal{G} \in y^{\uparrow}$ would violate U 's (G.12). Hence by inspection of the reverse miniswaps, T has $\bullet_{\mathcal{G}} \in y$ or $\bullet_{\mathcal{G}} \in y^{\leftarrow}$, in accordance with T 's (G.12).

Case 3: (U has \mathcal{F} or $(\mathcal{F}) \in \bar{x}$ and $\mathcal{F}' \in z$): Since $\mathcal{F}' \in z$, $\bullet_{\mathcal{G}^+} \notin z$, so $y \neq z$. By U 's (G.4), $\mathcal{F} < \text{label}_U(x)$ whereas by U 's (G.9), $\text{label}_U(x) \prec \mathcal{G}^+$; hence $\mathcal{F} < \mathcal{G}^+$. Therefore also $\mathcal{F}' < \mathcal{G}$ and so $\mathcal{F}' \in z$ is marked in U . The box z is not part of a ladder, so T has $\mathcal{F}' \in z$. No

reverse swap can move the $\mathcal{F} \in \bar{x}$ North. If it moves South, it will be southWest of z in T , so no (G.12) violation ensues. Hence we may assume T has $\mathcal{F} \in \bar{x}$. By Lemma 5.4, y^\rightarrow contains a marked label \mathcal{E}^\downarrow , and so by Lemma 5.5, U has a (possibly virtual) $\mathcal{G}' \in \underline{y^\rightarrow}$ with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$.

The reverse miniswap involving y is not L1.1, for $\mathcal{G} \in y^\uparrow$ violates U 's (G.12), together with $\mathcal{G}' \in \underline{y^\rightarrow}$. Thus, T has $\bullet_{\mathcal{G}} \in y$ or $\bullet_{\mathcal{G}} \in y^\leftarrow$. Unless T has $\bullet_{\mathcal{G}} \in y^\leftarrow$ and $x = y^\leftarrow$, this does not violate T 's (G.12). Suppose $x = y^\leftarrow$ and T has $\bullet_{\mathcal{G}} \in x$. The reverse miniswap involving y is L4. Hence U has $\mathcal{G} \in x$. By Lemma 5.5, either $\text{family}(\mathcal{E}) = \text{family}(\mathcal{F})$ or U has some $\mathcal{F}'' \in \underline{y^\rightarrow}$ with $\text{family}(\mathcal{F}'') = \text{family}(\mathcal{F})$. By Lemma 5.5, $N_{\mathcal{G}'} = N_{\mathcal{E}} = N_{\mathcal{F}''}$. By U 's (G.11) and (G.12), y 's column does not contain a label of \mathcal{F} 's family. Hence by U 's (G.6) and (G.7), $\mathcal{F}'' = \mathcal{F}^+$. By U 's (G.9) and (G.12), if y 's column contains a label of the same family as \mathcal{G} , it is on y . By U 's (G.6) and (G.7), it can only be \mathcal{G}^+ ; this contradicts U 's (G.12) (last sentence). Thus U has $\mathcal{G}^+ \notin \underline{y}$, and so $\mathcal{G}' = \mathcal{G}^+$. The reverse miniswap is L4.4 or L4.5. We have $N_{\mathcal{F}} = N_{\mathcal{G}}$. Thus $\mathcal{F} \in Z$ (in the notation of L4.4 and L4.5), contradicting that T has $\mathcal{F} \in \bar{x}$.

Case 4: (U has $\mathcal{F} \in a$ southwest of $\mathcal{F}' \in b$): Say b is in row r . No reverse miniswap can move \mathcal{F} North or move \mathcal{F}' further South than b^\downarrow . Hence unless a is in row r , T has \mathcal{F} southwest of \mathcal{F}' and no (G.12) violation ensues. So assume a is in row r .

If the reverse swap moves \mathcal{F} , then it cannot also move \mathcal{F}' ; hence T has \mathcal{F} southwest of \mathcal{F}' and no (G.12) violation ensues. Thus assume T has $\mathcal{F} \in a$. To violate (G.12), the reverse swap must move \mathcal{F}' South. So in T , $\mathcal{F}' \in \underline{b}$, $\mathcal{F}' \in \underline{b^\rightarrow}$ or $\mathcal{F}' \in b^\downarrow$.

Subcase 4.1: (T has $\mathcal{F}' \in \underline{b}$): Here $\mathcal{F}' = \mathcal{G}$ and the reverse miniswap involving b is L2.2 or L4.2. Although T has \mathcal{F} NorthWest of \mathcal{F}' , T has $\bullet_{\mathcal{G}} \in b$ to avoid violating (G.12). (We avoid violating the last sentence of T 's (G.12) by U 's (G.3) in the L2.2 case and by T 's $\bullet_{\mathcal{G}} \in b$ in the L4.2 case.)

Subcase 4.2: (T has $\mathcal{F}' \in \underline{b^\rightarrow}$): Here U has $\bullet_{\mathcal{G}^+} \in b^\rightarrow$ and the reverse miniswap involving b^\rightarrow is L4.4 or L4.5. Then T has $\bullet_{\mathcal{G}} \in b$, so although T has \mathcal{F} NorthWest of \mathcal{F}' , they do not violate (G.12).

Subcase 4.3: (T has $\mathcal{F}' \in b^\downarrow$): Here U has $\bullet_{\mathcal{G}^+} \in b^\downarrow$ and $\mathcal{F}' = \mathcal{G}$. Hence by U 's (G.2), $\text{label}_U(a^\downarrow)$ is a genetic label. Since a^\downarrow is northWest of $a \bullet_{\mathcal{G}^+}$, $\text{label}_U(a^\downarrow)$ is not marked. Hence by U 's (G.4), $\mathcal{F} < \text{label}_U(a^\downarrow)$. But by U 's (G.9), $\text{label}_U(a^\downarrow) \prec \mathcal{G}^+$, a contradiction.

Case 5: (U has $\mathcal{F} \in \underline{a}$ southwest of $\mathcal{F}' \in b$): Say b is in row r . No reverse miniswap can move \mathcal{F} further North than a or \mathcal{F}' further South than b^\downarrow . Therefore unless a is in r , T has \mathcal{F} southwest of \mathcal{F}' and no (G.12) violation ensues. Hence assume a is in r .

Subcase 5.1: (The reverse swap moves \mathcal{F} North): Here U has $\bullet_{\mathcal{G}^+} \in a$; the reverse miniswap involving a is L4.1, L4.2 or L4.3; and T has $\mathcal{F} \in a$. By U 's (G.2), \mathcal{F}' takes part in no reverse miniswap, so T has $\mathcal{F}' \in b$. Hence T has \mathcal{F} southwest of \mathcal{F}' and no (G.12) violation ensues.

Subcase 5.2: (T has $\mathcal{F} \in \underline{a}$): To have \mathcal{F} NorthWest of \mathcal{F}' in T , the reverse swap must move \mathcal{F}' to b^\downarrow . Hence U has $\bullet_{\mathcal{G}^+} \in b^\downarrow$ and $\mathcal{F}' = \mathcal{G}$. Since U has $\mathcal{F} \in \underline{a}$ and $\bullet_{\mathcal{G}^+} \in b^\downarrow$, it has a label in a^\downarrow . By U 's (G.2), it is a genetic label \mathcal{H} . By U 's (G.4), $\mathcal{F} < \mathcal{H}$. Hence since $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$, $\mathcal{G} < \mathcal{H}$. But by U 's (G.9), $\mathcal{H} \prec \mathcal{G}^+$, a contradiction.

Case 6: (U has $\mathcal{F} \in a$ southwest of $\mathcal{F}' \in b$): Say a is in row r . No swap can move \mathcal{F} North. No swap can move \mathcal{F}' further South than $\underline{b^\downarrow}$. Hence unless $b \in r^\uparrow$, T has \mathcal{F} southwest of

\mathcal{F}' and no (G.12) violation ensues. Thus, assume $b \in r^\uparrow$. To obtain a (G.12) violation, \mathcal{F}' must move South to $\underline{b^\downarrow}$ or $\underline{b^{\downarrow\rightarrow}}$.

Subcase 6.1: (T has $\mathcal{F}' \in \underline{b^\downarrow}$): Here U has $\bullet_{\mathcal{G}^+} \in b^\downarrow$, T has $\mathcal{F} \in a$ and $\mathcal{F}' = \mathcal{G}$. So T has $\bullet_{\mathcal{G}} \in b^\downarrow$. It remains to show we do not violate the last sentence of (G.12). By U 's (G.12), $\text{family}(\text{label}_U(b^{\downarrow\rightarrow})) \neq \text{family}(\mathcal{G})$; hence the same is true in T . Hence if $\text{label}_T(b^{\downarrow\rightarrow})$ is marked, then $\text{label}_U(b^{\downarrow\rightarrow})$ is marked. Then by Lemma 5.5, U has a label on $\underline{b^{\downarrow\rightarrow}}$ of the same family as \mathcal{G} , contradicting U 's (G.12). Thus (G.12) is confirmed in T .

Subcase 6.2: (T has $\mathcal{F}' \in \underline{b^{\downarrow\rightarrow}}$): Here U has $\bullet_{\mathcal{G}^+} \in b^{\downarrow\rightarrow}$ and $\mathcal{G} \in b^\downarrow$, while T has $\mathcal{F} \in a$ and $\bullet_{\mathcal{G}} \in b^\downarrow$. Thus, although T has \mathcal{F} NorthWest of \mathcal{F}' , they do not violate (G.12).

Case 7: (U has $\mathcal{F} \in \underline{a}$ southwest of $\mathcal{F}' \in \underline{b}$): Say b is in row r . No swap can move \mathcal{F} further North than a . No swap can move \mathcal{F}' further South than $\underline{b^\downarrow}$. Hence if a is South of row r^\downarrow , then T has \mathcal{F} southwest of \mathcal{F}' and no (G.12) violation ensues.

Case 7.1: (a is in row r^\downarrow): T has \mathcal{F} southwest of \mathcal{F}' , unless T has both $\mathcal{F} \in a$ and \mathcal{F}' in $\underline{b^\downarrow}$ or $\mathcal{F}' \in \underline{b^{\downarrow\rightarrow}}$. Suppose these both occur. Then U has $\bullet_{\mathcal{G}^+} \in a$ and either $\bullet_{\mathcal{G}^+} \in b^\downarrow$ or $\bullet_{\mathcal{G}^+} \in b^{\downarrow\rightarrow}$. Since by U 's (G.4), a is West of b , this contradicts U 's (G.2).

Case 7.2: (a is in row r): Suppose T has both $\mathcal{F} \in a$ and \mathcal{F}' in $\underline{b^\downarrow}$, $\mathcal{F}' \in \underline{b^{\downarrow\rightarrow}}$ or $\mathcal{F}' \in \underline{b^{\downarrow\rightarrow}}$. Then U has $\bullet_{\mathcal{G}^+} \in a$ and either $\bullet_{\mathcal{G}^+} \in b^\downarrow$ or $\bullet_{\mathcal{G}^+} \in b^{\downarrow\rightarrow}$, contradicting U 's (G.2). Hence the reverse swap cannot both move \mathcal{F} North and move \mathcal{F}' South.

Suppose T has $\mathcal{F} \in a$. Then T has $\mathcal{F}' \in \underline{b}$ and U has $\bullet_{\mathcal{G}^+} \in a$. Then $\mathcal{F} = \mathcal{G}^+$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G}) = \text{family}(\mathcal{G}^+)$. The reverse miniswap involving a is L4.1, L4.2 or L4.3. Hence U has $\mathcal{G} \in a^\leftarrow$. By U 's (G.4), $\text{label}_U(b) < \mathcal{F}'$, so it is marked. By Lemma 5.4, $\text{label}_U(a^\rightarrow)$ is marked. This contradicts the last sentence of U 's (G.12).

Suppose T has \mathcal{F}' in $\underline{b^\downarrow}$, $\underline{b^{\downarrow\rightarrow}}$ or $\underline{b^{\downarrow\rightarrow}}$. Then T has $\mathcal{F} \in \underline{a}$ and $\bullet_{\mathcal{G}} \in b^\downarrow$, while U has $\bullet_{\mathcal{G}^+} \in b^\downarrow$ or $b^{\downarrow\rightarrow}$. Hence although T has \mathcal{F} NorthWest of \mathcal{F}' , they do not violate (G.12).

Consistency of the prescribed virtual labels:

Lemma B.4. *Let U be a \mathcal{G}^+ -good tableau in which we apply L2.3, L4.3 or L4.5, and let $T \in \text{revswap}_{\mathcal{G}^+}(U)$. Then all prescribed \mathcal{G} 's from the outputs of L2.3, L4.3 and L4.5 are valid virtual labels in the sense of (V.1)–(V.3).*

Proof. Consider such a reverse miniswap. We may assume T contains an output with a prescribed \mathcal{G} , say in \underline{x} . Let T^* be T with \mathcal{G} added in \underline{x} .

((V.1) holds, i.e., $\mathcal{G} \in \underline{x}$ is not marked in T^*): Every \bullet in T^* is $\bullet_{\mathcal{G}}$, so by definition no \mathcal{G} in T^* is marked.

((V.2) holds, i.e., T has a \mathcal{G} West of x): By assumption of the reverse miniswaps, U has some \mathcal{G} West of x . Hence by Lemma B.3, T also has a \mathcal{G} West of x .

((G.1) holds in T^*): Immediate from U 's (G.1).

((G.4) holds in T^*): Let $\mathcal{H} = \text{label}_T(x^\downarrow)$. We must show $\mathcal{G} < \mathcal{H}$. By Lemma 8.4, x^\downarrow is not in a ladder of U , so $\mathcal{H} = \text{label}_U(x^\downarrow)$. Hence by U 's (G.4), we get $\mathcal{G} < \mathcal{H}$ in the case of L2.3 and L4.3. In the case of L4.5, by U 's (G.11), $\text{family}(\mathcal{H}) \geq \text{family}(\mathcal{G})$. But by U 's (G.12), $\text{family}(\mathcal{H}) \neq \text{family}(\mathcal{G})$, so $\mathcal{G} < \mathcal{H}$.

Let $\mathcal{F} = \text{label}_T(x^\uparrow)$. We must show $\mathcal{F} < \mathcal{G}$ in the L2.3 and L4.3 cases. (In the L4.5 case, there is nothing to confirm here.) Since x^\uparrow is not part of a ladder in U , $\mathcal{F} = \text{label}_U(x^\uparrow)$ as well. Hence $\mathcal{F} < \mathcal{G}$ follows from U 's (G.4).

((G.5) holds in T^*): Suppose not. Then there is some $\mathcal{G}' \in \underline{x}$ in T with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$. This \mathcal{G}' is not the result of any reverse miniswap, so $\mathcal{G}' \in \underline{x}$ in U . In the case of L2.3 or L4.3, this contradicts U 's (G.4). In the case of L4.5, note that by U 's (G.6) and (G.7), $\mathcal{G}' = \mathcal{G}^+$; this contradicts the assumption of L4.5.

((G.6) holds in T^*): This follows from Lemma B.3, as in the proof of T 's (G.6) above.

((G.8) holds in T^*): In the case of L4.5, this is immediate from the assumption that $N_{\mathcal{G}} = N_{\mathcal{E}}$ for every $\mathcal{E} \in Z$. In the case of L2.3, consider the tableau \tilde{T} differing from T only in the box x , where we choose the other output of L2.3. By \tilde{T} 's (G.8), \tilde{T} is ballot. But \tilde{T} and T^* have the same column reading words. Hence T^* is ballot.

Finally consider the case of L4.3. By T 's (G.8), T is ballot. Hence if there is a genotype G of T^* that is not ballot, G uses $\mathcal{G} \in \underline{x}$. Let \mathcal{F} be the gene with $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G}) - 1$ and $N_{\mathcal{F}} = N_{\mathcal{G}}$. Since G is not ballot, \mathcal{F} appears after \mathcal{G} in $\text{word}(G)$. Say \mathcal{F} appears in column c in G . By inspection of the reverse miniswaps, \mathcal{F} appears in U either in column c or column c^{\leftarrow} . Thus considering this \mathcal{F} and U 's $\mathcal{G} \in \underline{x}$, we contradict Lemma 5.6 for U .

((G.9) holds in T^*): \underline{x} is southeast of a $\bullet_{\mathcal{G}}$ in T . Hence by T 's (G.2), \underline{x} is not northwest of a $\bullet_{\mathcal{G}}$ in T , and this (G.9) verification is vacuous.

((G.12) holds in T^*): For L2.3, consider the tableau \tilde{T} differing from T only in the box x , where we choose the other output of L2.3. By \tilde{T} 's (G.12), any label \mathcal{G}' of \tilde{T} with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$ that is West of x must be no further North than the upper edge of x 's row. Since T^* and \tilde{T} are identical outside of x , this is also true of T^* . Since T^* has $\bullet_{\mathcal{G}} \in x$ and $\mathcal{G} \notin x^{\leftarrow}$, T^* 's (G.12) then follows.

For L4.3, observe that in light of the $\mathcal{G}^+ \in x^{\rightarrow}$ in T , it follows from T 's (G.12) that any label \mathcal{G}' of T with $\text{family}(\mathcal{G}') = \text{family}(\mathcal{G})$ that is West of x must be no further North than the upper edge of x 's row. Since T^* has $\bullet_{\mathcal{G}} \in x$ and $\mathcal{G} \notin x^{\leftarrow}$, T^* 's (G.12) then follows.

The L4.5 case is proved by repeating the arguments above for T 's (G.12) (Cases 3–5). \square

(G.13): For every marked label $\mathcal{E}^!$ in T , U has \mathcal{E} or $\mathcal{E}^!$ in the same position. Thus our analysis splits into two cases:

Case 1: (U has $\mathcal{E}^! \in \underline{b}$ or \underline{b}): Let ℓ be this instance of $\mathcal{E}^!$ and let \underline{b} be in column c . U also has some \mathcal{F} or \mathcal{F} in \underline{b} with $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$. Since ℓ is marked, $\mathcal{E} \preceq \mathcal{G}$.

Subcase 1.1: ($\text{family}(\mathcal{E}) \leq \text{family}(\mathcal{G}) - 2$): Such a marked label is not moved by any reverse miniswap. Hence, ℓ is in the same position in U and T . By Lemma 5.5, U has $\mathcal{F} \in \underline{b}$ (rather than \mathcal{F}). This \mathcal{F} is also not moved by any reverse miniswap, so T has $\mathcal{F} \in \underline{b}$ as well, and (G.13) is satisfied.

Subcase 1.2: ($\text{family}(\mathcal{E}) = \text{family}(\mathcal{G}) - 1$): Here, $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$. As in Subcase 1.1, ℓ is in the same position in U and T . Suppose U has $\mathcal{F} \in \underline{b}$ (rather than \mathcal{F}). By U 's (G.11) $\bullet_{\mathcal{G}^+} \notin c$ in U . Hence there is no reverse miniswap that affects $\mathcal{F} \in \underline{b}$. So T has $\mathcal{F} \in \underline{b}$ and (G.13) holds.

Hence, assume \mathcal{F} in \underline{b} in U . By Lemma 5.9, $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G}^+)$ and U has $\bullet_{\mathcal{G}^+} \in \underline{b}^{\leftarrow}$. Let T^* be T with $\mathcal{F} \in \underline{b}$. By U 's (V.1), $\mathcal{G} \prec \mathcal{F}$. Hence T^* satisfies (V.1). By the proof of Lemma 9.4, it satisfies (V.2). It remains to show that T^* satisfies (G.1), (G.4)–(G.6), (G.8), (G.9) and (G.12).

(G.1): Since U has $(\mathcal{F}) \in \underline{b}$, this follows from T 's (G.1).

(G.4): By T 's (G.4), since $\mathcal{E} < \mathcal{F}$, T has $\mathcal{Q} < \mathcal{F}$ for any \mathcal{Q} north of \underline{b} in c . By U 's (G.4), since no reverse miniswap affects labels of family greater than $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G})$, we also have $\mathcal{Q} > \mathcal{F}$ for any \mathcal{Q} appearing south of \underline{b} in c in T . Hence (G.4) holds in T^* .

(G.5): By U 's (G.5), U has no label of family $\text{family}(\mathcal{F})$ on \underline{b} . Indeed by U 's (G.4), U has no label of that family in c . By U 's (G.11), U has $\bullet_{\mathcal{G}^+} \notin c$. Hence no ladder of U intersects c and T has no label of family $\text{family}(\mathcal{F})$ on \underline{b} . Hence (G.5) holds in T^* .

(G.6): As no ladder of U intersects c , the genes West of c in U are exactly the genes West of c in T , and the genes East of c in T are a subset of those East of c in U . Hence (G.6) holds in T^* .

(G.8): As no ladder of U intersects c , c is the same in U and T . Since $N_{\mathcal{E}} = N_{\mathcal{F}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$, it suffices to check no \mathcal{E} is read in T^* after $\mathcal{F} \in \underline{b}$. By U 's (G.4), there is no \mathcal{E} in U South of \underline{b} in c . Thus, this is also true in T . By Lemma 5.7, ℓ is the westmost \mathcal{E} in U . But the genes that appear West of c in U are exactly the genes that appear West of c in T . Hence no \mathcal{E} appears West of \underline{b} in T , no \mathcal{E} is read in T^* after \underline{b} and T^* 's (G.8) holds.

(G.9): Since U has $\bullet_{\mathcal{G}^+} \in \mathbf{b}^{\leftarrow}$, by U 's (G.2), \underline{b} is not northwest of a $\bullet_{\mathcal{G}^+}$. Hence by inspection of the reverse miniswaps, \underline{b} is not northwest of a $\bullet_{\mathcal{G}}$ in T , so (G.9) holds in T^* .

(G.12): Suppose i is a label of \mathcal{F} 's family appearing in \underline{a} or \underline{a} NorthWest of \underline{b} in T . First suppose i does not appear in the same position in U . Then i was involved in a reverse miniswap. If it was L1.1, then $\mathcal{G} \in \mathbf{a}^{\uparrow}$ in U , contradicting U 's (G.12). It obviously was not L1.2. If it was an L2 miniswap, $i \in \underline{a}$ in U , contradicting U 's (G.12). The same holds for L3. For L4.1–3 to apply, $\underline{a} = \mathbf{b}^{\leftarrow}$ by U 's (G.2), but this contradicts U 's (G.12) (last sentence). In L4.4 and L4.5, $Z \neq \emptyset$ and the only labels of concern are the \mathcal{G} 's. Since they satisfy (G.12) in U , they do in T^* .

Now, suppose that i is in the same position in U and T . Hence i is NorthWest of \underline{b} in U . Let \underline{b} be in row r . By U 's (G.12), either U has $i \in \underline{a}$ and $\underline{a} \in r$, or else U has $i \in \underline{a}$ and $\underline{a} \in r^{\uparrow}$. Also \underline{a} is West of \mathbf{b}^{\leftarrow} . Since the labels in question do not move, it remains to check that $\bullet_{\mathcal{G}}$ appears in row r in T East of \underline{a} and west of \underline{b} . We are clearly only concerned when the reverse miniswap involving \mathbf{b}^{\leftarrow} is L1.1, or is L4 with $\mathbf{b}^{\leftarrow} = \mathbf{a}^{\rightarrow}$. If it is L1.1, U has $\mathcal{G} \in \mathbf{b}^{\leftarrow\uparrow}$, contradicting U 's (G.12). If it is L4 with $\mathbf{b}^{\leftarrow} = \mathbf{a}^{\rightarrow}$, then i would not appear in the same position in T , contradicting our assumption.

Now, suppose i is a label of \mathcal{F} 's family appearing in \underline{a} or \underline{a} SouthEast of \underline{b} in T . First, suppose that i does not appear in the same position in U . Then i was involved in a reverse miniswap. By U 's (G.2), this can only be a L2 reverse miniswap. It is obviously not L2.1. If it is L2.2 or L2.3, then by U 's (G.12), the \mathcal{G} or $(\mathcal{G}) \in \underline{a}$ is not a (G.12) violation in T because of T 's $\bullet_{\mathcal{G}} \in \underline{a}$.

Otherwise i is in the same position in U and T . Hence i is SouthEast of \underline{b} in U . By U 's (G.12), U has a $\bullet_{\mathcal{G}^+}$ SouthEast of \underline{b} . Given U 's $\bullet_{\mathcal{G}^+} \in \mathbf{b}^{\leftarrow}$, this contradicts U 's (G.2).

We conclude that the desired (\mathcal{F}) appears on \underline{b} in T .

Subcase 1.3: ($\text{family}(\mathcal{E}) = \text{family}(\mathcal{G})$): Suppose ℓ is moved by the reverse swap. Recall $\mathcal{E} \prec \mathcal{G}^+$. By inspection, no reverse swap will move such a $\mathcal{E} \prec \mathcal{G}^+$ with $\text{family}(\mathcal{E}) = \text{family}(\mathcal{G})$ unless $\mathcal{E} = \mathcal{G}$. Since the \bullet 's in T are $\bullet_{\mathcal{G}}$'s, no \mathcal{E} will be marked in T , so T 's (G.13) check is vacuous.

Hence assume that ℓ is unaffected by $\text{revswap}_{\mathcal{G}^+}$ and indeed that $\mathcal{E} \prec \mathcal{G}$. Since $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G}) + 1$, no reverse miniswap affects any instance of \mathcal{F} . In particular, if U has $\mathcal{F} \in \underline{b}$ (instead of \mathcal{F}), then T will also have $\mathcal{F} \in \underline{b}$ and satisfy (G.13).

Hence further assume that U has $\mathcal{F} \in \underline{b}$. We need $\mathcal{F} \in \underline{b}$ in T . By Lemma 5.9, U has $\bullet_{\mathcal{G}^+} \in \mathfrak{b}^{\leftarrow}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{G}^+)$. Let c be \mathfrak{b} 's column. By U 's (G.11), U has $\bullet_{\mathcal{G}^+} \notin c$. By U 's (G.4), since $\mathcal{E} \prec \mathcal{G}$, U has $\mathcal{G} \notin c$. Hence no ladder of U intersects c and so column c in T is identical to column c in U .

Let T^*, U^* be T, U respectively with $\mathcal{F} \in \underline{b}$ added. We show T^* satisfies (V.1)–(V.3).

(V.1): Since $\mathcal{F} > \mathcal{G}$, this is obvious.

(V.2): By U 's (V.2), U has an \mathcal{F} West of c . Hence there is an \mathcal{F} West of c in T , as needed.

(V.3): We show T^* satisfies (G.1), (G.4), (G.5), (G.6), (G.8), (G.9) and (G.12). We know T satisfies these.

(G.1): Immediate from T 's (G.1), given U 's $\mathcal{F} \in \underline{b}$.

(G.4) and (G.5): These hold in T^* since they hold in U^* , and column c of T is identical to column c of U .

(G.6): The genes West of c in U are exactly the genes West of c in T , and the genes East of c in T are a subset of those East of c in U . Now T^* 's (G.6) follows.

(G.8): Immediate from U 's $\mathcal{F} \in \underline{b}$ and the facts that

- the genes West of c in T are exactly the genes West of c in U ,
- the genes East of c in T are a subset of those East of c in U , and
- column c in T is identical to column c in U .

(G.9): Since ℓ is marked in T , T has a $\bullet_{\mathcal{G}}$ northwest of \underline{b} . Hence by T 's (G.2), \underline{b} is not northwest of a $\bullet_{\mathcal{G}}$ in T^* , so this condition is vacuous.

(G.12): Take \mathcal{F}' with $\text{family}(\mathcal{F}') = \text{family}(\mathcal{F})$. Suppose \mathcal{F}' is NorthWest of \underline{b} in T^* . Since $\text{family}(\mathcal{F}') > \text{family}(\mathcal{G})$, \mathcal{F}' appears in the same positions in both T^* and U^* . Hence \mathcal{F}' is NorthWest of \underline{b} in U^* . But then this \mathcal{F}' is northwest of U^* 's $\bullet_{\mathcal{G}^+} \in \mathfrak{b}^{\leftarrow}$, contradicting U^* 's (G.9). Thus T^* has no such \mathcal{F}' NorthWest of \underline{b} . Similarly T^* has no \mathcal{F}' SouthEast of \underline{b} .

Case 2: (ℓ is a marked label in T that is not marked in U): Suppose ℓ is an instance of \mathcal{E}^1 on \mathfrak{b} or \underline{b} in T . Since ℓ is marked and every bullet in T is $\bullet_{\mathcal{G}}$, $\mathcal{E} \prec \mathcal{G}$. Hence $\mathcal{E} \prec \mathcal{G}^+$, and any instance of \mathcal{E} southeast of a $\bullet_{\mathcal{G}^+}$ is marked in U .

Case 2.1: ($\text{family}(\mathcal{E}) = \text{family}(\mathcal{G})$): No reverse miniswap affects any instance of \mathcal{E} . Hence ℓ is in the same position in U as in T . Since ℓ is unmarked in U , U has no $\bullet_{\mathcal{G}^+}$ northwest of \underline{b} . Hence, since U has $\mathcal{E} \in \mathfrak{b}$ or \underline{b} , by U 's (G.3) and (G.4), U has no \mathcal{G} northwest of \underline{b} . But there is $\bullet_{\mathcal{G}}$ northwest of \underline{b} in T ; this contradicts Lemma B.1.

Case 2.2: ($\text{family}(\mathcal{E}) < \text{family}(\mathcal{G})$): If ℓ is not moved by $\text{revswap}_{\mathcal{G}^+}$, we obtain a contradiction exactly as in Case 2.1. Otherwise, it is moved by a L4.4 or L4.5 reverse miniswap. Then by definition $N_{\mathcal{E}} = N_{\mathcal{G}}$. Let \mathcal{F} be the gene (which must exist) with $N_{\mathcal{F}} = N_{\mathcal{E}} = N_{\mathcal{G}}$ and $\text{family}(\mathcal{F}) = \text{family}(\mathcal{E}) + 1$. By Lemma 5.8, \mathcal{F} appears in $\mathfrak{b}^{\leftarrow}$'s column in U . Hence by U 's (G.4), U has $\mathcal{F} \in \mathfrak{b}^{\leftarrow}$ or $\overline{\mathfrak{b}^{\leftarrow}}$. It is then in the set A'' or $A''' \cup \{\mathcal{G}\}$ (in the notation of L4.4/L4.5) and appears on \underline{b} in T (possibly virtual), as desired. \square

APPENDIX C. BLOCK DECOMPOSITION; COMPLETION OF PROOF OF
PROPOSITION 12.3(II)

Below, we define, for each $S \in \text{Snakes}_{\mathcal{G}}$, a set \mathcal{B}_S containing S . Clearly $\cup_{S \in \text{Snakes}_{\mathcal{G}}} \mathcal{B}_S = \text{Snakes}_{\mathcal{G}}$. Along the way, we will argue that if $S, S' \in \text{Snakes}_{\mathcal{G}}$ and $S' \in \mathcal{B}_S$, then $\mathcal{B}_S = \mathcal{B}_{S'}$. This proves (D.1).

Recall $\Gamma_i := \{T \in P_{\mathcal{G}} : T \text{ contains a snake from } \mathcal{B}_i\}$. From the construction, the following two additional conditions will also be essentially clear:

(D.3) Suppose $A, B \in \Gamma_i$. For each snake in \mathcal{B}_i and A , there is a snake of \mathcal{B}_i in B and in the exact same location.

(D.4) $A, B \in \Gamma_i$ are identical outside of the snakes in \mathcal{B}_i .

The bulk of the work is to establish (D.2). This will be done simultaneously with the description of each \mathcal{B}_S . To establish (D.2), we must verify (12.3) by considering the boxfactors, edg factors and virtual factors from every box and edge of the common shape ν/λ . Except where otherwise noted, by inspection, these factors do not change for boxes/edges not in \mathcal{B}_S . Thus, the majority of our discussion concerns the region defined by \mathcal{B}_S . For simplicity, we assume $\mathcal{B}_J = \emptyset$. The modifications for the general case are straightforward, using (D.3) and (D.4).

Assume $\mathcal{G} := i_k$. Let $S \in \text{Snakes}_{\mathcal{G}}$ be in the tableau U . We break into cases according to the type of $\text{head}(S)$. We write T_1, T_2, \dots for the fine tableau in $\text{swap}_S(U)$, in the order illustrated in each case. We write U_j for T_j together with its coefficient in $\text{swap}_S(U)$.

Case 0: ($\text{head}(S) = \emptyset$): By Definition-Lemma 6.8, either the southmost row of S contains a single box, or else it consists of two boxes x, x^{\rightarrow} with $\bullet \in x$ and a marked label in x^{\rightarrow} . Thus, $\text{body}(S)$ is either empty, or it falls under case B1 or B3.

Subcase 0.1: ($\text{body}(S)$ is B1): Let $\mathcal{B}_S = \{S\}$. Since S contains no $\bullet_{\mathcal{G}}$, $\text{swapset}_{\mathcal{B}_S}(U) = U$. Thus $\text{wt}(\text{swapset}_{\mathcal{B}_S}(U)) = \text{wt}(U)$, which implies (12.3).

Subcase 0.2: ($\text{body}(S)$ is B3 or $\text{body}(S) = \emptyset$): By Definition-Lemma 6.8(III) either S has at least two rows, or else $S = \text{tail}(S)$. In either case, $\text{tail}(S) \neq \emptyset$.

Subcase 0.2.1: ($\text{tail}(S)$ is T1): Let $\mathcal{B}_S = \{S\}$. Locally at the snake S , this swap looks like $\begin{array}{|c|c|} \hline \bullet & \mathcal{G} \\ \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \\ \hline \end{array} \mapsto \prod_{x \ni \bullet_{\mathcal{G}}} \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{G} & \\ \hline \mathcal{G} & \bullet \\ \hline \bullet & \\ \hline \end{array}$. Note that $\text{body}(S)$ is nonempty in this case. This swap

does not affect the locations or weights of edge labels or virtual labels in U . Hence $\text{edgewt}(U) = \text{edgewt}(T_1)$ and $\text{virtualwt}(U) = \text{virtualwt}(T_1)$. One checks that a box outside S is productive in U if and only if it is productive in T_1 . (The critical checks are for the box immediately east of the northmost box of S and the box immediately west of the southmost box of S .) Also, each such productive box has the same boxfactor in U and T_1 . The productive boxes of S_1 are the boxes $\{x\}$ containing \mathcal{G} , while in S they are the boxes $\{x^{\downarrow}\}$ containing \mathcal{G} . For each productive box x of S_1 with $\text{boxfactor}(x) := w_x$, there is a corresponding productive box x^{\downarrow} in S with $\text{boxfactor}(x^{\downarrow}) = \hat{\beta}(x)w_x$.

Thus $\text{wt } U = \text{wt } U_1$ follows from

$$(-1)^{d(U)} \prod_{x: \text{label}_U(x) = \bullet_{\mathcal{G}}} \hat{\beta}(x)w_x = \left(\prod_{x: \text{label}_U(x) = \bullet_{\mathcal{G}}} \hat{\beta}(x) \right) \cdot (-1)^{d(U)} \prod_{x: \text{label}_U(x) = \bullet_{\mathcal{G}}} w_x.$$

Subcase 0.2.2: ($\text{tail}(S)$ is T2): Let $\mathcal{B}_S = \{S\}$. This case is similar to Subcase 0.2.1; we have something like

$$\begin{array}{|c|c|} \hline \bullet & \mathcal{G} \\ \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \\ \hline \end{array} \mapsto - \prod_{x \ni \bullet} \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{G} & \bullet \\ \hline \mathcal{G} & \bullet \\ \hline \bullet & \\ \hline \end{array}.$$

This swap preserves locations and weights of edge labels. Hence $\text{edgewt}(U) = \text{edgewt}(T_1)$. A box outside S is productive in U if and only if it is productive in T_1 . Also, each such productive box has the same boxfactor in U and T_1 . Let y be the box containing the Northmost \mathcal{G} in S . In S_1 , the productive boxes are the boxes $\{x\}$ containing \mathcal{G} . The productive boxes of S are the boxes $\{x^\downarrow\}$ containing \mathcal{G} in all but the northmost row of S , and y if $\mathcal{G}^+ \notin y^\rightarrow$ with $\text{family}(\mathcal{G}) = \text{family}(\mathcal{G}^+)$. For each productive box x in S_1 with $\text{boxfactor}(x) := w_x$, there is a corresponding productive x^\downarrow of S with $\text{boxfactor}(x^\downarrow) = \hat{\beta}(x)w_x$.

The box y is productive in S with $\text{boxfactor}(y) := y$ if and only if $(\mathcal{G}) \in \bar{y}$ in S_1 with $\text{virtualfactor}_{\mathcal{G}}(\bar{y}) = y$. The swap does not otherwise affect the location or weight contribution of virtual labels. Finally note $(-1)^{d(U)} = (-1)^{d(T_1)-1}$. If y is productive in S , then $\text{wt } U = \text{wt } U_1$ follows from

$$(-1)^{d(U)} \cdot y \prod_{x:\text{label}_U(x)=\bullet\mathcal{G}} \hat{\beta}(x)w_x = \left(- \prod_{x:\text{label}_U(x)=\bullet\mathcal{G}} \hat{\beta}(x) \right) \cdot (-1)^{d(U)-1} \cdot y \cdot \prod_{x:\text{label}_U(x)=\bullet\mathcal{G}} w_x.$$

If y is not productive in S , we use the same identity without y .

Subcase 0.2.3: ($\text{tail}(S)$ is T3): This is again similar to Subcase 0.2.1. Locally at S , we have something like

$$\begin{array}{|c|c|} \hline \bullet & \mathcal{G}^+ \\ \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \\ \hline \end{array} \mapsto \prod_{x \ni \bullet} \hat{\beta}(x) \cdot \begin{array}{|c|c|} \hline \mathcal{G} & \mathcal{G}^+ \\ \hline \mathcal{G} & \bullet \\ \hline \bullet & \\ \hline \end{array} - \alpha \cdot \begin{array}{|c|c|} \hline \mathcal{G} & \bullet \\ \hline \mathcal{G} & \mathcal{G}^\pm \\ \hline \bullet & \\ \hline \end{array}.$$

By Lemma 6.9(V), S has at least two rows. Let y be the box containing \mathcal{G}^+ in S . Here $\alpha := \prod_{x \ni \bullet} \hat{\beta}(x)$ if $\bullet \notin y^\uparrow$ and $\alpha := 0$ otherwise.

The locations and weights of virtual labels are unaffected by the swap. Therefore, $\text{virtualwt}(U) = \text{virtualwt}(T_1) = \text{virtualwt}(T_2)$. Furthermore the edgefactors and boxfactors from labels outside of S are the same in each of U, T_1, T_2 , so we restrict attention to the boxfactors and edgefactors from labels inside S .

Subcase 0.2.3.1: ($\bullet_{\mathcal{G}} \notin y^\uparrow$): Let $\mathcal{B}_S = \{S\}$. The productive boxes of S_1 and S_2 are the boxes $\{x\}$ containing \mathcal{G} and not in the northmost row, and possibly also y (depending on what label, if any, appears in y^\rightarrow). The productive boxes of S are those boxes $\{x^\downarrow\}$ containing \mathcal{G} not in the second row from the top, the box z containing \mathcal{G} in the second row from the top, and possibly also y . One sees y is productive in any of S, S_1, S_2 if and only if it is productive in all of them. Further, if it is productive, then it has the same boxfactor, say a , in each one.

For each productive box x in each of S_1, S_2 , with $\text{boxfactor}(x) := w_x$, there is a corresponding productive box x^\downarrow in S , with $\text{boxfactor}(x^\downarrow) = \hat{\beta}(x)w_x$. Let $\text{edgefactor}_{\underline{z} \in T_2}(\mathcal{G}^+) := 1 - b$. Then $\text{boxfactor}_U(z) = \hat{\beta}(z^\uparrow)b$.

Now (12.3) is the statement $\text{wt } U = \text{wt}(U_1 + U_2)$. If y is productive in U , this follows from the identity

$$a\hat{\beta}(z^\uparrow)b \prod_x \hat{\beta}(x)w_x = \left(\hat{\beta}(z^\uparrow) \prod_x \hat{\beta}(x) \right) \cdot \left[a \prod_x w_x - (1-b) \cdot a \prod_x w_x \right]$$

Otherwise we use the same identity without a .

Subcase 0.2.3.2: ($\bullet_{\mathcal{G}} \in y^\uparrow$): Here $\alpha = 0$, so we ignore T_2 . Let S' be the snake containing y^\uparrow and let $\mathcal{B}_S = \{S, S'\}$. By Lemma 6.6, $S' = \{y^\uparrow\}$ participates in a trivial H3 or H8 miniswap.

The productive boxes $\{x\}$ of S_1 are those containing \mathcal{G} (even the box y^\leftarrow) and possibly also y (depending on what label, if any, appears in y^\rightarrow). The productive boxes of S are the boxes $\{x^\downarrow\}$ containing \mathcal{G} and possibly also y . One checks that y is productive in S if and only if it is productive in S_1 . If productive, $\text{boxfactor}_U(y) = \text{boxfactor}_{T_1}(y) := y$. In S' , y^\uparrow is productive if and only if it is in S'_1 ; if it is productive, it contributes the same $\text{boxfactor}(y^\uparrow) := q$ to both.

For each productive x in S_1 with $\text{boxfactor}(x) := w_x$, there is a corresponding productive x^\downarrow in S with $\text{boxfactor}(x^\downarrow) = \hat{\beta}(x)w_x$. If y and y^\uparrow are productive in U , (12.3) follows from

$$qy \prod_{x:\text{label}_U(x)=\bullet_{\mathcal{G}}} \hat{\beta}(x)w_x = q \left(\prod_{x:\text{label}_U(x)=\bullet_{\mathcal{G}}} \hat{\beta}(x) \right) \cdot y \cdot \prod_{x:\text{label}_U(x)=\bullet_{\mathcal{G}}} w_x.$$

Otherwise we use the same identity without q, y or both.

Subcase 0.2.4: ($\text{tail}(S)$ is T4): Let $\text{tail}(S) = \{x, y := x^\rightarrow\}$. By (G.7), the $\mathcal{G} \in \underline{y}$ is westmost in its gene, so $\text{body}(S) = \emptyset$.

Subcase 0.2.4.1: ($\text{tail}(S)$ is T4.1): Set $\mathcal{B}_S = \{S\}$. Locally at S , $\boxed{\bullet_{\mathcal{G}, \mathcal{H}} \mathcal{F}^\dagger} \mapsto \boxed{\bullet_{\mathcal{G}^\dagger, \mathcal{H}^\dagger} \mathcal{F}^\dagger}$, where we have $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$. Virtual labels appear in the same places in U and T_1 . In particular, neither U nor T_1 can have $(\mathcal{G}) \in \underline{x}$, since it would be West of every \mathcal{G} . Further no labels move. As no weights change, trivially $\text{wt } U = \text{wt } U_1$, which implies (12.3).

Subcase 0.2.4.2: ($\text{tail}(S)$ is T4.2):

Subcase 0.2.4.2.1: ($\mathcal{G}^- \in x^\leftarrow$ with $\text{family}(\mathcal{G}^-) = \text{family}(\mathcal{G})$): Set $\mathcal{B}_S = \{S\}$. Locally at S , the swap is $\boxed{\mathcal{G}^- \bullet_{\mathcal{G}, \mathcal{H}} \mathcal{F}^\dagger} \mapsto \boxed{\mathcal{G}^- \bullet_{\mathcal{G}^\dagger, \mathcal{H}^\dagger} \mathcal{F}^\dagger} + \hat{\beta}(x) \cdot \boxed{\mathcal{G}^- \mathcal{G}^\dagger \bullet}$. Here (12.3) is equivalent to $\text{wt } U = \text{wt } T_1 + \hat{\beta}(x) \text{wt } T_2$.

The $\mathcal{F}^\dagger \in \underline{y}$ is productive in U and T_1 if and only if $(\mathcal{F}) \in \bar{y}$ in T_2 . If these boxes are productive,

$$\text{boxfactor}_U(y) = \text{boxfactor}_{T_1}(y) = \text{virtualfactor}_{\bar{y} \in T_2}((\mathcal{F})) := a.$$

The box x is not productive in U or T_1 , but it is in T_2 . Let $\text{boxfactor}_{T_2}(x) := u$. Then $\text{virtualfactor}_{\underline{y} \in U}((\mathcal{H})) = u$ and $\text{virtualfactor}_{\underline{y} \in T_1}((\mathcal{H})) = u - 1$. Let $w := \text{edgefactor}_{\underline{y} \in T}(\mathcal{G})$. Then we have $\text{edgefactor}_{\underline{y} \in T_1}(\mathcal{G}^\dagger) = w$ and $\text{edgefactor}_{\bar{x} \in T_2}(\mathcal{F}) = w$. The box x^\leftarrow is productive in U and T_1 , but not in T_2 . We have $\text{boxfactor}_U(x^\leftarrow) = \text{boxfactor}_{T_1}(x^\leftarrow) = \hat{\beta}(x)u$.

If y is productive in U , $\text{wt } U = \text{wt } T_1 + \hat{\beta}(x) \text{wt } T_2$ follows from the identity on \mathcal{B}_S -contributions: $\hat{\beta}(x)au^2w = \hat{\beta}(x)au(u-1)w + \hat{\beta}(x)auw$. If y is not productive in U , it follows from the same identity after cancelling a 's.

Subcase 0.2.4.2.2: ($\mathcal{G}^- \notin x^\leftarrow$ or $\mathcal{G}^- < \mathcal{G}$): Let $\bar{T} := \phi_4^{-1}(T) \in P_{\mathcal{G}}$. Let \bar{S} be the snake of \bar{T} containing x . Set $\mathcal{B}_S = \{S, \bar{S}\}$; thus $\Gamma = \{T, \bar{T}\}$.

Locally at S , the swap is $\boxed{\bullet \mid \mathcal{F}^!}_{\mathcal{G} \oplus \mathbb{Z}} \mapsto \boxed{\bullet \mid \mathcal{F}^!}_{\mathcal{G} \oplus \mathbb{Z}} + \hat{\beta}(x) \cdot \boxed{\mathcal{G} \mid \bullet}_{\mathcal{F} \cup \mathbb{Z}}$. Locally at \bar{S} , $\boxed{\bullet \mid \mathcal{F}^!}_{\mathcal{G} \oplus \mathbb{Z}} \mapsto 0$. By Proposition 12.21, $[T]P_{\mathcal{G}} = [\bar{T}]P_{\mathcal{G}}$. Hence (12.3) is equivalent to $\text{wt } U + \text{wt } \bar{U} = \text{wt } U_1 + \text{wt } U_2$. Here $U(\Gamma) = \{U, \bar{U}\}$.

The $\mathcal{F}^! \in y$ is productive in U, T_1 and \bar{U} if and only if $(\mathcal{F}) \in \bar{y}$ in T_2 . If these boxes are productive,

$$\text{boxfactor}_U(y) = \text{boxfactor}_{T_1}(y) = \text{boxfactor}_{\bar{U}}(y) = \text{virtualfactor}_{\bar{y} \in T_2}((\mathcal{F})) := a.$$

The box x is not productive in U, T_1 or \bar{U} . Let $w = \text{edgefactor}_{y \in T}(\mathcal{G})$. Then we have $\text{virtualfactor}_{y \in \bar{T}}((\mathcal{G})) = -w$, $\text{edgefactor}_{y \in T_1}(\mathcal{G}^!) = w$ and $\text{edgefactor}_{x \in T_2}(\mathcal{F}) = w$. Let $u = \text{virtualfactor}_{y \in U}(\mathcal{H})$. Then $\text{virtualfactor}_{y \in T_1}(\mathcal{H}) = u - 1$. Let $1 - v = \text{edgefactor}_{x \in \bar{U}}(\mathcal{G})$. Then $\text{boxfactor}_{T_2}(x) = \frac{v}{\hat{\beta}(x)}$.

If y is productive in U , $\text{wt } U + \text{wt } \bar{U} = \text{wt } U_1 + \text{wt } U_2$ follows from the identity on \mathcal{B}_S -contributions: $auw - a(1-v)w = a(u-1)w + a\hat{\beta}(x)\frac{v}{\hat{\beta}(x)}w$. The same is true if y is not productive in U except that a does not appear.

Subcase 0.2.4.3: ($\text{tail}(S)$ is T4.3):

Subcase 0.2.4.3.1: ($\mathcal{G}^- \in x^\leftarrow$ with $\text{family}(\mathcal{G}^-) = \text{family}(\mathcal{G})$): Set $\mathcal{B}_S = \{S\}$. Locally at S , the swap is $\boxed{\mathcal{G}^- \mid \bullet \mid \mathcal{F}^!}_{z \cup \mathcal{G}} \mapsto \hat{\beta}(x) \cdot \boxed{\mathcal{G}^- \mid \mathcal{G} \mid \bullet}_{\mathcal{F} \cup \mathbb{Z}}$. Here (12.3) is equivalent to $\text{wt } U = \hat{\beta}(x) \text{wt } T_1$.

For every label $\ell \in Z \cup \mathcal{G}$ in U , there is a unique label $\ell_1 \in \mathcal{F} \cup Z$ with $\text{family}(\ell_1) = \text{family}(\ell) - 1$. Also, $\text{edgefactor}_U(\ell) = \text{edgefactor}_{T_1}(\ell_1) := a_\ell$.

In U , y is productive if and only if y^\rightarrow does not contain a label of the same family as \mathcal{F} . Hence y is productive in U if and only if $(\mathcal{F}) \in \bar{y}$ in T_1 . If y is productive in U , then $\text{boxfactor}_U(y) = \text{boxfactor}_{\bar{U}}(y) = \text{virtualfactor}_{\bar{y} \in T_1}((\mathcal{F})) := b$.

The box x is productive in T_1 , but not U . Let $w := \text{boxfactor}_{T_1}(x)$. The box x^\leftarrow is productive in U , but not in T_1 . We have $\text{boxfactor}_U(x^\leftarrow) = \hat{\beta}(x)w$.

Hence if y is productive in U , $\text{wt } U = \hat{\beta}(x) \text{wt } T_1$ follows from

$$\hat{\beta}(x)wb \prod_{\ell \in Z \cup \mathcal{G}} a_\ell = \hat{\beta}(x)bw \prod_{\ell_1 \in \mathcal{F} \cup Z} a_\ell.$$

Otherwise, we use the same identity after canceling b .

Subcase 0.2.4.3.2: ($\mathcal{G}^- \notin x^\leftarrow$ or $\mathcal{G}^- < \mathcal{G}$): Let $\bar{T} := \phi_4^{-1}(T) \in P_{\mathcal{G}}$. Let \bar{S} be the snake of \bar{T} containing x and set $\mathcal{B}_S = \{S, \bar{S}\}$.

Locally at S , the swap is $\boxed{\bullet \mid \mathcal{F}^!}_{z \cup \mathcal{G}} \mapsto \hat{\beta}(x) \cdot \boxed{\mathcal{G} \mid \bullet}_{\mathcal{F} \cup \mathbb{Z}}$. Locally at \bar{S} , $\boxed{\bullet \mid \mathcal{F}^!}_{z \cup \mathcal{G}} \mapsto 0$. By Proposition 12.21, $[T]P_{\mathcal{G}} = [\bar{T}]P_{\mathcal{G}}$. Hence (12.3) is equivalent to $\text{wt } U + \text{wt } \bar{U} = \text{wt } U_1$.

For every label $\ell \in Z \cup \mathcal{G}$ in U , there is a unique label $\ell_1 \in \mathcal{F} \cup Z$ with $\text{family}(\ell_1) = \text{family}(\ell) - 1$. Also, $\text{edgefactor}_U(\ell) = \text{edgefactor}_{T_1}(\ell_1) := a_\ell$. If $\ell \in Z$ in U , then there is unique $\bar{\ell} \in Z$ in \bar{U} with $\text{family}(\ell) = \text{family}(\bar{\ell})$, and $\text{edgefactor}_{\bar{U}}(\bar{\ell}) = a_\ell$. Further $\text{virtualfactor}_{y \in \bar{U}}(\mathcal{G}) = -a_\mathcal{G}$.

In U , y is productive if and only if y^\rightarrow does not contain a label of the same family as \mathcal{F} . Hence y is productive in U if and only if it is productive in \bar{U} and further if and only if $(\mathcal{F}) \in \bar{y}$ in T_1 . If y is productive in U , then $\text{boxfactor}_U(y) = \text{boxfactor}_{\bar{U}}(y) = \text{virtualfactor}_{y \in T_1}(\mathcal{F}) := b$.

The box x is productive in T_1 . Let $w := \text{boxfactor}_{T_1}(x)$. Observe $\text{edgefactor}_{x \in \bar{U}}(\mathcal{G}) = 1 - \hat{\beta}(x)w$. Hence $\text{wt } U + \text{wt } \bar{U} = \text{wt } U_1$ follows from

$$b \prod_{\ell \in Z \cup \mathcal{G}} a_\ell + b(1 - \hat{\beta}(x)w)(-a_\mathcal{G}) \prod_{\ell \in Z} a_\ell = \hat{\beta}(x)bw \prod_{\ell \in Z \cup \mathcal{G}} a_\ell$$

if y is productive in U . If it is not productive, we use the same identity without b .

Subcase 0.2.5: ($\text{tail}(S)$ is T5): Either $\text{body}(S) = \emptyset$ and $\text{tail}(S) = S$, or else $\text{body}(S) \neq \emptyset$. Set $B_S = \{S\}$. Let $\text{tail}(S) = \{x, y := x^\rightarrow\}$.

Subcase 0.2.5.1: ($\text{body}(S) = \emptyset$): Locally at S , $\begin{array}{|c|} \hline \bullet \quad \mathcal{F}! \\ \hline \oplus_{U \cup Z} \end{array} \mapsto \hat{\beta}(x) \cdot \begin{array}{|c|} \hline \mathcal{G} \quad \bullet \\ \hline \end{array}$. In U , x is not productive, while y is productive if and only if y^\rightarrow does not contain a label of the same family as \mathcal{F} . In T_1 , x is productive, but y is not. y is productive in U if and only if $(\mathcal{F}) \in \bar{y}$ in T_1 . If y is productive in U , then $\text{boxfactor}_U(y) = \text{virtualfactor}_{y \in T_1}(\mathcal{F}) := a$.

For every edge label $\ell \in Z$ in U , there is a unique $\ell_1 \in \mathcal{F} \cup Z$ in T_1 with $\text{family}(\ell_1) = \text{family}(\ell) - 1$. Furthermore $\text{edgefactor}_U(\ell) = \text{edgefactor}_{T_1}(\ell_1) := b_\ell$. Let ℓ_1^M be greatest label of Z in T_1 , and let $\text{edgefactor}_{T_1}(\ell_1^M) := w$. Then $\text{virtualfactor}_{y \in U}(\mathcal{G}) = -w$.

Let $\text{boxfactor}_{T_1}(x) := c$. Then $\text{virtualfactor}_{x \in U}(\mathcal{G}) = \hat{\beta}(x)c$. Since T_1 has one more \mathcal{G} than U , $d(U) = d(T_1) - 1$.

In this case (12.3) is equivalent to $\text{wt } U = \hat{\beta}(x) \text{wt } T_1$. This follows from

$$(-1)^{d(U)} \cdot \left(\prod_{\ell \in Z} b_\ell \right) \cdot (-w) \hat{\beta}(x)c \cdot a = \hat{\beta}(x) \cdot (-1)^{d(U)+1} \cdot w \left(\prod_{\ell \in Z} b_\ell \right) \cdot a \cdot c,$$

if y is productive in U . If it is not productive, we use the same identity without a .

Subcase 0.2.5.2: ($\text{body}(S) \neq \emptyset$): Locally at S , the swap looks like

$$\begin{array}{|c|c|} \hline \bullet & \mathcal{F}! \\ \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \end{array} \mapsto - \prod_{z \in \bullet} \hat{\beta}(z) \cdot \begin{array}{|c|c|} \hline \mathcal{G} & \bullet \\ \hline \mathcal{G} & \bullet \\ \hline \end{array}.$$

Let A be the set of boxes of S_1 containing \mathcal{G} . The productive boxes of S are $\{z^\perp : z \in A\}$, as well as perhaps y , which is productive if and only if y^\rightarrow contains a label of the same family as \mathcal{F} in U . The productive boxes of S_1 are A . For each $z \in A$, let $a_z := \text{boxfactor}_{T_1}(z)$. Then $\text{boxfactor}_U(z^\perp) = \hat{\beta}(z)a_z$. If y is productive in U , let $b := \text{boxfactor}_U(y)$. Observe y is productive in U if and only if $(\mathcal{F}) \in \bar{y}$ in T_1 . Furthermore if y is productive in U , then $\text{virtualfactor}_{y \in T_1}(\mathcal{F}) = b$.

For every edge label $\ell \in Z$ in U , there is a unique $\ell_1 \in \mathcal{F} \cup Z$ in T_1 with $\text{family}(\ell_1) = \text{family}(\ell) - 1$. Furthermore $\text{edgefactor}_U(\ell) = \text{edgefactor}_{T_1}(\ell_1) := c_\ell$. Let ℓ_1^M be greatest label of Z in T_1 , and let $\text{edgefactor}_{T_1}(\ell_1^M) := w$. Then $\text{virtualfactor}_{y \in U}(\mathcal{G}) = -w$.

In this case (12.3) is equivalent to $\text{wt } U = -\prod_{z \in A} \hat{\beta}(z) \text{wt } T_1$. This follows from

$$\left(\prod_{\ell \in Z} c_\ell \right) \cdot (-w) \cdot b \prod_{z \in A} (\hat{\beta}(z) a_z) = \left(-\prod_{z \in A} \hat{\beta}(z) \right) \cdot w \left(\prod_{\ell \in Z} c_\ell \right) \cdot b \cdot \prod_{z \in A} a_z.$$

Subcase 0.2.6: ($\text{tail}(S)$ is T6): This case is covered by Subcases 0.2.4.2 and 0.2.4.3.

Case 1: ($\text{head}(S)$ is H1): Set $B_S = \{S\}$. Let x be the unique box of S . Locally at S , the swap is $\begin{array}{|c|} \hline \bullet \\ \hline \mathcal{G} \\ \hline \end{array} \mapsto \beta(x) \cdot \begin{array}{|c|} \hline \mathcal{G} \\ \hline \end{array} + \gamma \cdot \begin{array}{|c|} \hline \bullet \\ \hline \end{array}$, where $\gamma := 0$ if x is in row i and $\gamma := 1$ otherwise.

Let $\text{boxfactor}_{T_1}(x) := a$. The box x is productive in U if and only if $\text{family}(\text{label}(x^\leftarrow)) = \text{family}(\mathcal{G})$, in which case $\text{boxfactor}_U(x) = a$. Observe $\text{edgefactor}_{x \in U}(\mathcal{G}) = 1 - \hat{\beta}(x)a$, $\text{edgefactor}_{\bar{x} \in T_2}(\mathcal{G}) = 1 - a$, and x is not productive in T_2 . Notice further that $\gamma = 0$ if and only if $a = 1$.

If x^\leftarrow is empty, then x^\leftarrow is not productive in any of U, T_1, T_2 . If x^\leftarrow is nonempty and $\text{label}(x^\leftarrow) < \mathcal{G}$, then x^\leftarrow is productive in all three tableaux, and $\text{boxfactor}_U(x^\leftarrow) = \text{boxfactor}_{T_1}(x^\leftarrow) = \text{boxfactor}_{T_2}(x^\leftarrow) := b$. If x^\leftarrow is nonempty and $\text{family}(\text{label}(x^\leftarrow)) = \text{family}(\mathcal{G})$, then x^\leftarrow is productive in T_2 , but not in U or T_1 . Moreover by (G.6), $\mathcal{G}^- \in x^\leftarrow$, so $\text{boxfactor}_{T_2}(x^\leftarrow) = \hat{\beta}(x)a$.

In this case (12.3) is equivalent to $\text{wt } U = \beta(x) \text{wt } T_1 + \gamma \text{wt } T_2$. Since $\gamma = 1$ whenever $\text{wt } T_2 \neq 0$, it suffices to show $\text{wt } U = \beta(x) \text{wt } T_1 + \text{wt } T_2$. If x^\leftarrow is nonempty and $\text{label}(x^\leftarrow) < \mathcal{G}$, this follows from

$$(1 - \hat{\beta}(x)a) \cdot b = \beta(x) \cdot ab + (1 - a) \cdot b.$$

If x^\leftarrow is empty, we use the same identity without b . If $\text{family}(\text{label}(x^\leftarrow)) = \text{family}(\mathcal{G})$, we use the identity

$$(1 - \hat{\beta}(x)a) \cdot a = \beta(x) \cdot a + (1 - a) \cdot \hat{\beta}(x)a.$$

Case 2: ($\text{head}(S)$ is H2): Set $B_S = \{S\}$. Let x be the unique box of S . Locally at S , $\begin{array}{|c|} \hline \bullet \\ \hline \mathcal{G} \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \bullet \\ \hline \end{array} + \beta(x) \cdot \begin{array}{|c|} \hline \mathcal{G} \\ \hline \end{array}$.

Let $\text{boxfactor}_{T_2}(x) := a$. Then $\text{virtualfactor}_{x \in U}(\mathcal{G}) := a\hat{\beta}(x)$. In T_1 , $\mathcal{G} \in \bar{x}$ with $\text{virtualfactor}_{\bar{x} \in T_1}(\mathcal{G}) = a$. Due to T_2 's extra $\mathcal{G} \in x$, $d(T_2) = d(U) + 1 = d(T_1) + 1$.

In this case (12.3) is equivalent to $\text{wt } U = \text{wt } T_1 + \beta(x) \text{wt } T_2$. This follows from

$$(-1)^{d(U)} \cdot a\hat{\beta}(x) = (-1)^{d(U)} \cdot a + \beta(x) \cdot (-1)^{d(U)+1} \cdot a.$$

Case 3: ($\text{head}(S)$ is H3): Here $S = \{x\}$. Let $y = x^\leftarrow$. Locally at S , $\begin{array}{|c|} \hline \bullet \\ \hline i_k \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \bullet \\ \hline i_{k+1} \\ \hline \end{array}$.

Subcase 3.1: ($i_{k+1} \in x^\leftarrow = y^\rightarrow$, $i_k \in y$, no \bullet West of y in the same row): In U the y is not productive, whereas in T_1 , y is productive. Let $a := \text{boxfactor}_{T_1}(y)$.

Subcase 3.1.1: (y contains the only i_k in T): Let $\bar{T} := \phi_2^{-1}(T)$. Let S' be the snake in T containing y , \bar{S} be the snake in \bar{T} containing x and \bar{S}' the snake in \bar{T} containing y . Set $B_S = \{S, S', \bar{S}, \bar{S}'\}$. Locally at $S \cup S'$ and at $\bar{S} \cup \bar{S}'$ the swaps are respectively,

$$\begin{array}{|c|c|} \hline & \bullet \\ \hline i_k & i_{k+1} \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline & \bullet \\ \hline i_k & i_{k+1} \\ \hline \end{array} \quad \text{and} \quad \begin{array}{|c|c|} \hline & \bullet \\ \hline \bullet & i_{k+1} \\ \hline \end{array} \mapsto \beta(y) \begin{array}{|c|c|} \hline & \bullet \\ \hline i_k & i_{k+1} \\ \hline \end{array},$$

where $\bullet = \bullet_{i_k}$ before the swap and $\bullet = \bullet_{i_{k+1}}$ after the swap.

By Proposition 12.12, $[T]P_{i_k} = -[\overline{T}]P_{i_k}$. Hence, (12.3) is equivalent to

Claim C.1. $\text{wt}(U - \overline{U}) = \text{wt}(T_1 - \beta(y)T_1)$.

Proof. First, x is not productive in U, \overline{U} or T_1 . Second, x^\downarrow is productive in U, \overline{U} and T_1 and moreover

$$\text{boxfactor}_U(x^\downarrow) = \text{boxfactor}_{\overline{U}}(x^\downarrow) = \text{boxfactor}_{T_1}(x^\downarrow) := b.$$

Third, y is not productive in \overline{U} .

Next, $\text{edgefactor}_{y \in \overline{U}}(i_k) = 1 - a\hat{\beta}(y)$. Hence the claim follows from

$$b - (1 - a\hat{\beta}(y)) \cdot b = \hat{\beta}(y) \cdot ab = (1 - \beta(y))\text{wt}(T_1). \quad \square$$

Subcase 3.1.2: (Subcase 3.1.1 does not apply): Let $\overline{T} := \phi_1^{-1}(T)$. Let S' be the snake in T containing y , \overline{S} be the snake in \overline{T} containing x and \overline{S}' the snake in \overline{T} containing y . Set $B_S = \{S, S', \overline{S}, \overline{S}'\}$. Locally at $S \cup S'$ and at $\overline{S} \cup \overline{S}'$ the swaps are respectively,

$$\begin{array}{|c|c|} \hline & \bullet \\ \hline i_k & i_{k+1} \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline & \bullet \\ \hline i_k & i_{k+1} \\ \hline \end{array} \quad \text{and} \quad \begin{array}{|c|c|} \hline & \bullet \\ \hline \bullet & i_{k+1} \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline & \bullet \\ \hline \bullet & i_{k+1} \\ \hline \end{array} + \beta(y) \begin{array}{|c|c|} \hline & \bullet \\ \hline i_k & i_{k+1} \\ \hline \end{array},$$

where $\bullet = \bullet_{i_k}$ before the swap and $\bullet = \bullet_{i_{k+1}}$ after the swap. Note that $T_1 = \overline{T}_2$.

By Proposition 12.8, $[T]P_{i_k} = -[\overline{T}]P_{i_k}$. Thus (12.3) is equivalent to

Claim C.2. $\text{wt } U - \text{wt } \overline{U} = \text{wt } T_1 - \text{wt } \overline{T}_1 - \beta(y) \text{wt } T_1$.

Proof. Firstly, x is not productive in U, \overline{U}, T_1 or \overline{T}_1 . Secondly, x^\downarrow is productive in U, \overline{U}, T_1 and \overline{T}_1 . Moreover,

$$\text{boxfactor}_U(x^\downarrow) = \text{boxfactor}_{\overline{U}}(x^\downarrow) = \text{boxfactor}_{T_1}(x^\downarrow) = \text{boxfactor}_{\overline{T}_1}(x^\downarrow) := b.$$

Note y is not productive in U, \overline{U} or \overline{T}_1 . Observe $\text{virtualfactor}_{y \in \overline{U}}(\bullet_{i_k}) = a\hat{\beta}(y)$. Finally, $d(U) = d(T_1) = d(\overline{U}) + 1 = d(\overline{T}_1) + 1$. The claim then follows from

$$(-1)^{d(U)} \cdot b - (-1)^{d(U)-1} \cdot a\hat{\beta}(y) \cdot b = (-1)^{d(U)} \cdot ab - (-1)^{d(U)-1} \cdot b - \beta(y) \cdot (-1)^{d(U)} \cdot ab. \quad \square$$

Subcase 3.2: (Subcase 3.1 does not apply):

Subcase 3.2.1: (x^\downarrow is part of a T3 tail): Let S' be the snake containing x^\downarrow , and let $B_S = B_{S'}$. The remaining discussion of this case is found with the discussion of S' ; see e.g., Subcase 0.2.3.

Subcase 3.2.2: (Subcase 3.2.1 does not apply): Set $B_S = \{S\}$. Recall that locally at S , $\begin{array}{|c|} \hline \bullet_{i_k} \\ \hline \end{array} \mapsto \begin{array}{|c|} \hline \bullet_{i_{k+1}} \\ \hline \end{array}$. The swap affects no weight factors.

Case 4: (head(S) is H4): We argue the case $S = \text{head}(S)$. When S is a multirow, weight preservation follows by combining the present argument with that of Case 0.2.

Assume $S = \{x, x^\rightarrow\}$. Let $\overline{T} = \phi_{3, \{x, x^\rightarrow\}}(T)$, and let \overline{S} be the snake containing x in \overline{T} . Set $B_S = \{S, \overline{S}\}$. Locally at S, \overline{S} respectively the swaps are

$$\begin{array}{|c|c|} \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \bullet \\ \hline \end{array} \mapsto 0 \quad \text{and} \quad \begin{array}{|c|c|} \hline \bullet & \mathcal{G} \\ \hline \bullet & \mathcal{G} \\ \hline \end{array} \mapsto \hat{\beta}(x) \begin{array}{|c|c|} \hline \bullet & \mathcal{G} \\ \hline \mathcal{G} & \bullet \\ \hline \end{array}.$$

Since by Proposition 12.17, $[T]P_{\mathcal{G}} = [\overline{T}]P_{\mathcal{G}}$, (12.3) is equivalent to:

Claim C.3. $\text{wt } U + \text{wt } \overline{U} = \text{wt } \overline{T}_1$.

Proof. In U and \bar{U} , x is not productive. However x is productive in \bar{T}_1 . Let $\text{boxfactor}_{\bar{T}_1}(x) := a$. In \bar{T}_1 , x^\rightarrow is not productive. In U , x^\rightarrow is productive if and only if x^\rightarrow does not contain a label of family i . Further, x^\rightarrow is productive in U if and only if it is productive in \bar{U} and if and only if $(\mathcal{G}) \in \bar{x}^\rightarrow$ in \bar{T}_1 . In this case, $\text{boxfactor}_U(x^\rightarrow) = \text{boxfactor}_{\bar{U}}(x^\rightarrow) = \text{virtualfactor}_{\bar{x}^\rightarrow \in \bar{T}_1}((\mathcal{G})) := b$.

In T , $\text{edgefactor}(x) = 1 - \hat{\beta}(x)a$. Finally $d(U) = d(\bar{U}) + 1 = d(\bar{T}_1) + 1$. If x^\rightarrow is productive in U , then the claim follows from

$$(-1)^{d(U)} \cdot (1 - \hat{\beta}(x)a) \cdot b + (-1)^{d(U)-1} \cdot b = \hat{\beta}(x)(-1)^{d(U)-1} \cdot ab.$$

Otherwise we use the same identity with b 's removed. \square

Case 5: ($\text{head}(S)$ is H5): We only explicitly argue the case $S = \text{head}(S)$. The case S is a multirow ribbon follows by combining the present arguments with those from Case 0.2. Hence $S = \text{head}(S) = \{x, y := x^\rightarrow\}$.

Subcase 5.1: ($\text{head}(S)$ is H5.1): Let $\mathcal{B}_S = \{S\}$. Locally at S the swap is $\boxed{\bullet \begin{array}{|c|} \mathcal{G} \\ \mathcal{H} \end{array}} \mapsto \boxed{\bullet \begin{array}{|c|} \mathcal{G}' \\ \mathcal{H} \end{array}}$, where $\mathcal{H} \in \underline{y}$, $\text{family}(\mathcal{H}) = \text{family}(\mathcal{G}) + 1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$. Since no labels move, (12.3) follows trivially.

Subcase 5.2: ($\text{head}(S)$ is H5.2): Locally at S the swap is $\boxed{\bullet \begin{array}{|c|} \mathcal{G} \\ \mathcal{H} \end{array}} \mapsto \boxed{\bullet \begin{array}{|c|} \mathcal{G}' \\ \mathcal{H} \end{array}} + \hat{\beta}(x) \cdot \boxed{\mathcal{G} \bullet}$. By Lemma 5.6, y contains the westmost instance of \mathcal{G} and that hence $(\mathcal{G}) \notin \underline{x}$.

Subcase 5.2.1: ($\mathcal{G}^- \in x^\leftarrow$ with $\text{family}(\mathcal{G}^-) = i$): Let $\mathcal{B}_S = \{S\}$. Now, y is productive in U if and only if it is productive in T_1 and if and only if $(\mathcal{G}) \in \bar{y}$ in T_2 . In this case,

$$\text{boxfactor}_U(y) = \text{boxfactor}_{T_1}(y) = \text{virtualfactor}_{\bar{y} \in T_2}((\mathcal{G})) := a.$$

Let $\text{virtualfactor}_{y \in U}(\mathcal{H}) := b$. The box x is productive only in T_2 , with $\text{boxfactor}_{T_2}(x) = b$. The box x^\leftarrow is productive in U and T_1 ; x^\leftarrow is not productive in T_2 . Furthermore, $\text{boxfactor}_U(x^\leftarrow) = \text{boxfactor}_{T_1}(x^\leftarrow) = \hat{\beta}(x)b$. Observe $\text{virtualfactor}_{y \in T_1}(\mathcal{H}) = b - 1$. In this case, (12.3) is equivalent to $\text{wt } U = \text{wt } T_1 + \hat{\beta}(x) \text{wt } T_2$. If y is productive in U , this follows from the identity

$$b \cdot a \hat{\beta}(x)b = (b - 1) \cdot a \hat{\beta}(x)b + \hat{\beta}(x) \cdot a \cdot b.$$

If y is not productive in U , we use the same identity without a .

Subcase 5.2.2: ($\mathcal{G}^- \notin x^\leftarrow$ or $\text{family}(\mathcal{G}^-) \neq i$): Observe $T \in \mathcal{S}'_3$. Let $\bar{T} = \phi_3^{-1}(T)$. Note that $(\mathcal{H}) \notin \underline{y}$ in \bar{T} . Let \bar{S} be the snake of \bar{T} containing x . Set $\mathcal{B}_S = \{S, \bar{S}\}$. The swap at S is illustrated at the start of Case 5.2 above. Locally at \bar{S} , the swap is $\boxed{\bullet \begin{array}{|c|} \mathcal{G} \end{array}} \mapsto 0$.

Observe that y is productive in U if and only if it is productive in \bar{U} and if and only if it is productive in T_1 ; y is not productive in T_2 . Also if y is productive in U , then $\text{boxfactor}_U(y) = \text{boxfactor}_{\bar{U}}(y) = \text{boxfactor}_{T_1}(y) := a$. There is $(\mathcal{G}) \in \bar{y}$ in T_2 if and only if y is productive in U . In this case, $\text{virtualfactor}_{\bar{y} \in T_2}((\mathcal{G})) = a$.

Let $b := \text{virtualfactor}_{y \in U}(\mathcal{H})$ and $1 - c := \text{edgefactor}_{x \in \bar{U}}(\mathcal{G})$. Consequently we have $\text{virtualfactor}_{y \in T_1}(\mathcal{H}) = b - 1$, while $\text{boxfactor}_{T_2}(x) = c/\hat{\beta}(x)$. Observe $d(U) =$

$d(T_1) = d(T_2) = d(\bar{U}) - 1$. By Proposition 12.17, $[T]P_{\mathcal{G}} = [\bar{T}]P_{\mathcal{G}}$, to (12.3) is equivalent to $\text{wt } U + \text{wt } \bar{U} = \text{wt } T_1 + \hat{\beta}(x) \text{wt } T_2$. If y is productive in U , this follows from

$$(-1)^{d(U)} \cdot b \cdot a + (-1)^{d(U)-1} \cdot (1-c) \cdot a = (-1)^{d(U)} \cdot (b-1) \cdot a + \hat{\beta}(x) \cdot (-1)^{d(U)} \cdot a \cdot \frac{c}{\hat{\beta}(x)}.$$

Otherwise it follows from the same identity without a 's.

Case 5.3: ($\text{head}(S)$ is H5.3):

Subcase 5.3.1: ($(\mathcal{G}) \in \underline{x}$): Set $B_S = \{S\}$. Let $S = \{x, y := x^{\rightarrow}\}$. Locally at S , the swap is $\boxed{\bullet \mid \mathcal{G}} \mapsto \hat{\beta}(x) \boxed{\mathcal{G} \mid \bullet}$. Let $a := \text{virtualfactor}_{\underline{x} \in U}(\mathcal{G})$. In U , x is not productive, while y is productive if and only if y^{\rightarrow} does not contain a label of family i . Further y is productive in U if and only if $(\mathcal{G}) \in \bar{y}$ in T_1 . If y is productive in U , then $\text{boxfactor}_U(y) = \text{virtualfactor}_{\bar{y} \in T_1}(\mathcal{G}) := b$. In T_1 , x is productive, but y is not; $\text{boxfactor}_{T_1}(x) = \frac{a}{\hat{\beta}(x)}$. Here (12.3) is equivalent to $\text{wt } U = \hat{\beta}(x) \text{wt } T_1$. If y is productive in U , this follows from $a \cdot b = \hat{\beta}(x) \cdot b \cdot \frac{a}{\hat{\beta}(x)}$. Otherwise we use the same identity without b .

Subcase 5.3.2: ($\mathcal{G}^- \in x^{\leftarrow}$ with $\text{family}(\mathcal{G}^-) = i$): By (G.12) and Lemma 5.10, no label of family i appears in x 's column. Hence the $\mathcal{G} \in y$ is the Westmost \mathcal{G} . In particular, $(\mathcal{G}) \notin \underline{x}$, so this case is disjoint from Subcase 5.3.1. Set $B_S = \{S\}$. Locally at S , the swap is $\boxed{\bullet \mid \mathcal{G}} \mapsto \hat{\beta}(x) \boxed{\mathcal{G} \mid \bullet}$.

Let $a := \text{boxfactor}_U(x^{\leftarrow})$. In U , x is not productive, while y is productive if and only if y^{\rightarrow} does not contain a label of family i if and only if $(\mathcal{G}) \in \bar{y}$ in T_1 . In this case $\text{boxfactor}_U(y) = \text{virtualfactor}_{\bar{y} \in T_1}(\mathcal{G}) := b$. In T_1 , x^{\leftarrow} and y are not productive, while x is productive and $\text{boxfactor}_{T_1}(x) = a/\hat{\beta}(x)$. Here (12.3) is equivalent to $\text{wt}(U) = \hat{\beta}(x) \text{wt}(T_1)$. If $y \in U$ is productive then this follows from $ab = \hat{\beta}(x) \cdot b \cdot a/\hat{\beta}(x)$; otherwise the same is true after removing the b 's.

Subcase 5.3.3: (Subcases 5.3.1 and 5.3.2 do not apply): There is no $\mathcal{G}^- \in x^{\leftarrow}$ because we are not in Subcase 5.3.2. Since we are not in Case 5.1 there is not $\mathcal{H} \in \underline{y}$ with $\text{family}(\mathcal{H}) = i+1$ and $N_{\mathcal{H}} = N_{\mathcal{G}}$. By the assumption that we are in Case 5, there can be no label of $\text{family}(\mathcal{G})$ in the column of x . Thus if the $\mathcal{G} \in y$ were not Westmost then $(\mathcal{G}) \in \underline{x}$ and we would be in Subcase 5.3.1, a contradiction. Therefore we conclude $T \in S'_3$. Let $\bar{T} := \phi_3^{-1}(T)$. Let \bar{S} be the snake in \bar{T} containing x . Then set $\mathcal{B}_S = \{S, \bar{S}\}$.

Locally at S we have the swap is $\boxed{\bullet \mid \mathcal{G}} \mapsto \hat{\beta}(x) \boxed{\mathcal{G} \mid \bullet}$. Locally at \bar{S} , the swap is $\boxed{\bullet \mid \mathcal{G}} \mapsto 0$. By Proposition 12.17, $[T]P_k = [\bar{T}]P_k$. Therefore (12.3) is equivalent to $\text{wt}(U) + \text{wt}(\bar{U}) = \hat{\beta}(x) \text{wt}(T_1)$. This is exactly proved (up to renaming of tableaux) in Case 4.

Case 6: ($\text{head}(S)$ is H6): Here $S = \{x, x^{\rightarrow}\}$.

Subcase 6.1: ($\bullet_{\mathcal{G}} \notin x^{\rightarrow\uparrow}$): Let $\mathcal{B}_S = \{S\}$. Locally at S , $\boxed{\bullet \mid \mathcal{G}^+} \mapsto \beta(x) \cdot \boxed{\mathcal{G} \mid \mathcal{G}^+} + \hat{\beta}(x) \cdot \boxed{\mathcal{G} \mid \bullet_{\mathcal{G}^+}}$.

In U, T_1 and T_2 , x is not productive. Now, x^{\rightarrow} is productive in T_1 if and only if it is productive in T_2 if and only if it is productive in U ; in case of productivity,

$$a := \text{boxfactor}_U(x^{\rightarrow}) = \text{boxfactor}_{T_1}(x^{\rightarrow}) = \text{boxfactor}_{T_2}(x^{\rightarrow}).$$

Next, let $\text{edgefactor}_{\underline{x} \in U}(\mathcal{G}) := 1 - b$. Thus $\text{edgefactor}_{\underline{x} \rightarrow \in T_2}(\mathcal{G}^+) = 1 - b/\hat{\beta}(\underline{x})$.

Finally, (12.3) is equivalent to $\text{wt}(U) = \beta(\underline{x})\text{wt}(T_1) + \hat{\beta}(\underline{x})\text{wt}(T_2)$. If $\underline{x} \rightarrow$ is productive in U , then this follows from

$$(1 - b) \cdot a = \beta(\underline{x}) \cdot a + \hat{\beta}(\underline{x}) \cdot (1 - b/\hat{\beta}(\underline{x})) \cdot a;$$

otherwise we are done by the same expression without the a 's.

Subcase 6.2: ($\bullet_{\mathcal{G}} \in \underline{x} \rightarrow \uparrow$): Thus $T \in \mathcal{S}_2$. Let $\bar{T} := \phi_2(T)$. Let \bar{S} be the snake of \bar{T} containing \underline{x} . Let S' and \bar{S}' be the snakes of T and \bar{T} containing $\underline{x} \rightarrow \uparrow$, respectively. Set $\mathcal{B}_S = \{S, S', \bar{S}, \bar{S}'\}$.

Notice S' falls into Case 3 and in fact the $\mathcal{B}_{S'}$ defined there equals the current \mathcal{B}_S . Hence (12.3) holds by Case 3.

Case 7: ($\text{head}(S)$ is H7): Here $S = \{\underline{x}, \underline{x} \rightarrow\}$.

Subcase 7.1: ($\bullet_{\mathcal{G}} \notin \underline{x} \rightarrow \uparrow$): Let $\mathcal{B}_S = \{S\}$. Locally at S ,

$$\boxed{\begin{array}{c} \bullet \\ \mathcal{G} \end{array}} \boxed{\mathcal{G}^+} \mapsto \boxed{\bullet} \boxed{\mathcal{G}^+} + \beta(\underline{x}) \cdot \boxed{\mathcal{G}} \boxed{\mathcal{G}^+} + \hat{\beta}(\underline{x}) \cdot \boxed{\mathcal{G}} \boxed{\bullet_{\mathcal{G}^+}}$$

First, \underline{x} is not productive in U, T_1, T_2 or T_3 whereas $\underline{x} \rightarrow$ is productive in each of U, T_1, T_2 and T_3 or otherwise not productive in any of these tableaux. If $\underline{x} \rightarrow$ is productive then

$$a := \text{boxfactor}_U(\underline{x} \rightarrow) = \text{boxfactor}_{T_i}(\underline{x} \rightarrow) \text{ for } i = 1, 2, 3.$$

Second, let $b := \text{virtualfactor}_{\underline{x} \in U}(\mathcal{G})$. Third, $\text{edgefactor}_{\underline{x} \rightarrow \in T_3}(\mathcal{G}^+) = 1 - b/\hat{\beta}(\underline{x})$. Fourth, $d(T_1) = d(U)$, $d(T_2) = d(T_3) = d(U) + 1$.

Here (12.3) is equivalent to $\text{wt}(U) = \text{wt}(T_1) + \beta(\underline{x})\text{wt}(T_2) + \hat{\beta}(\underline{x})\text{wt}(T_3)$. If $\underline{x} \rightarrow$ is productive in U , then this follows from

$$(-1)^{d(U)} b \cdot a = (-1)^{d(U)} a + \beta(\underline{x}) (-1)^{d(U)+1} a + \hat{\beta}(\underline{x}) (-1)^{d(U)+1} \left(1 - \frac{b}{\hat{\beta}(\underline{x})}\right) \cdot a;$$

otherwise we are done by the same identity without the a 's.

Subcase 7.2: ($\bullet_{\mathcal{G}} \in \underline{x} \rightarrow \uparrow$): Thus $T \in \mathcal{S}_1$. Let $\bar{T} := \phi_1(T)$. Let \bar{S} be the snake of \bar{T} containing \underline{x} . Let S' and \bar{S}' be the snakes of T and \bar{T} containing $\underline{x} \rightarrow \uparrow$, respectively. Set $\mathcal{B}_S = \{S, S', \bar{S}, \bar{S}'\}$.

Notice S' falls into Case 3 and in fact the $\mathcal{B}_{S'}$ defined there equals the current \mathcal{B}_S . Hence (12.3) holds by Case 3.

Case 8: ($\text{head}(S)$ is H8): Here $\text{head}(S) = S$. The definitions of \mathcal{B}_S and subsequent analysis are exactly the same as in Case 3.

Case 9: ($\text{head}(S)$ is H9): Let $\mathcal{B}_S = \{S\}$. If $\text{head}(S) = S$, then since swapping at S does nothing (including no change to any \bullet indices), (12.3) is trivially true. If $\text{head}(S) \neq S$, then we use an argument similar to Subcase 0.2.

ACKNOWLEDGMENTS

AY thanks Hugh Thomas for his collaboration during [ThYo13]. Both authors thank him for his crucial role during the early stages of this project. OP was supported by an Illinois Distinguished Fellowship, an NSF Graduate Research Fellowship and NSF MCTP grant DMS 0838434. AY was supported by NSF grants and a Helen Corley Petit fellowship at UIUC.

REFERENCES

- [AnGrMi11] D. Anderson, S. Griffeth and E. Miller, *Positivity and Kleiman transversality in equivariant K-theory of homogeneous spaces*, J. Eur. Math. Soc., **13** (2011), 57–84.
- [Br05] M. Brion, *Lectures on the geometry of flag varieties*. Topics in cohomological studies of algebraic varieties, 33–85, Trends Math., Birkhäuser, Basel, 2005.
- [Bu02] A. Buch, *A Littlewood-Richardson rule for the K-theory of Grassmannians*, Acta Math. **189** (2002), 37–78.
- [Bu15] A. Buch, *Mutations of puzzles and equivariant cohomology of two-step flag varieties*, Ann. of Math. **182** (2015), 173–220.
- [BKPT13] A. Buch, A. Kresch, K. Purbhoo and H. Tamvakis, *The puzzle conjecture for the cohomology of two-step flag manifolds*, preprint, 2013. arXiv:1401.1725
- [BuKrTa03] A. Buch, A. Kresch and H. Tamvakis, *Gromov-Witten invariants on Grassmannians*, J. Amer. Math. Soc., **16** (2003), 901–915.
- [BuMi11] A. Buch and L. Mihalcea, *Quantum K-theory of Grassmannians*, Duke Math. J., **156** (2011), no. 3, 501–538.
- [Co09] I. Coşkun, *A Littlewood-Richardson rule for two-step flag varieties*, Invent. Math. **176** (2009), 325–395.
- [CoVa05] I. Coşkun and R. Vakil, *Geometric positivity in the cohomology of homogeneous spaces and generalized Schubert calculus*, in “Algebraic Geometry — Seattle 2005” Part 1, 77–124, Proc. Sympos. Pure Math., 80, Amer. Math. Soc., Providence, RI, 2009.
- [FuLa94] W. Fulton and A. Lascoux, *A Pieri formula in the Grothendieck ring of a flag bundle*, Duke Math. J., **76** (1994), no. 3, 711–729.
- [Gr01] W. Graham, *Positivity in equivariant Schubert calculus*, Duke Math. J., **109** (2001), no. 3, 599–614.
- [GrKu08] W. Graham and S. Kumar, *On positivity in T-equivariant K-theory of flag varieties*, Inter. Math. Res. Notices, Vol. 2008, Art. ID rnn093, 43 pages.
- [HLMvW11] J. Haglund, K. Luoto, S. Mason and S. van Willigenburg. *Refinements of the Littlewood-Richardson rule*, Trans. Amer. Math. Soc., **363** (2011), 1665–1686.
- [Kn10] A. Knutson, *Puzzles, positroid varieties, and equivariant K-theory of Grassmannians*, preprint, 2010. arXiv:1008.4302
- [KnMiYo09] A. Knutson, E. Miller and A. Yong, *Gröbner geometry of vertex decompositions and of flagged tableaux*, J. Reine Angew. Math. (Crelle’s J.) **630** (2009), 1–31.
- [KnTa03] A. Knutson and T. Tao, *Puzzles and (equivariant) cohomology of Grassmannians*, Duke Math. J. **119** (2003), no. 2, 221–260.
- [KoKu90] B. Kostant and S. Kumar, *T-equivariant K-theory of generalized flag varieties*, J. Differential Geom., **32** (1990), 549–603.
- [Kr05] V. Kreiman, *Schubert classes in the equivariant K-theory and equivariant cohomology of the Grassmannian*, preprint, 2005. arXiv:math/0512204
- [LaSc82] A. Lascoux and M.-P. Schützenberger, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris, **295** (1982), 629–633.
- [LePo07] C. Lenart and A. Postnikov, *Affine Weyl groups in K-theory and representation theory*, Int. Math. Res. Not., 2007, no. 12, Art. ID rnm038, 65 pp.
- [LiRi34] D. E. Littlewood and A. R. Richardson, *Group Characters and Algebra*, Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **233** (1934), no. 721–730, 99–141.
- [MoSa99] A. Molev and B. Sagan, *A Littlewood-Richardson rule for factorial Schur functions*, Trans. Amer. Math. Soc. **351** (1999), no. 11, 4429–4443.
- [PeYo15] O. Pechenik and A. Yong, *Genomic tableaux and combinatorial K-theory*, Discrete Math. Theor. Comput. Sci. Proc., to appear, 2015. FPSAC 2015, Daejeon, Korea.
- [RoYo13] C. Ross and A. Yong, *Combinatorial rules for three bases of polynomials*, Sémin. Lothar. Combin., to appear, 2014. arXiv:1302.0214
- [Sc77] M.-P. Schützenberger, *Combinatoire et représentation du groupe symétrique* (Actes Table Ronde CNRS, Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976), pp. 59–113. Lecture Notes in Math., Vol. 579, Springer, Berlin, 1977.
- [ThYo09a] H. Thomas and A. Yong, *A combinatorial rule for (co)minuscule Schubert calculus*, Adv. Math. **222** (2009), no. 2, 596–620.

- [ThYo09b] H. Thomas and A. Yong, *A jeu de taquin theory for increasing tableaux, with applications to K -theoretic Schubert calculus*, *Algebra Number Theory* **3** (2009), no. 2, 121–148.
- [ThYo13] H. Thomas and A. Yong, *Equivariant Schubert calculus and jeu de taquin*, *Ann. Inst. Fourier (Grenoble)*, to appear, 2013.
- [Wi06] M. Willems, *K -théorie équivariante des tours de Bott. Application à la structure multiplicative de la K -théorie équivariante des variétés de drapeaux*, *Duke Math. J.*, **132** (2006), no. 2, 271–309.
- [WoYo12] A. Woo and A. Yong, *A Gröbner basis for Kazhdan-Lusztig ideals*, *Amer. J. Math.*, **134**(2012), 1089–1137.

DEPT. OF MATHEMATICS, U. ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801, USA

E-mail address: pecheni2@illinois.edu, ayong@uiuc.edu